

List of Figures

		Page no.
Figure 4.1	XRD of synthesized nano crystalline zirconia samples	37
Figure 4.2	SEM of different samples of synthesized nano crystalline zirconia	39
Figure 4.3	TEM of nano zirconia samples 1A and 1B	41
Figure 4.4	TEM of nano zirconia samples 2A and 2B	42
Figure 4.5	TEM of nano zirconia samples 2A and 2B	43
Figure 4.6	DTA /TGA of Zr (OH) ₄ (0.075 M)	44
Figure 4.7	FTIR of synthesized nano zirconia samples	45
Figure 4.8	XRD diffractogram of nano crystalline iron oxide/hydroxide samples	47
Figure 4.9	FTIR of nano crystalline iron oxide/ hydroxide samples	48
Figure 4.10	XPS of nano crystalline iron oxide/ hydroxide samples	50
Figure 4.11	Magnetization curves of nano crystalline iron oxide/ hydroxide samples	51
Figure 4.12	Raman spectrums of synthesized nano crystalline iron oxide/ hydroxide samples	52
Figure 4.13	SEM images of iron oxide/ hydroxide	53
Figure 4.14	TEM images of iron oxide/ hydroxide	54
Figure 5.1	Effect of contact time on removal (%) of chromium from aqueous solution on nano crystalline zirconia (initial concentration = 10 mg L ⁻¹ , pH = 6.6, adsorbent dose = 2 g L ⁻¹ , agitation speed= 120 rpm temperature =303K)	56
Figure 5.2	A) Surface plot of 'chromium removal (%) vs. pH' and initial concentration mg L ⁻¹) at hold values of adsorbent dose and temperature at 4 g L ⁻¹ and 303K respectively B) Contour plot of 'chromium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold values of adsorbent dose and temperature at 4 g L ⁻¹ and 303K respectively C) Surface plot of 'chromium removal (%) vs. adsorbent dose and initial concentration (mg L ⁻¹)' at hold values of pH and temperature at 3 and 303K respectively D) Contour plot of 'chromium removal (%) vs. adsorbent dose and initial concentration (mg L ⁻¹)' at hold values of pH and temperature at 3 and 303K respectively	61
Figure 5.3	Surface plot of 'chromium removal (%) vs. pH and adsorbent dose' at hold values of initial concentration and temperature at 20 mg L ⁻¹ and 303K respectively B) Contour plot of 'chromium removal (%) vs. pH and dose' at hold values of initial concentration and temperature at 20 mg L ⁻¹ and 303K respectively	62

Figure 5.4	Optimization plots for removal of chromium from aqueous solution utilizing nano crystalline zirconia	63
Figure 5.5	Linear Langmuir isotherm plot of chromium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	65
Figure 5.6	Linear Freundlich isotherm plot of chromium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	65
Figure 5.7	Nonlinear Freundlich isotherm plot of chromium removal using nano crystalline zirconia obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	67
Figure 5.8	Nonlinear Langmuir isotherm plot of chromium removal using nano crystalline zirconia obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	67
Figure 5.9	Nonlinear Langmuir isotherm plot of chromium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	69
Figure 5.10	Nonlinear Freundlich isotherm plot of chromium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	69
Figure 5.11	Linear pseudo-first order plot of chromium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	70
Figure 5.12	Linear pseudo-second order plot of chromium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	71
Figure 5.13	Nonlinear pseudo-first order plot of chromium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	72
Figure 5.14	Nonlinear pseudo-second order plot of chromium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	73
Figure 5.15	Nonlinear pseudo-first order plot of chromium removal using nanocrystalline zirconia obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	73

Figure 5.16	Nonlinear pseudo-second order plot of chromium removal using nano crystalline zirconia obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	74
Figure 5.17	Intraparticle diffusion plot for removal of chromium from aqueous solution utilizing nano crystalline zirconia	76
Figure 5.18	Boyd model plot for removal of chromium from aqueous using nano crystalline zirconia	76
Figure 6.1	Effect of contact time on removal (%) of chromium from aqueous solution on nano crystalline iron oxide/hydroxide (initial concentration = 10 mg L ⁻¹ , pH = 6.6, adsorbent dose = 2 g L ⁻¹ , agitation speed= 120 rpm temperature = 303K)	82
Figure 6.2	(A) Effect of pH (adsorbent dose = 4 g L ⁻¹ , initial concentration = 20 mg L ⁻¹ , temperature = 303 K) (B) Effect of adsorbent dose (Initial pH=2, initial concentration = 20 mg L ⁻¹ , temperature = 303 K) (C) Effect of initial concentration on removal of chromium using nano iron oxide/hydroxide (adsorbent dose = 4 g L ⁻¹ , initial pH=2)	83
Figure 6.3	Surface plot of 'chromium removal (%) vs. adsorbent dose (g L ⁻¹) and initial concentration (mg L ⁻¹)' at hold values of pH and temperature at 2 and 293 K respectively	87
Figure 6.4	Surface plot of 'chromium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold values of adsorbent dose and temperature at 8 g L ⁻¹ and 313 K respectively	88
Figure 6.5	Surface plot of 'chromium removal (%) vs. pH and adsorbent dose (g L ⁻¹)' at hold values of initial concentration and temperature at 50 mg L ⁻¹ and 313 K respectively	88
Figure 6.6	Contour plot of 'chromium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold values of adsorbent dose and temperature at 8 g L ⁻¹ and 313 K respectively	90
Figure 6.7	Contour plot of 'chromium removal (%) vs. pH and adsorbent dose (g L ⁻¹)' at hold values of initial concentration and temperature at 50 mg L ⁻¹ and 313 K respectively	90
Figure 6.8	Contour plot of 'chromium removal (%) vs. adsorbent dose (g L ⁻¹) and initial concentration (mg L ⁻¹)' at hold values of pH and temperature at 2 and 293 K respectively	91
Figure 6.9	Optimisation plot of removal of chromium from aqueous solution by iron oxide/hydroxide	92
Figure 6.10	Linear Langmuir isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	93

Figure 6.11	Linear Freundlich isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	94
Figure 6.12	Nonlinear Langmuir isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	95
Figure 6.13	Nonlinear Freundlich isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	95
Figure 6.14	Nonlinear Langmuir isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	97
Figure 6.15	Nonlinear Freundlich isotherm plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	97
Figure 6.16	Linear pseudo-first order plot of chromium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	98
Figure 6.17	Linear pseudo-second order plot of chromium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	99
Figure 6.18	Nonlinear pseudo-first order plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	100
Figure 6.19	Nonlinear pseudo-second order plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	100
Figure 6.20	Nonlinear pseudo-first order plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	101
Figure 6.21	Nonlinear pseudo-second order plot of chromium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	102

Figure 6.22	Intraparticle diffusion plot for adsorption of chromium using nano iron oxide/hydroxide	103
Figure 6.23	Boyd plot of for adsorption of chromium using nanocrystalline iron oxide/hydroxide	104
Figure 7.1	Effect of adsorbent dose on removal (%) of cadmium from aqueous solutions on nano crystalline zirconia (Initial pH=7, Initial concentration 10 mg L ⁻¹ , Temperature 303K)	110
Figure 7.2	Effect of initial pH on removal (%) of cadmium from aqueous solutions on nano crystalline zirconia (Initial concentration= 10 mg L ⁻¹ , adsorbent dose 2 g L ⁻¹ Temperature= 303 K)	111
Figure 7.3	Effect of initial concentration on removal (%) of cadmium from aqueous solutions on nano crystalline zirconia (Initial pH= 7, adsorbent dose 2g L ⁻¹ , Temperature 303 K)	111
Figure 7.4	Effect of contact time on removal (%) of cadmium from aqueous solutions on nano crystalline zirconia (Initial concentration 1 mg L ⁻¹ , Initial pH =7, adsorbent dose 2 g L ⁻¹ , Temperature= 303 K)	112
Figure 7.5	Surface plot of 'cadmium removal (%) vs. pH and adsorbent dose (g L ⁻¹)' at hold value of initial concentration at 1mg L ⁻¹	116
Figure 7.6	Contour plot of 'cadmium removal (%) vs. pH and adsorbent dose (g L ⁻¹)' at hold value of initial concentration at 1 mg L ⁻¹	116
Figure 7.7	Surface plot of 'cadmium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold value of adsorbent dose at 8 g L ⁻¹	117
Figure 7.8	Contour plot of 'cadmium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold value of adsorbent dose at 8 g L ⁻¹	117
Figure 7.9	Surface plot of 'cadmium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold value of pH at 7	118
Figure 7.10	Contour plot of 'cadmium removal (%) vs. pH and initial concentration (mg L ⁻¹)' at hold value of pH at 7	118
Figure 7.11	Optimisation plot of removal of cadmium from aqueous solution by nano crystalline zirconia	119
Figure 7.12	Linear Langmuir isotherm plot of cadmium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	121
Figure 7.13	Linear Freundlich isotherm plot of cadmium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	121
Figure 7.14	Nonlinear Freundlich isotherm plot of cadmium removal using nano crystalline zirconia obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	123

Figure 7.15	Nonlinear Langmuir isotherm plot of cadmium removal using nano crystalline zirconia obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	123
Figure 7.16	Nonlinear Langmuir isotherm plot of cadmium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	125
Figure 7.17	Nonlinear Freundlich isotherm plot of cadmium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	125
Figure 7.18	Linear pseudo-first order plot of cadmium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	126
Figure 7.19	Linear pseudo-second order plot of cadmium removal using nano crystalline zirconia (dots represent the experimental data and lines represent the data estimated by the model)	126
Figure 7.20	Nonlinear pseudo-first order plot of cadmium removal using nano crystalline zirconia obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	128
Figure 7.21	Nonlinear pseudo-second order plot of cadmium removal using nano crystalline zirconia obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	128
Figure 7.22	Nonlinear pseudo-first order plot of cadmium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	129
Figure 7.23	Nonlinear pseudo-second order plot of cadmium removal using nano crystalline zirconia obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	130
Figure 7.24	Intraparticle diffusion plot for adsorption of cadmium using nanocrystalline zirconia	131
Figure 7.25	Boyd plot for adsorption of cadmium using nanocrystalline zirconia	131
Figure 8.1	Effect of initial concentration on removal (%) of cadmium from aqueous solution on nano crystalline iron oxide/hydroxide (Initial pH= 6, adsorbent dose 2 g L ⁻¹ , Temperature 303 K)	137
Figure 8.2	Effect of initial pH on removal (%) of cadmium from aqueous solution on nano crystalline iron oxide/hydroxide (Initial concentration= 20 mg L ⁻¹ , adsorbent dose 2 g L ⁻¹ Temperature=	138

	303K)	
Figure 8.3	Effect of adsorbent dose on removal (%) of cadmium from aqueous solution on nano crystalline iron oxide/hydroxide (Initial pH=6, Initial concentration 20 mg L ⁻¹ , Temperature 303K)	138
Figure 8.4	Effect of contact time on removal (%) of cadmium from aqueous solution on nano crystalline iron oxide/hydroxide (Initial concentration 20 mg L ⁻¹ , Initial pH =7, adsorbent dose 4g L ⁻¹ , Temperature= 303K)	139
Figure 8.5	Surface plot of 'cadmium removal (%) vs. pH and adsorbent dose' at hold values of initial concentration at 20 mg L ⁻¹	143
Figure 8.6	Contour plot of 'cadmium removal (%) vs. pH and adsorbent dose' at hold values of initial concentration at 20 mg L ⁻¹	144
Figure 8.7	Surface plot of 'cadmium removal (%) vs. adsorbent dose and initial concentration' at hold value of pH at 7	144
Figure 8.8	Contour plot of 'cadmium removal (%) vs. adsorbent dose and initial concentration' at hold value of pH at 7	145
Figure 8.9	Contour plot of 'cadmium removal (%) vs. pH and initial concentration' at hold values of adsorbent dose at 4 g L ⁻¹	145
Figure 8.10	Surface plot of 'cadmium removal (%) vs. pH and initial concentration' at hold values of adsorbent dose at 4 g L ⁻¹	146
Figure 8.11	Optimization plot of removal of cadmium with iron oxide/hydroxide	146
Figure 8.12	Linear Langmuir isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	148
Figure 8.13	Linear Freundlich isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	148
Figure 8.14	Nonlinear Freundlich isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	150
Figure 8.15	Nonlinear Langmuir isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by error analysis method (dots represent the experimental data and lines represent the data estimated by the model)	150
Figure 8.16	Nonlinear Langmuir isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	152
Figure 8.17	Nonlinear Freundlich isotherm plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by customized	152

	Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	
Figure 8.18	Linear pseudo-first order plot of cadmium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	153
Figure 8.19	Linear pseudo-second order plot of cadmium removal using nano crystalline iron oxide/hydroxide (dots represent the experimental data and lines represent the data estimated by the model)	153
Figure 8.20	Nonlinear pseudo-first order plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	154
Figure 8.21	Nonlinear pseudo-second order plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by error analysis function (dots represent the experimental data and lines represent the data estimated by the model)	155
Figure 8.22	Nonlinear pseudo-first order plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	157
Figure 8.23	Nonlinear pseudo-second order plot of cadmium removal using nano crystalline iron oxide/hydroxide obtained by customized Microcal origin function (dots represent the experimental data and lines represent the data estimated by the model)	157
Figure 8.24	Intraparticle diffusion plot for adsorption of cadmium using nanocrystalline iron oxide/hydroxide	158
Figure 8.25	Boyd plot for adsorption of cadmium using nanocrystalline iron oxide/ hydroxide	159