

# TABLE OF CONTENTS

---

---

<b>Titles</b>	<b>Page No.</b>
<b>Certificate</b>	<b>i</b>
<b>Declaration by the Candidate</b>	<b>ii</b>
<b>Copyright Transfer Certificate</b>	<b>iii</b>
<b>Acknowledgement</b>	<b>iv-v</b>
<b>Table of Contents</b>	<b>vi-x</b>
<b>List of Abbreviations</b>	<b>xi-xii</b>
<b>List of Figures</b>	<b>xiii-xiv</b>
<b>List of Tables</b>	<b>xv</b>
<b>Preface</b>	<b>xvii</b>
<b>Chapter 1: Introduction</b>	
<b>1 Introduction</b>	
1.1 Sleep wake cycle	
1.2 Common sleep deprivation consequences	
1.3 Difference between circadian rhythm and biological clock	
1.4 Different types of cycle	
1.4.1 Circadian rhythm	
1.4.2 Diurnal rhythm	
1.4.3 Ultradian rhythm	
1.4.4 Infradian rhythm/ Circalunar rhythm	
1.4.5 Circannual rhythm	
1.5 Circadian clock of photic and nonphotic cues	
1.6 Type of disrupted circadian rhythm disorder	
1.6.1 Delayed phase sleep wake syndrome (DSP)	
1.6.2 Early phase sleep wake syndrome (ASP)	
1.6.3 Jet lag disorder	
1.6.4 Shift work disorder	
1.6.5 Irregular sleep wake syndrome	
1.6.6 Free running (nonentrained)	
1.7 Transcription/Translation Feedback Loops (TTFL) of the Mammalian Clock	
1.8 Importance of light as a source of energy	
1.9 Circadian control of mitochondrial respiration	
1.10 Light source that we exposed in our daily life and their consequences	
1.11 Mitochondrial modulators	
1.11.1 N-Acetylcysteine (NAC)	
1.11.2 Melatonin	
1.12 Rationale	

- 1.13 Lacunae in the existing literature, relevance of the study
- 1.14 Hypothesis
- 1.15 Objectives

## **Chapter 2: Effect of normal and disrupted circadian rhythm on the mitochondria**

### **2 Introduction**

- 2.1 Materials and methods
  - 2.1.1 Animals and housing
  - 2.1.2 Experimental design
  - 2.1.3 Locomotor activity measurements
  - 2.1.4 Isolation of brain suprachiasmatic mitochondria
  - 2.1.5 Measurement of mitochondrial bioenergetics
  - 2.1.6 Measurement of mitochondrial DNA
  - 2.1.7 Quantification of clock genes expression
  - 2.1.8 Estimation of corticosterone
  - 2.1.9 Statistical analyses
- 2.2 Results
  - 2.2.1 Wheel running locomotors activity
  - 2.2.2 Respiratory control rate
  - 2.2.3 Mitochondrial DNA
  - 2.2.4 Clock genes
  - 2.2.5 Corticosterone
- 2.3 Discussion
- 2.4 Summary

## **Chapter 3: Evaluation of NAC effect on central clock related circadian rhythm**

### **3 Introduction**

- 3.1 Materials and Methods
  - 3.1.1 Animals
  - 3.1.2 Drugs
  - 3.1.3 Experimental Design
  - 3.1.4 Isolation of mitochondria from brain suprachiasmatic nuclei (SCN)
  - 3.1.5 Measurement of mitochondrial respiration
  - 3.1.6 Measurement of GSH fluorometrically
  - 3.1.7 Measurement of glutamate
  - 3.1.8 Measurement of total ATP content
  - 3.1.9 Measurement of plasma melatonin
  - 3.1.10 Estimation of corticosterone
  - 3.1.11 Quantification of clock genes
  - 3.1.12 Statistical analyses

### 3.2 Results

- 3.2.1 NAC improved the mitochondrial RCR and states in the SCN
- 3.2.2 NAC dose dependently improved the GSH and glutamate level in the SCN
- 3.2.3 NAC dose dependently improved total ATP content
- 3.2.4 NAC improved Melatonin
- 3.2.5 NAC decreases the CORT level in LL exposed mice
- 3.2.6 NAC dose dependently upregulated the Clock genes

### 3.3 Discussion

### 3.4 Summary

## **Chapter 4: Evaluation of N-acetylcysteine effect on peripheral clock related circadian rhythm**

### **4 Introduction**

#### 4.1 Materials and methods

- 4.1.1 Animals
- 4.1.2 Drugs
- 4.1.3 Experimental design
- 4.1.4 Determination of body weight and diurnal feed intake
- 4.1.5 Isolation of mitochondria from the liver
- 4.1.6 Measurement of mitochondrial respiration
- 4.1.7 Staining of succinate dehydrogenase
- 4.1.8 Measurement of GSH fluorometrically
- 4.1.9 Measurement of total ATP content
- 4.1.10 Measurement of leptin and ghrelin
- 4.1.11 Estimation of plasma glucose
- 4.1.12 Estimation of lipid profile
- 4.1.13 Statistical analyses

#### 4.2 Results

- 4.2.1 NAC recovered the body weight elevated by LL exposure
- 4.2.2 NAC restored the altered diurnal feed intake in LL exposed mice
- 4.2.3 NAC improved the mitochondrial RCR and states in the Liver
- 4.2.4 NAC improved Succinate dehydrogenase (SDH) activity in liver
- 4.2.5 NAC dose dependently improved the GSH in the liver
- 4.2.6 NAC dose dependently improved ATP level in the liver
- 4.2.7 NAC decreases the blood glucose level, total cholesterol, triglyceride and increases HDL in LL exposed mice
- 4.2.8 Leptin and ghrelin

#### 4.3 Discussion

#### 4.4 Summary

## **Chapter 5: Development of ex-vivo method for evaluation of mitochondrial modulators**

## **5. Introduction**

### 5.1 Materials and Methods

#### 5.1.1 Animals

#### 5.1.2 Chemicals

#### 5.1.3 Experimental design

#### 5.1.4 Determination of optimal DMSO concentration

#### 5.1.5 Isolation of mitochondria from the liver and brain

#### 5.1.6 Ex-vivo incubation and measurement of mitochondrial respiration

#### 5.1.7 Mitochondrial ETC enzyme activities

##### 5.1.7.1 NADH dehydrogenase (Complex-1) activity

##### 5.1.7.2 Succinate dehydrogenase (Complex-2) activity

##### 5.1.7.3 Cytochrome-C oxidase (Complex-4) activity

##### 5.1.7.4 F1F0 ATP synthase (Complex-5) activity

##### 5.1.7.5 Mitochondrial membrane potential (MMP)

#### 5.1.8 Statistical analysis

### 5.2 Results

#### 5.2.1 DMSO was toxic at >5% to isolated mitochondria

#### 5.2.2 MET inhibited mitochondrial respiration; Q10 improved MET inhibited mitochondrial respiration

#### 5.2.3 Q10 exerts a concentration-dependent effect on MET induced mitochondrial respiration

#### 5.2.4 Q10 exerts concentration-dependent effect on MET-induced mitochondrial dysfunction on complex-1, 5 with no effect on complex-2 and 4

#### 5.2.5 Q10 improved MET-induced changes in mitochondrial membrane potential (MMP)

### 5.3 Discussion

### 5.4 Summary

## **Chapter 6: Summary and conclusion**

### 6.1 Important outcomes

### 6.2 Limitation of the study

### 6.2 Future studies

## **References**

### **List of Publication from Thesis**