

LIST OF FIGURES

Fig. No.	Figure Title	Pg. No.
2.1.	<i>Curcuma longa</i> L. Plant	9
2.2.	Major phytochemical constituents from <i>Curcuma longa</i>	12
4.1	HPLC Chromatogram of A) <i>Curcuma longa</i> extract 1H sample B) <i>Curcuma longa</i> extract 2B sample	76
4.2	HPLC Chromatogram of A) <i>Curcuma longa</i> extract 3R sample B) <i>Curcuma longa</i> extract 4M sample	77
4.3	HPLC Chromatogram of A) pure curcumin B) turmeric oil	78
4.4	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight of stressed A) male mice B) female mice	79
4.5	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on basal rectal temperature of stressed A) male mice B) female mice	80
4.6	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on stress induced hyperthermia of stressed A) male mice B) female mice	81
4.7	Log dose response curve of A) male mice B) female mice	82
4.8	Effect of different <i>Curcuma longa</i> extracts (CLE) on A) body weight and B) basal rectal temperature of male mice	83
4.9	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on stress induced hyperthermia in male mice	84
4.10	Effect of different <i>Curcuma longa</i> extracts (CLE) on A) immobility time and B) onset and duration of sleep in stressed male mice	84
4.11	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and turmeric oil on A) body weight and B) basal rectal temperature of stressed male mice	85
4.12	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and turmeric oil on stressed induced hyperthermia of stressed male mice	86
4.13	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and turmeric oil on A) immobility time and B) onset and duration of stressed male mice	86
4.14	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) body weight and B) basal rectal temperature of stressed male mice	87
4.15	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) hot plate reaction time B) reaction time difference	87
4.16	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin treated stressed and non-stressed male rats in forced swim test	88

4.17	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight of male rats A) pooled data of all corresponding saline groups B) data from all groups	89
4.18	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on basal rectal temperature of male rats A) pooled data of corresponding saline group B) data from all groups	89
4.19	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) flinches per minute and B) paw edema in saline and formalin injected rats	90
4.20	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on food intake of male rats A) pooled data of corresponding NDW group and B) data from all groups	91
4.21	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on water intake of male rats A) pooled data of corresponding NDW group and B) data from all groups	91
4.22	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight of male rats A) pooled data of corresponding NDW group and B) data from all groups	92
4.23	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on plasma glucose, insulin, and corticosterone levels of A) normal drinking water and B) fructose water consuming male rats	92
4.24	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on total cholesterol, triglyceride, HDL and LDL levels of A) normal drinking water and B) fructose drinking water consuming male rats	93
4.25	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) body weight and B) basal rectal temperature of normal and stressed diabetic rats	94
4.26	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on stress induced hyperthermia of normal and stressed diabetic rats	95
4.27	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on spontaneous locomotor activity A) photo beam count for the first 60 sec and B) photo beam count for the last 60 sec of total 10 min period in actophotometer of normal and diabetic rats	95
4.28	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) numbers of entries and B) time spent in open and closed arms on elevated plus maze in normal and stressed diabetic rats	96
4.29	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on marble burying behavior in normal and diabetic rats	96
4.30	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on the weights of A) spleen B) adrenal glands and C) liver of normal and stressed diabetic rats	97

4.31	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) plasma glucose B) insulin and C) corticosterone level of normal and stressed diabetic rats	98
4.32	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on stomach ulceration index of A) non-stressed non diabetic B) stressed non diabetic C) stressed diabetic D) diabetic+ CLE-3R E) diabetic+ MET	99
4.33	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on levels of A) glutamate oxaloacetate transaminase (GOT) and B) glutamate pyruvate transaminase (GTP) in normal and stressed diabetic rats	99
4.34	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on A) blood and B) brain antioxidant status in normal and diabetic rats	100
4.35	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on Glyoxalase-I enzyme activity in normal and stressed diabetic rats	100
4.36	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on Paraoxonase 1 activity in normal and stressed diabetic rats	101
4.37	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on monoamines A) Noradrenaline B) Dopamine and C) 5-Hydroxytryptamine levels in brain samples of normal and stressed diabetic rats	101
4.38	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on monoamine oxidase (MAO) enzyme level A) MAO-A and B) MAO-B activity in brain samples of normal and diabetic rats	102
4.39	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on acetylcholinesterase (AChE) and butyrylcholinesterase (BChE) activity in A) blood B) brain cortex and C) brain hippocampus of normal and stressed diabetic rats	103
4.40	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on iNOS, NO and NF- κ B level in A) blood B) brain cortex and C) hippocampus of normal and diabetic rats	104

LIST OF TABLES

Table No.	Table Title	Pg. No.
2.1	Reported preclinical observations made with diverse types of <i>Curcuma longa</i> extracts	17
2.2	Some reported pharmacological activities of Curcumin	19
2.3	Marketed major formulations containing <i>Curcuma longa</i> extracts	21
2.4	Marketed major formulations containing curcumin	25
4.1	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 5	105
4.2	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 7	105
4.3	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of male mice on day 10	106
4.4	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 5	106
4.5	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 7	107
4.6	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on body weight difference of female mice on day 10	107
4.7	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on tail suspension test in male mice	108
4.8	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on tail suspension test in female mice	108
4.9	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on pentobarbital induced sleep test in male mice	109
4.10	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on pentobarbital induced sleep test in female mice	109
4.11	Effect of 95.4% curcuminoids containing <i>Curcuma longa</i> extract (CLE-3R) and metformin on glucose, insulin and corticosterone level in non-stressed and stressed male rats	110