

Table of contents

List of figures	xix
List of publications	xxxv
1 Introduction	1
1.1 Kinetics of phase transition	1
1.1.1 Liquid-liquid unmixing	3
1.2 Characterization of domain evolution and growth laws	7
1.2.1 Radial distribution function	8
1.2.2 Local number density distribution	9
1.2.3 Correlation function and the structure factor	9
1.2.4 Average domain length and growth laws	11
1.3 General examples of phase separating systems	16
1.3.1 Binary simple fluid	17
1.3.2 Binary polymer blend	19
1.3.3 Block copolymer melt	21
1.4 DPD simulation method	23
1.4.1 Modeling polymer chains	26
1.4.2 Updates in position and velocity	27
1.4.3 Model parameters	28

1.5	Overview of the thesis	29
1.6	Summary and discussion	31
2	Phase separation kinetics of BCP melts under on-off light cycles	33
2.1	Introduction	33
2.2	Photo-controlled reactions mechanism	36
2.3	System details	37
2.4	Effect of alternate light cycles	38
2.4.1	Reaction kinetics	41
2.4.2	Quantitative study of evolution	43
2.4.3	Characterization functions	45
2.4.4	Growth laws	48
2.5	Summary and discussion	50
3	Random photo-illumination on segregating polymeric fluids	53
3.1	Introduction	53
3.2	Modeling random photo-illumination	55
3.3	System details	56
3.4	Polymer blend system	57
3.4.1	Domain morphology and reaction kinetics	57
3.4.2	Radial distribution function	60
3.4.3	Characterization functions and anisotropy	61
3.4.4	Growth laws	64
3.5	BCP melt system	66
3.5.1	Domain morphology and reaction kinetics	67
3.5.2	Characterization functions	69
3.5.3	Growth laws	71

3.6	Summary and discussion	73
4	Block copolymer melts confined under moving parallel walls	75
4.1	Introduction	75
4.2	System details	78
4.3	Calculaion of velocity profile and shear-viscosity	79
4.4	Anisotropy parameter	80
4.5	Effect of external shear on viscosity and velocity profile	81
4.5.1	Velocity profile depends on different parameters	82
4.6	Average bond length and instantaneous temperature	86
4.7	Case 1: Both walls fixed	87
4.7.1	RDF and density distribution	88
4.7.2	Characterization functions	89
4.8	Case 2: Only top wall moves	91
4.8.1	RDF and density distribution	92
4.8.2	Characterization functions	94
4.9	Case 3: Both walls move in the same direction	96
4.9.1	RDF and density distribution	97
4.9.2	Characterization functions	98
4.9.3	Anisotropy developed in case 2 and case 3	100
4.10	Case 4: Both walls move in opposite directions	101
4.10.1	Characterization of domain coarsening	102
4.10.2	Anisotropy for different shear rates	105
4.11	BCP melt system with smaller chain length	108
4.12	Effect of shear on asymmetric BCP melt system	110
4.13	Summary and discussion	113

5	Segregation kinetics of binary polymer fluids: effect of external shear	116
5.1	Introduction	116
5.2	System details	119
5.3	Critical mixture of polymer blend	120
5.3.1	Case 1: Both walls fixed	123
5.3.2	Case 2: Only top wall moves	125
5.3.3	Case 3: Both walls move in the same direction	127
5.3.4	Case 4: Both walls move in opposite directions	128
5.4	Off-critical mixture of polymer blend	133
5.4.1	Case 1: Both walls fixed	133
5.4.2	Case 2: Only top wall moves	135
5.4.3	Case 3: Both walls move in the same direction	137
5.4.4	Case 4: Both walls move in opposite directions	138
5.5	Summary and discussion	142
6	Conclusion and future plan	144
6.1	Conclusion	144
6.2	Future Plan	146
	References	148

List of figures

- 1 Effect of on- and off-light cycles on the phase separation kinetics of BCP melt systems. xi
- 2 Random photo-illumination on polymer blend systems. xii
- 3 Effect of external shear on the phase separation kinetics of BCP melts. xiii
- 4 Impact of external shear on critical and off-critical polymer blend systems. xiv

- 1.1 Phase diagram for ferromagnet [1]. Here T and h represents the temperature and the magnetic field, respectively. The line ($T < T_c, h = 0$) denotes the first-order transition, and the point ($T_c, h_c = 0$) signifies the second-order critical point. When $T > T_c$, the system is in a disordered state, whereas for $T < T_c$, it falls in magnetized state with spins aligned either upwards or downwards. 3
- 1.2 Phase diagram for (AB) binary mixture [1]. ϕ depict the volume fraction and T represents the temperature. At the second-order critical point, $T/T_c = 1$, and $\phi = 0.5$. A solid line represents the coexistence curve. The system is in the homogeneously mixed state above the coexistence curve and starts the phase separating below it. The dashed line is a spinodal curve representing the metastable states between spinodal and coexistence curves. 6

1.3	(a) Schematic for calculating RDF at a distance r from the reference particle. (b) The general behavior of RDF representing the distribution of B -type particles around A in binary fluids.	8
1.4	Plots for the correlation function and corresponding structure factor curves for binary mixture are shown in (a) and (b), respectively. The black dashed line in (b) with a slope of -4 represents the Porod's tail in $3d$	11
1.5	Schematic represents the repeating units of simple fluids, polymer blends, and block copolymer melts.	16
1.6	(a-e) The evolution morphology of phase separating binary fluid at different time intervals, indicated alongside each image. (f) $g_{AB}(r)$ vs. r for different time intervals.	17
1.7	(a) Plots for $C(r, t)$ vs. $r/R(t)$. (b) The structure factor $S(k, t)$ vs. $kR(t)$ at various time intervals, as indicated in the legends. The solid black line represents the porod's tail as $k \rightarrow \infty$. Inset of (a) and (b) plot the unscaled correlation and structure factor. (c) $R(t)$ vs. t for a simple fluid. Solid lines, characterized by slopes of 1 and $2/3$, signify the phases of viscous and inertial hydrodynamic growth experienced by the system. . .	19
1.8	We plot the evolved morphology of the polymer blend system in (a-e) at different time intervals. (f) plots the corresponding RDF curves.	20
1.9	In (a) and (b), we depict the scaled $C(r, t)$ vs. $r/R(t)$ and $S(k, t)$ vs. $kR(t)$, respectively, for different time intervals as indicated in the legends. Inset of (a) and (b) plot the unscaled correlation and structure factor. (c) The time-dependent behavior of $R(t)$	21
1.10	In (a-e), snapshots of the domain morphology of the BCP melt system are shown at various time intervals. Corresponding RDF plots are depicted in (f).	22

-
- 1.11 In correspondence to the evolved morphology displayed in Fig. 1.10, we plot the unscaled and scaled correlation function in (a) and (b), respectively. (c) and (d) represent the unscaled and scaled structure factor. 23
- 1.12 Temporal variation of characteristic length scale $R(t)$ vs. t 24
- 2.1 (a) Schematic of bond-breaking and combination reactions during the on- and off-light cycles. P_b and P_c depict the probability of bond-breaking and recombination reactions. (b) The homogeneous mixture of BCP melt system (1 : 1 ratio of A and B -type beads: critical composition) after the equilibration. (c) Representation of color marks of the beads: Inactive A and B -type beads are represented by blue and yellow. At the same time, the corresponding active radicals are shown with glowing cyan and glowing orange. 37
- 2.2 (a) Homogeneous mixture of BCP ($A_nB_n; n = 16$) melt at $t = 0$. (b-f) Subsequent snapshots depict the evolution after each cycle in a sequence of off \rightarrow on \rightarrow off \rightarrow on \rightarrow off. The blue and yellow colors represent the A and B -type beads, respectively. 39
- 2.3 Evolution snapshots for set 1 after the first and second cycles at different bond-breaking probabilities: (a) $P_b = 0.1$, (b) $P_b = 0.5$, and (c) $P_b = 1.0$. The first column depicts the microphase-separated structures at $t = 600$. The second column shows the snapshots after cycle 2. The interface between A and B -type domains is illustrated in the third column at $t = 1200$. 40

- 2.4 Evolution snapshots for set 2 after the completion of cycle 1 (on) and cycle 2 (off) at different bond-breaking probabilities: (a) $P_b = 0.1$, (b) $P_b = 0.5$, and (c) $P_b = 1.0$. The first column shows the macrophase-separated evolution morphology at $t = 600$. The second column displays the snapshots after cycle two. The interface between A and B-type domains is illustrated in the third column, corresponding to snapshots shown in column two. 40
- 2.5 We normalize the cumulative number of bonds broken (N_b) during on-cycles and a cumulative number of bonds recombined (N_c) during off-cycles by the total number of photo-sensitive bonds N_{pc} at $t = 0$. Corresponding normalized numbers are denoted by n_b and n_c , respectively, and plotted as a function of time. (a) Plots for n_b vs. t during on-cycles and n_c vs. t during off-cycles of set 1. (b) Variation of n_b with time for different P_b values for cycle 2 in set 1. (c) Time variation of n_b and n_c corresponding to different cycles of set 2 is plotted with different symbol types. The dashed lines represent the exponential fits, $n_b(t) \sim (1 - e^{-\lambda P_b t})$ followed by each plot, where λ is the fitting parameter. 42
- 2.6 (a) Plots for radial distribution function $g_{AB}(r)$ vs. r after each cycle of set 1. (b) Local number density distribution of A-type beads along z -direction for the same data set plotted in (a) with various symbol types. For each plot, the probability P_b and P_c are fixed at 1.0. 43
- 2.7 Comparison of $g_{AB}(r)$ vs. r for different probabilities in cycle 2 (on-cycle) of set 1. The inset plot shows the RDF for cycle 4 (on-cycle) with different P_b values indicated by different symbols as specified in the legends. . . . 44

- 2.8 (a) and (b) plots the spherically averaged scaled correlation function and corresponding structure factor after the completion of different on- and off-light cycles in set 1. Various symbol types represent the data at different time intervals. The solid line in (a) depicts the zero crossing of the correlation function. The tail of the structure factor follows Porod's law, illustrated with a solid black line in (b) with slope -4 . (c-d) Scaled correlation function $C(r, t)$ vs. $r/R(t)$ and the structure factor $S(k, t)$ vs. $kR(t)$ for different time intervals in cycle 2 set 1 (light-on). (e-f) Same set of observables plotted in (c) and (d) for cycle 3 set 1 (light-off). 46
- 2.9 (a-b) Scaled correlation and structure factor after ending each light cycle for set 2. Various symbol types represent the different time intervals. (c-d) Plots for $C(r, t)$ vs. $r/R(t)$ and $S(k, t)$ vs. $kR(t)$ for the evolution through cycle 1 set 2 (bond-breaking on). (e-f) Same set of observables as in (c) and (d) for the evolution through cycle 2 set 2 (bonds are recombining). 47
- 2.10 (a) Average domain size $R(t)$ vs. t is plotted for microscopic domain growth during off-cycles (1, 3, and 5) of set 1 when bond recombination reactions reform the BCP chains. A solid black line represents the expected diffusive growth ($\phi \sim 1/3$) at early times. (b) Plots for average length scale $R(t)$ vs. t during macroscopic domain growth in cycles 2 and 4, when bond-breaking reactions turn the BCP melt system into a binary polymer blend. The solid line displays the inertial hydrodynamic growth ($\phi \sim 2/3$) for $d = 3$ fluids. 49
- 2.11 Logarithmic plots for the average domain growth $R(t)$ as a function of time. Data corresponding to on-cycles (1, 3, and 5) of set 2 are plotted in (a) while (b) plots the data for off-cycles (2 and 4). Solid lines represent the growth laws followed by the system in different cycles. 50

-
- 3.1 Schematic of bond-breaking and combination reactions under random photo-illumination. The inactive A and B -type beads are represented by blue and yellow. At the same time, the corresponding active radicals are shown with glowing cyan and glowing orange. 56
- 3.2 (a-d) Domain evolution snapshots of polymer blend system for different bond breaking probabilities at $t = 3.0 \times 10^3$. Blue and yellow colors represent A and B beads, respectively. (e) Time variation of the normalized cumulative number of bonds broken (n_b) at various P_b values. (f) n_b vs. P_b at two different time intervals $t = 2.0 \times 10^2$ and 3.0×10^3 plotted with black and red curves, respectively. Inset plots the n_b vs P_b on log-log scale at higher P_b values. (g) Scaling of the normalized cumulative number of bonds broken at low P_b values ($0.01 - 0.1$). Corresponding scaled data is denoted as n_{bs} . The scaling at higher P_b values ($0.2 - 1.0$) is shown in (h). Legends at the top represent the different P_b values depicted with various symbol types. 58
- 3.3 (a) Radial distribution function, $g_{AB}(r)$, at $P_b = 0.5$. Different symbol types in the legends represent different time intervals. (b) Comparison of $g_{AB}(r)$ vs, r at different P_b values at a late time $t = 3.0 \times 10^3$ 60
- 3.4 Spherically averaged scaled correlation function $C(r, t)$ vs. $r/R(t)$ at different time intervals are plotted with different symbols at $P_b = 0.01$ in (a) and $P_b = 0.5$ in (b). The solid black line represents the zero-crossing. 61

- 3.5 (a-b) Scaled correlation function $C(r, t)$ vs. $r/R(t)$ for different bond-breaking probability $P_b = 0.01, 0.1, 0.5,$ and 1.0 plotted with black, red, green, and blue curves respectively for $t = 1.2 \times 10^3$ in (a), and $t = 3.0 \times 10^3$ in (b). Corresponding evolved morphologies are displayed in Figs. 3.2(a)-3.2(d). (c-d) Plots for the structure factor along $x, y,$ and z direction at $t = 3.0 \times 10^3$ are displayed in (c) for $P_b = 0.01$ and (d) for $P_b = 0.5$ to discuss the anisotropy developed within the system. 63
- 3.6 (a) The characteristic length scale $R(t)$ vs. t for different bond-breaking probabilities $P_b \in (0.0, 1.0)$ at a fixed bond recombination probability $P_c = 1.0$. The solid lines represent the growth laws followed by the system ($1/3$ for the diffusive growth and $2/3$ for the inertial hydrodynamic growth). For a reference case, the length scale of pure BCP melt and pure polymer blend is plotted with the black and magenta curves. (b) Plots for $R(t)$ vs. P_b at different time intervals are depicted with various symbol types in the legends. (c) ϕ_{eff} vs. $1/t$ for different bond-breaking probabilities plotted with various symbols given in the legends. 65
- 3.7 (a-d) Domain evolution of BCP melt system at different light intensities: ($P_b = 0.01, 0.1, 0.5,$ and 1.0) at $t = 1.2 \times 10^4$. (e) Time-dependent behavior of n_b at different bond-breaking probabilities. (f) n_b vs. P_b at two different time intervals: $t = 1.2 \times 10^3$ (black curve) and $t = 1.2 \times 10^4$ (red curve). Inset plots the data at high P_b values representing the power law growth. (g) and (h) plot the n_b vs. t data after scaling with the power law equation with the unit and lower growth exponents, respectively. 67

- 3.8 (a-b) Comparison of the scaled correlation function and structure factor for $P_b = 0.01$ at various time intervals, depicted by different symbols. In (a), $C(r, t)$ vs. $r/R(t)$ curves are plotted on a linear scale, while (b) shows $S(k, t)$ vs. $kR(t)$ on a logarithmic scale. (c-d) The same quantities are plotted for $P_b = 0.5$ 69
- 3.9 (a-b) Comparison of the correlation and the structure factor curves for different bond-breaking probabilities P_b : 0.01 (black), 0.1 (red), 0.5 (green), and 1.0 (blue), at a late-time regime, $t = 1.2 \times 10^4$ 70
- 3.10 (a) The characteristic domain size, $R(t)$ vs. t on a logarithmic scale for various P_b values. The length scale for a pure BCP melt system is plotted with a black dashed curve for a reference case. The inset of (a) represents the plots for $R(t)$ vs. P_b at different time intervals indicated by various symbol types in the legends. (b) The effective growth exponent, ϕ_{eff} is plotted as a function of $1/t$ corresponding to the length scale presented in (a). 72
- 4.1 (a) Plots of the average velocity profile, $\langle v_x \rangle$ vs. z , at a high shear rate ($\dot{\gamma} = 1.56 \times 10^{-2}$), with different symbols representing the various scenarios. (b) Plots of shear viscosity, $\langle \eta \rangle$, vs. $\dot{\gamma}$ at the late time step ($t = 5 \times 10^3$) for all three scenarios, each indicated by distinct symbols. 81

- 4.2 (a-c) Plots for the velocity profile, $\langle v_x \rangle$ along z -direction. (a) $\langle v_x \rangle$ vs. z at different shear rates mentioned in the legend at a fixed chain length, $N_p = 32$. (b) Effect of BCP chain length on $\langle v_x \rangle$ vs. z at a fixed shear rate, $\dot{\gamma} = 1.56 \times 10^{-2}$. The three black, red, and green curves represent the data for chain length $N_p = 8, 16,$ and 32 , respectively. In (a) and (b), The height of the box is fixed at $L = 64$ in a cubic simulation box. We vary the height of the box in (c) and compare the velocity profile by keeping the box length ($L = 64$) fixed in x - and y -directions at $\dot{\gamma} = 1.56 \times 10^{-2}$ and $N_p = 32$. In (d-f), the velocity profiles are plotted for case 3 with the variation of the same set of parameters as discussed in (a-c). 83
- 4.3 Plots for average velocity profile against time for all three cases of applied shear are mentioned at the top. (a-c) Plots for $\langle v_x \rangle$ vs. t for different shear rates mentioned in the legends. Data is taken at a plane near the top wall. In (d-f) $\langle v_x \rangle$ is plotted at different xy -planes for a moderate shear rate $\dot{\gamma} = 7.8 \times 10^{-3}$. (g-i) plots the same data set as in (d-f), for high shear rate $\dot{\gamma} = 1.56 \times 10^{-2}$ 85
- 4.4 (a), (c), and (e) represent the time-dependent average bond length $\langle l_b \rangle$ for cases 2-4 mentioned at extreme right. Various symbols in the legends represent the different shear rates applied to the system. In the second column (b), (d), and (f), we plotted the instantaneous temperature over time for the same cases as in the first column. 86
- 4.5 Comparison of late stage morphology of phase separating critical BCP melt for (a) $N_p = 8$, (b) $N_p = 16$, and (c) $N_p = 32$. The second right frame shows the corresponding isosurfaces. 87

- 4.6 (a) Plots for the radial distribution function $g_{AB}(r)$ vs. r for BCP chain length $N_p = 32$ for case 1. Different symbols types mentioned in the legends represent the various time intervals. (b) Average number density distribution function $\rho_A(y)$ vs. y of A -type beads, corresponding to the same time intervals as in (a). 89
- 4.7 (a-b) Spherically averaged scaled correlation $C(r, t)$ vs. $r/R(t)$ and corresponding structure factor $S(k, t)$ vs. $kR(t)$ are plotted for different time intervals with various symbols mentioned in the legends. The solid black line in (a) represents the zero crossing of correlation function. The structure factor curves follow Porod's law: $S(k, t) \sim k^{-4}$ at $k \rightarrow \infty$ represented by a solid black line with slope -4 . (c) The characteristic length scale over time for case 1. The solid black line with slope $1/3$ represents the diffusive growth at early times which saturates later. (d) Plots for the effective growth exponent ϕ_{eff} vs. $1/R(t)$ correspond to the length scale plotted in (c). The solid black line illustrates the reference value of $\phi_{eff} \sim 1/3$ for early diffusive growth. 90
- 4.8 Morphology evolution of BCP melts system for case 2 for three different shear rates (a) $\dot{\gamma} = 1.56 \times 10^{-3}$, (b) $\dot{\gamma} = 7.80 \times 10^{-3}$, and (c) $\dot{\gamma} = 1.56 \times 10^{-2}$. The snapshots are taken at two different time intervals: $t = 1.2 \times 10^2$ (first column) and $t = 5.4 \times 10^3$ (second column). The direction of the wall motion is shown with a magenta arrow. In the third column, the isosurfaces are plotted, corresponding to the evolution shown in the second column. 92

- 4.9 (a) and (b) represents the plots of $g_{AB}(r)$ vs. r and $\rho(y)$ of A -beads along y -direction, respectively for case 2. These data sets are computed at a fixed shear rate of $\dot{\gamma} = 7.80 \times 10^{-3}$ for different time intervals mentioned in the legends. (c) $g_{AB}(r)$ vs. r for different shear rates at a fixed time interval of $t = 5.4 \times 10^3$. (d) $\rho(y)$ versus y for the same data set as in (c). 93
- 4.10 The characterization functions are plotted at a fixed time step $t = 5.4 \times 10^3$ for different shear rates for case 2. The scaled correlation function $C(r, t)$ versus $r/R(t)$ is displayed in (a), While the corresponding structure factor $S(k, t)$ versus $kR(t)$ on a logarithmic scale is shown in (b). (c) The characteristic length scale is plotted against time for given shear rates. To calculate the growth exponent, we display the ϕ_{eff} as a function of $1/R(t)$ in (d). 95
- 4.11 The evolution snapshots for case 3, where both walls move in the same direction. Snapshots are plotted at two-time steps $t = 1.2 \times 10^2$ and $t = 5.4 \times 10^3$. The shear rates applied to the system is varied as: (a) $\dot{\gamma} = 1.56 \times 10^{-3}$, (b) $\dot{\gamma} = 7.80 \times 10^{-3}$, and (c) $\dot{\gamma} = 1.56 \times 10^{-2}$. The third column represents the isosurfaces corresponding to the snapshots in the second column. The magenta arrows display the shear direction. 97
- 4.12 Case 3: Both walls are moving in x -direction with similar velocities. (a) Plots the $g_{AB}(r)$ versus r , and (b) the average number density profiles of A -type beads in y -direction at $t = 5.4 \times 10^3$ for different shear rates; related morphologies are shown in Fig. (4.11) 98

- 4.13 (a) Spherically averaged $C(r, t)$ vs. $r/R(t)$ for different shear rates; related morphologies are illustrated in Fig. 4.11. The corresponding $S(k, t)$ versus $kR(t)$ curves are plotted in (b) on a log-log scale. (c) $R(t)$ versus t for different shear rates. (d) Plots for related effective growth exponents ϕ_{eff} versus $1/R(t)$ 99
- 4.14 Plots for the unidirectional structure factor $S(k_x, k_y, k_z)$ versus k_y (black curve) and k_z (red curve) to discuss the structural anisotropy in the system. The data is calculated at a fixed shear rate of $\dot{\gamma} = 7.80 \times 10^{-3}$ and plotted for case 2 in (a) while for case 3 in (b). The inset represents the $S(k_x, k_y, k_z)$ versus k_y along diagonal and cross-diagonal directions with green and blue curves, respectively, in yz -plane. 100
- 4.15 Case 4: The evolution snapshots at two different time steps $t = 1.2 \times 10^2$ and $t = 5.4 \times 10^3$ for different shear rates (a) $\dot{\gamma} = 1.56 \times 10^{-3}$, (b) $\dot{\gamma} = 7.80 \times 10^{-3}$, and (c) $\dot{\gamma} = 1.56 \times 10^{-2}$. To see the bulk structures, we have plotted the isosurfaces in column three corresponding to the snapshots at $t = 5.4 \times 10^3$ 102
- 4.16 Case 4: (a) Comparison of $g_{AB}(r)$ versus r , (b) $\rho(y)$ versus y of A -type beads, (c) $C(r, t)$ versus $r/R(t)$, and (d) $S(k, t)$ versus $kR(t)$ at $t = 5.4 \times 10^3$ for different shear rates mentioned in the legends. (e) The characteristic average domain size $R(t)$ vs. t for the evolution displayed in Fig. 4.15. (f) Plots for the effective growth exponents ϕ_{eff} vs. $1/R(t)$ for different shear rates. 103
- 4.17 We plot the unidirectional structure factor $S(k_x, k_y, k_z)$ versus k_y (black curve) and k_z (red curve) for different shear rates of $\dot{\gamma} = 0.0$ in (a), $\dot{\gamma} = 1.56 \times 10^{-3}$ in (b), $\dot{\gamma} = 7.80 \times 10^{-3}$ in (c), and $\dot{\gamma} = 1.56 \times 10^{-2}$ in (d) to analyze the structural anisotropy developed in the system. 106

- 4.18 Time variation of anisotropy parameters, D_{xy} , D_{yz} , and D_{xz} are plotted in (a-d). (a) represent the reference case 1 ($\dot{\gamma} = 0.0$). (b-d) plots the same data for case 2 for different shear rates, $\dot{\gamma} = 1.56 \times 10^{-3}$, 7.80×10^{-3} , and 1.56×10^{-2} , respectively. Time variations of D_{yz} are compared in (e) and (f) for case 3 and case 4 at different shear rates depicted by various symbol types. 107
- 4.19 Effect of shear on the BCP melt system with a smaller chain length of A_4B_4 . We plot the late time evolution, and related isosurfaces of BCP melt for all three scenarios 2-4 at a fixed moderate shear rate of $\dot{\gamma} = 7.80 \times 10^{-3}$ and time step $t = 5.4 \times 10^3$. Corresponding snapshots for $\dot{\gamma} = 0.0$ is shown in Fig. 4.5(a). 108
- 4.20 The evolution snapshots and related isosurfaces for a BCP melt system with a smaller chain length of A_8B_8 . The shear is applied with cases 2-4 with a moderate shear rate of $\dot{\gamma} = 7.80 \times 10^{-3}$ and time step $t = 5.4 \times 10^3$. Corresponding snapshots for $\dot{\gamma} = 0.0$ is shown in Fig. 4.5(b). 109
- 4.21 Domain evolution for asymmetric BCP melt system confined between two parallel rigid walls. (a) Evolution snapshot and related isosurface plot of BCP melt system with the ratio of 1:3 between A and B beads, at the time step $t = 5.4 \times 10^3$ for zero shear rate. (b) Snapshot at the same time and zero shear for the BCP melt system with a 1:7 ratio of A and B beads. 110
- 4.22 Comparison of the effect of shear in different cases 2-4 in BCP melt system with 1:3 ratio of A and B beads. The evolution snapshots are taken at two different time intervals plotted in the first two columns at $t = 1.2 \times 10^2$ and $t = 5.4 \times 10^3$, respectively. The third column plots the isosurface related to the snapshots at the late time. 111

- 4.23 Evolution snapshots of BCP melt system with 1:7 ratio of A and B beads for cases 2-4. The first two columns represent the morphology at $t = 1.2 \times 10^2$ and $t = 5.4 \times 10^3$, respectively. The third column depicts the isosurface corresponding to the evolution shown in column two. . . . 112
- 5.1 (a) Variation of average velocity profile $\langle v_x \rangle$ along z -direction for all cases mentioned in the legends. The data is plotted for a fixed shear rate $\dot{\gamma} = 1.56 \times 10^{-2}$ at $t = 1.2 \times 10^3$. (b) Comparison of shear viscosity $\langle \eta \rangle$ as a function of $\dot{\gamma}$ at $t = 1.2 \times 10^3$. (c-e) $\langle v_x \rangle$ vs. z for different shear rates for cases 2-4, respectively. 121
- 5.2 The evolution snapshots are shown at $t = 0$ in (a) and $t = 1.2 \times 10^3$ in (b); the red, yellow, and olive colors represent the A , B , and wall-type beads, respectively. (c) The spatial intensity variation of $S(k_x, k_z)$ at $t = 1.2 \times 10^3$ along xz -plane. 123
- 5.3 (a) The RDF $g_{AB}(r)$ vs. r is compared at different time steps for zero shear rate. (b) The scaled correlation function $C(r, t)$ vs. $r/R(t)$. The solid black line represents the zero crossing. (c) The characteristic domain growth $R(t)$ against time is plotted after averaging along x , y , and z directions, as shown in the legends. The inset figure represents the spherically averaged length scale, $R(t)$ vs. t . Solid black lines with slopes 1 and $2/3$ represent the viscous and inertial hydrodynamic growth laws. (d) The unidirectional structure factor $S(k_x, k_y, k_z)$ as a function of k_x , k_y , and k_z . The inset correspond to $S(k_x, k_y, k_z)$ vs. k_y along diagonal and cross-diagonal directions. 124

- 5.4 (a-c) The evolution snapshots and corresponding xz -cross section are displayed for case 2 with different shear rates: (a) $\dot{\gamma} = 1.56 \times 10^{-2}$, (b) $\dot{\gamma} = 7.80 \times 10^{-3}$, and (c) $\dot{\gamma} = 1.56 \times 10^{-3}$. Related isosurfaces are also provided in the inset of (d). (d) and (e) show $C(r, t)$ vs. $r/R(t)$ and $R(t)$ vs. t , respectively, for different shear rates at $t = 1.2 \times 10^3$. The solid black lines in (e) represent the growth laws followed by $R(t)$ 126
- 5.5 The evolution snapshots for case 3 and corresponding xz -cross-section plots at $t = 1.2 \times 10^3$ is displayed in (a-c) with different shear rates mentioned alongside them. Related isosurfaces are also provided in the inset of (d). Data for $C(r, t)$ vs. $r/R(t)$ and $R(t)$ vs. t are plotted in (d) and (e), respectively, for different shear rates shown in the legends at $t = 1.2 \times 10^3$. The solid black lines in (e) represent the growth laws followed by $R(t)$. . . 128
- 5.6 (a-c) The evolution morphology and corresponding xz -cross section plots for the middle layer ($y = 32$). These are the snapshots at the late time intervals $t = 1.2 \times 10^3$ for different shear rates mentioned alongside them. (d) $g_{AB}(r)$ vs. r for different $\dot{\gamma}$ values. Inset represents the isosurfaces corresponding to the evolution pictures given in (a-c). (e) $C(r, t)$ vs. $r/R(t)$ is plotted for the same data set as in (d). The respective plots for $R(t)$ vs. t averaged along the radial direction are illustrated in the inset of (e). (f-g) The average length scale along different directions is plotted for various $\dot{\gamma}$ values. Solid lines with slopes 1 and $2/3$ represent the viscous and inertial hydrodynamic growth laws followed by the system. 129

- 5.7 (a-d) The spatial intensity variation of the structure factor $S(k_x, k_z)$ along xz -plane for different $\dot{\gamma}$ values at $t = 1.2 \times 10^3$. To see the structural anisotropy, the unidirectional structure factor $S(k_x, k_y, k_z)$ vs. k_x (black curve), k_y (red curve), and k_z (green curve) is plotted in (e), (f), and (g) for $\dot{\gamma} = 1.56 \times 10^{-3}$, 7.80×10^{-3} , and 1.56×10^{-2} , respectively. The magenta and blue curves demonstrate $S(k_x, k_y, k_z)$ vs. k_y along diagonal and cross-diagonal directions. 132
- 5.8 The evolution snapshots are shown at $t = 0$ in (a) and $t = 1.2 \times 10^3$ in (b). (c) The spatial intensity variation of $S(k_x, k_z)$ at $t = 1.2 \times 10^3$ along xz -plane. 134
- 5.9 Off-critical polymer mixture with ratio $A : B = 3 : 1$: (a) $g_{AB}(r)$ vs. r at various time steps mentioned in the legends. (b) Plots for $C(r, t)$ vs. $r/R(t)$ for the same data set used in (a). (c) Characteristic length scale $R(t)$ vs. t is represented along different directions with various symbol types in the legends. Inset plots the time-dependent spherically averaged domain length. (d) Plots for $S(k_x, k_y, k_z)$ as a function of k_x , k_y , and k_z , representing the anisotropy along different directions. In the Inset, the magenta and blue curves plot $S(k_x, k_y, k_z)$ vs. k_y along diagonal and cross diagonal directions on xz -plane. 134
- 5.10 (a-c) Domain evolution and $2d$ cross-section plots along xz -plane for the case 2. These pictures are related to different shear rates mentioned at left for the late time $t = 1.2 \times 10^3$. Related isosurfaces are also plotted in the inset of (d). (d) Scaled correlation as a function of $r/R(t)$ for the same case as in (a-c). The corresponding time-dependent average domain size is plotted in (e). $R(t)$ curves follow the viscous and inertial hydrodynamic growth represented by solid black lines with slopes 1 and $2/3$, respectively. 136

- 5.11 (a-c) Evolution snapshots and corresponding xz -cross section plots for $y = 32$ at $t = 1.2 \times 10^3$. Related isosurfaces are also shown in the inset of (d). (d) Comparison of Scaled correlation $C(r, t)$ vs. $r/R(t)$ corresponding to the evolutions plotted in (a-c). (e) $R(t)$ vs. t for different shear rates mentioned in the legends. Solid black lines with slope 1 and $2/3$ represent the growth laws followed by $R(t)$ 137
- 5.12 (a-c) Plots for the late-time domain evolution and corresponding xz -cross section plots for $y = 32$ at different shear rates. (d) The RDF plots for different $\dot{\gamma}$ values at $t = 1.2 \times 10^3$. Inset represents the isosurfaces corresponding to the evolution pictures displayed in (a-c). (e) $C(r, t)$ vs. $r/R(t)$ for different shear rates, same as in (d). A solid black line depicts zero crossing. Inset represents the plots for $R(t)$ vs. t for different $\dot{\gamma}$ values. (f-h) $R(t)$ vs. t averaged along x , y , and z directions for different shear rates given in each figure. $R(t)$ curves show a cross-over from viscous to inertial hydrodynamic growth represented by solid lines. 139
- 5.13 (a-d) The intensity variation of the structure factor along xz -plane for different shear rates at $t = 1.2 \times 10^3$. Unidirectional structure factor $S(k_x, k_y, k_z)$ vs. k_x , k_y , and k_z is compared for $\dot{\gamma} = 1.56 \times 10^{-3}$ in (e), $\dot{\gamma} = 7.80 \times 10^{-3}$ in (f), and $\dot{\gamma} = 1.56 \times 10^{-2}$ in (g) at $t = 1.2 \times 10^3$. The inset of each picture depict the variation of structure factor as a function of k_y along diagonal and cross-diagonal directions. 141
- 6.1 Phase separation kinetics of different systems inside hydrogel. 147
- 6.2 Mixing and de-mixing in the dimer melt system on the application of alternate on-off light cycles. 147