

Table of contents

List of figures	xv
List of tables	xx
1 Introduction	1
1.1 Introduction	1
1.2 The Interstellar Medium (ISM)	3
1.3 Probing the ISM	5
1.4 Turbulence in the ISM	7
1.5 This Thesis	11
2 Techniques of Radio Interferometry	15
2.1 Introduction	15
2.2 Radio Interferometers	16
2.3 Data analysis methodologies	22
3 Trans-Alfvenic magnetohydrodynamic turbulence in the vicinity of supernova remnant Cassiopeia-A shocks	31
3.1 Introduction	31
3.2 Method	33
3.3 Analysis	34

3.4	Results	39
3.5	Discussion	40
4	H I Column Density Statistics of the Cold Neutral Medium from Absorption Studies	44
4.1	Introduction	44
4.2	Two point statistics of the optical depth from observed visibilities	46
4.2.1	Specific intensity of 21-cm radiation	46
4.2.2	Autocorrelation function of optical depth fluctuations	47
4.2.3	Measuring the optical depth autocorrelation from radio-interferometric observations	48
4.3	Relation of optical depth autocorrelation with statistics of the H I column density and spin temprature	49
4.4	Sensitivity of observables on the physical parameters	52
4.4.1	With isotropic column density fluctuations	52
4.4.2	Effect of anisotropic fluctuations	54
4.5	Discussion and Conclusion	57
5	H I absorption towards the Tycho's supernova remnant.	60
5.1	Tycho's Supernova remnant	61
5.2	Observation and Data Analysis	63
5.3	Spectral Analysis and Result	66
5.4	Discussion and Conclusion	75
6	Conclusions and Future Scope	78
	References	81

Table of contents	xxi
-------------------	------------

Appendix A	93
-------------------	-----------

A.1	93
---------------	----

List of figures

1.1	: Left: The figure shows the Galactic positions of the six supernovae remnants along with their names. Right: The corresponding Galactic spiral arms are also shown, from which H I absorption is expected toward these remnants. The left figure is taken from <i>Annu.Rev.Astron.Astrophys</i> (2012), originally made by R. Hurt (in collaboration with R. Benjamin) and available in Churchwell et al. (2009). The figure of Galactic structures (in the background) first appeared in <i>Astronomical Society of the Pacific</i> (Copyright 2009, Astron.Soc.Pac.)	13
2.1	: Image shows the GMRT antenna positions, taken from Swarup et al. (1991). The dot points inside the zoomed circle are the central part of the GMRT consisting 14 antennas.	20
2.2	: The above figure shows the observation strategy of a target source along with calibrators. Flux Calibrators (FC1 and FC2) and Bandpass Calibrators (BP1 and BP2) are observed at the beginning and end, while Phase Calibrator (PC) and Target source (TS) are observed alternatively. .	24

- 5.1 : a) Continuum image of the Tycho's SNR along with point source at 00:29:45.475, +63.58.40.727 (J2000) (inside circle) made using the baseline range 0-25K λ . Since in our image only this point source is visible, so we will call it the point source at all references for it, b) Low resolution (0-5K λ) continuum image of Tycho's SNR. The numbers given in the horizontal and vertical axes in images correspond to pixels. For the 0-25K λ image each pixel has a dimension of 4arcsec \times 4arcsec. For 0-5K λ , the pixel size is 8.0arcsec \times 8.0arcsec. 66

- 5.2 : Figure shows the six optical depth images averaged in the velocity ranges, as shown in the labels. These images are made using the baseline range of 0-5K λ . Optical depth images made by averaging in LSR velocity ranges from -25.85 km/s to -33.57 km/s, -23.28 km/s to -16.84 km/s, -16.84 km/s to -09.08 km/s, -09.08km/s to -00.08km/s and -00.08km/s to 08.94 km/s show significant absorptions at the overall face of Tycho's SNR, while in the range -45.20 km/s to -51.63 km/s, it shows higher absorption toward the eastern edge. The quantity shown in the image is optical depth and bar at the bottom of the image shows its magnitude. 67

- 5.3 : Optical depth spectra (with 1σ error bars) of H I absorption towards point source (PS) and Tycho's SNR in the baseline range $1.3-40K\lambda$ are shown as upper panels in the above figures. H I emission profiles, taken from LAB survey (Kalberla et al., 2005) in their directions with velocity resolution of 1.30 km/s are shown as lower panels. The typical per channel sensitivity in brightness temperature for the LAB survey is $\sim 0.1 \text{ K}$. Vertical lines, marked as I, II, III, IV and V in these spectra show velocities 0.59 km/s , -15.84 km/s , -26.48 km/s , -51.62 km/s and -62.25 km/s and are made in reference to the maximum absorptions in the direction of point source in the windows of its absorption as mentioned in the text. 68
- 5.4 : Left: Comparisons of the optical depth spectra of point source with two optical depth spectra toward the left and right face of Tycho's SNR. Right: Four spectra of optical depth toward the left region of Tycho's SNR. The velocity resolution of the spectra is 1 km/s . The marked vertical lines refers to the velocities as shown in figure 5.3. 70
- 5.5 : Optical depth spectra for different regions along with spectra of point source (PS) are shown. The figure helps to easily compare the spectra correlated with each other. 72
- 5.6 : Similarity index and Spearman's rank (S R) correlation between the spectra of the point source (reg 0) and four different regions (reg8, reg 9, reg13 and reg18) of Tycho's SNR. Vertical lines represents the marked velocities as shown in figure 5.3 73

5.7	: Approximate location of Tycho’s SNR (red dot) in our Galaxy and geometry to calculate the excess velocity of cloud in the direction of the point source. The left figure is taken from <i>Annu.Rev.Astron.Astrophys</i> (2012), originally made by R. Hurt (in collaboration with R. Benjamin) and available in Churchwell et al. (2009). The figure of galactic structures (in the background) first appeared in <i>Astronomical Society of the Pacific</i> (Copyright 2009,Astron.Soc.Pac.)	74
A.1	: The figure shows the modified version of figure 3.2. We have relatively scaled the amplitudes to clarify the differences in the spectra.	94
A.2	: Similar to the figure A.1, this figure also show the modified version of the figure 3.4. Amplitudes are scaled for the purpose of visualization.	94
A.3	: Normalized statistics of the magnetic field disturbances from the radial region of 0.526R to 0.820R.	95
A.4	: The figure shows the normalized magnetic energy spectra from the radial region of 0.826R to 1.093R	96

List of tables

3.1	The position and width information about the different regions of the supernova used in the analysis.	36
3.2	Fitted values of α for the spectra shown in figures 3.2 and 3.4.	39
4.1	Different auto-correlation functions and the corresponding power spectrum of the observables used in this chapter.	46
5.1	Various distance estimates of Tycho's SNR from different references.	62
5.2	Observation summary for GWB and GSB.	64
5.3	Details of the calibrators for this observation.	65