

Contents

1	Introduction and literature review	1
1.1	Introduction to the family of Hall effect	2
1.1.1	Ordinary Hall effect	2
1.1.2	Anomalous Hall effect	3
1.1.3	Spin Hall effect	4
1.1.4	Quantum Hall effect	5
1.1.5	Quantum spin Hall effect	5
1.1.6	Quantum anomalous Hall effect	6
1.1.7	Anomalous Nernst effect	7
1.1.8	Topological Hall effect	7
1.2	Origin of anomalous Hall effect	11
1.2.1	Extrinsic mechanism	11
1.2.2	Intrinsic mechanism	14
1.2.3	Scaling relation and progress in anomalous Hall effect	18
1.3	Topological materials	19
1.3.1	Topological insulators and Quantum Hall state	21
1.3.2	Topological semimetal	24
1.3.3	Topological semimetal: Source of Berry curvature	32
1.4	Heusler alloys	35
1.4.1	Heusler alloys as topological semimetal: Realization of large anomalous transport response	37

1.5	Objective of present thesis	39
2	Methods	66
2.1	Experimental method	67
2.1.1	Synthesis	67
2.1.2	X-ray diffraction	68
2.1.3	Synchrotron X-ray diffraction	70
2.1.4	Scanning electron microscope	72
2.1.5	Magnetization measurements	74
2.1.6	Transport measurements	75
2.2	Theoretical method	76
2.2.1	Density Functional Theory	76
2.2.2	Wannier interpolation	81
2.2.3	Micromagnetic simulation: A tool for simulation of skyrmions	86
3	Anomalous Hall effect due to gapped nodal line in Co_2FeGe Heusler compound	92
3.1	Introduction	93
3.2	Experiment and method	94
3.3	Result and discussion	95
3.3.1	Structural properties	95
3.3.2	Magnetization and resistivity measurements	97
3.3.3	Anomalous Hall	97
3.3.4	Theoretical calculations	100
3.4	Conclusion	103
4	Antisite disorder induced Berry phase and anomalous Hall conductivity in Co_2FeAl Heusler compound	111
4.1	Introduction	112
4.2	Methods	113
4.3	Result and discussion	114
4.3.1	Structural analysis	114

4.3.2	Magnetization and resistivity	116
4.3.3	Anomalous Hall	116
4.3.4	Theoretical calculations	119
4.4	Conclusion	121
5	Band splitting induced Berry flux and intrinsic anomalous Hall conductivity in NiCoMnGa quaternary Heusler compound	129
5.1	Introduction	130
5.2	Methods	131
5.3	Result and discussion	132
5.3.1	Structure and magnetization	132
5.3.2	Transport measurements	134
5.3.3	First principles calculation	138
5.4	Conclusion	140
6	Nodal-line symmetry breaking induced colossal anomalous Hall and Nernst effects in Cu₂CoSn Heusler compound	146
6.1	Introduction	147
6.2	Computational details	149
6.3	Result and discussion	150
6.4	Conclusion	156
7	Observation of the topological Hall effect and signature of skyrmions in high Curie-temperature Co₂FeAl Heusler compound	163
7.1	Introduction	164
7.2	Result and discussion	165
7.2.1	Sample preparation and characterization method	165
7.2.2	Magnetization and magnetoresistance measurements	166
7.2.3	Topological Hall effect	167
7.2.4	AC-susceptibility measurement	169
7.2.5	Micromagnetic simulation	171

7.3	Conclusion	172
8	Summary and suggestions for future work	181
8.1	Summary	182
8.2	Future work	184