

Contents

Title	Page No.
Title of thesis	i
Dedication	ii
Certificate	iii
Declaration by the candidate and certificate by the supervisor	iv
Copyright transfer certificate	v
Acknowledgment	vi-vii
Contents	viii-xii
List of Figures	xiii-xx
List of Tables	xxi
List of Symbols/Abbreviations	xxii
Preface	xxiii-xxvii
Chapter 1: Introduction	1-29
1. Introduction	1
1.1. Photoredox reaction	1
1.2. Photocatalysis with semiconductor	3
1.3. Thermodynamic feasibility of photoredox reactions	4
1.4. Current strategies to enhance charge transport and suppress charge recombination in photoredox reactions	5
1.4.1. Roles of cocatalysts in photocatalytic reactions	5
1.4.2 Categories of cocatalysts	6
1.4.3 Construction of heterojunction-based photocatalysts	7
1.4.4 Type-I, Type-II, Type-III, Z-Scheme, and S-Scheme heterojunctions	7
1.5 Metal halide perovskite photocatalyst	9
1.5.1. Application of MHP photocatalyst	10
1.6. Structural features of MHP	11
1.7. Optical properties of MHP quantum dots (QDs)	12
1.8. MHP QDs for photoredox reactions	13

1.9. Band structure of CsPbBr ₃ QDs	14
1.10. Stability of metal halide perovskite	15
1.11. Characterization techniques for charge behaviour induced by cocatalysts	16
1.12. Role of reactive oxygen species (ROS) in photocatalysis	16
1.13. Importance of amide, amine, and heterocyclic compounds	16
1.13.1. Importance of amides	16
1.13.2 Importance of Amines	18
1.13.3. Importance of isoquinoline compounds	20
1.14. Objectives of the thesis	21
1.15. References:	23
Chapter-2: Lead halide perovskite-nickel(II) complex improves visible light photoredox C–N coupling	30-59
2.1. Introduction	30
2.2. Chemicals	31
2.3. Instruments	31
2.4. Experimental	33
2.4.1 Synthesis of CsPbBr ₃ QDs	33
2.4.2. Synthesis of Nickel(II) dimethylglyoxime complex	34
2.4.3. Preparation of [Ni]-CsPbBr ₃	34
2.5. Photocatalytic activity test	35
2.6. Working electrode preparation and photoelectrochemical measurements	35
2.7. Results and discussion	36
2.7.1. Characterizations of the catalysts	36
2.7.2 Optoelectronic properties of catalysts	36
2.7.3. Role of cocatalyst on photoredox activity	41
2.8. Optimization of the photoredox reaction conditions	42
2.9. Substrate scope for the C–N bond formation	42
2.10. Cocatalyst effects on ultrafast dynamics in CsPbBr ₃ QDs	44

2.11. Reaction mechanism	49
2.12. Conclusion	51
2.13. ^1H NMR and ^{13}C NMR spectra of the product	56
2.14 References	57
Chapter-3: Halide perovskite-cobalt(III) complex photoredox catalyst for highly efficient N-Alkylation of amines with alcohols	60-83
3.1. Introduction	60
3.2. Chemicals	62
3.3. Instruments	62
3.4. Experimental	62
3.4.1. Synthesis of CsPbBr ₃ QDs	62
3.4.2. Synthesis of oxime ligands and metal complexes (Co-1, Co-2, Co-3, and Co-4)	62
3.4.3. Preparation of [Co]-CsPbBr ₃ catalysts	64
3.5. Results and discussion	64
3.5.1. Characterizations of CsPbBr ₃ QDs systems	64
3.6. Optical properties of the catalyst systems	65
3.7. Role of cocatalyst on charge transfer properties of CsPbBr ₃ QDs	66
3.8. Optimization of reaction conditions	67
3.9. Reaction mechanism	70
3.10. Control study of N-Alkylation of aniline	71
3.11. Substrate scope of amine formation	75
3.12. Conclusion	76
3.13. ^1H NMR and ^{13}C NMR spectra of the product	80
3.14. References	81
Chapter-4: Z-Scheme CsPbBr₃/BiOBr heterojunction for selective semihydrogenation of 1,2,3,4-tetrahydroisoquinoline to 3,4-dihydroisoquinoline	84-107
4.1. Introduction	84
4.2. Chemicals	86

4.3. Instruments	86
4.4. Experimental	86
4.4.1. Synthesis of CsPbBr ₃ QDs	86
4.4.2. Synthesis of ultrathin BiOBr	86
4.4.3. Construction of heterojunction (Z-20)	87
4.5. Results and discussion	88
4.5.1. Syntheses and characterizations of heterojunction catalysts	88
4.5.2. Z-scheme vs type-II heterojunction	90
4.5.3. Z-Scheme heterojunction	92
4.6. Charge transfer dynamics	93
4.7. Femtosecond transient absorption spectroscopy study	94
4.8. Photocatalytic semidehydrogenation reaction	98
4.9. Mechanistic study of semidehydrogenation	100
4.10. Quenching experiment	103
4.11. Conclusion	104
4.12. ¹ H NMR and ¹³ C NMR spectra of the product	105
4.13. References	106
Chapter-5: Photoredox molecular oxygen activation with lead-free metal halide perovskite for oxidative amidation of alcohols	108-129
5.1. Introduction	108
5.2. Chemicals	110
5.3. Instruments	110
5.4. Experimental	110
5.4.1. Synthesis of Cs ₂ CuBr ₄ by hot injection method (PC-1)	110
5.4.2. Synthesis of Cs ₂ CuBr ₄ at room temperature (PC-2)	111
5.5. Results and discussion	111
5.5.1. Syntheses and characterizations of the catalysts	111
5.6. Optoelectronic properties of the photocatalysts	114
5.7. Optimization of reaction conditions for the amidation of alcohols	116
5.8. Substrate scope for the amide synthesis	117
5.9. Reaction mechanism	119

5.10. Quenching experiments	120
5.11. Conclusions	122
5.12. ^1H NMR and ^{13}C NMR spectra of the product	126
5.13. References	127
Chapter-6: Summary and Future Perspective	130-134
6.1. Conclusions	130
6.2. Future scope and perspective	133
List of publications	135