

- Afiattalab F. Mutations affecting segment number and polarity in *Drosophila*. *Nature*. 1980;287:795-801.
- Akhtar Y, Isman MB, Niehaus LA, Lee C-H, Lee H-S. Antifeedant and toxic effects of naturally occurring and synthetic quinones to the cabbage looper, *Trichoplusia ni*. *Crop protection*. 2012;31(1):8-14.
- Araujo J, Logothetis C. Dasatinib: a potent SRC inhibitor in clinical development for the treatment of solid tumors. *Cancer treatment reviews*. 2010;36(6):492-500.
- Atun R, Davies JI, Gale EAM, Barnighausen T, Beran D, Kengne AP, et al. Diabetes in sub-Saharan Africa: from clinical care to health policy. *Lancet diabetes endocrinol*. 2017;5(8):622-67.
- Bachelder RE, Crago A, Chung J, Wendt MA, Shaw LM, Robinson G, et al. Vascular endothelial growth factor is an autocrine survival factor for neuropilin-expressing breast carcinoma cells. *Cancer research*. 2001;61(15):5736-40.
- Bajaj S, Roy PP, Singh J. Synthesis, thymidine phosphorylase inhibitory and computational study of novel 1, 3, 4-oxadiazole-2-thione derivatives as potential anticancer agents. *Computational biology and chemistry*. 2018;76:151-60.
- Bala BD, Muthusaravanan S, Choon TS, Ali MA, Perumal S. Sequential synthesis of amino-1, 4-naphthoquinone-appended triazoles and triazole-chromene hybrids and their antimycobacterial evaluation. *European journal of medicinal chemistry*. 2014;85:737-746.
- Barr FA, Silljé HH, Nigg EA. Polo-like kinases and the orchestration of cell division. *Nature reviews Molecular cell biology*. 2004;5(6):429-440.
- Barr MP, Bouchier-Hayes DJ, Harmey JJ. Vascular endothelial growth factor is an autocrine survival factor for breast tumour cells under hypoxia. *International journal of oncology*. 2008;32(1):41-48.
- Bar-Sagi D, Hall A. Ras and Rho GTPases: a family reunion. *Cell*. 2000;103(2):227-38.
- Basmaciyan L, Jacquet P, Azas N, Casanova M. A novel hydrolase with a pro-death activity from the protozoan parasite *Leishmania major*. *Cell death discovery*. 2019;5(1):99-112.
- Basu A. DNA damage, mutagenesis and cancer. *International journal of molecular sciences*. 2018;19(4):970-983.
- Bates DO. Vascular endothelial growth factors and vascular permeability. *Cardiovascular research*. 2010;87(2):262-271.
- Behrens J, Frixen U, Schipper J, Weidner M, Birchmeier W, editors. Cell adhesion in invasion and metastasis. *Seminars in cell biology*. 1992;13(3):169-178.
- Bhatia S, Krailo MD, Chen Z, Burden L, Askin FB, Dickman PS, et al. Therapy-related myelodysplasia and acute myeloid leukemia after Ewing sarcoma and primitive neuroectodermal tumor of bone: A report from the Children's Oncology Group. *Blood*. 2007;109(1):46-51.
- Bleyer A, Barr R, Hayes-Lattin B, Thomas D, Ellis C, Anderson B. The distinctive biology of cancer in adolescents and young adults. *Nature Reviews Cancer*. 2008;8(4):288-298.

- Bolognesi ML, Lizzi F, Perozzo R, Brun R, Cavalli A. Synthesis of a small library of 2-phenoxy-1,4-naphthoquinone and 2-phenoxy-1,4-anthraquinone derivatives bearing anti-trypanosomal and anti-leishmanial activity. *Bioorganic & medicinal chemistry letter*. 2008 1;18(7):2272-2276.
- Boocock CA, Charnock-Jones DS, Sharkey AM, McLaren J, Barker PJ, Wright KA, et al. Expression of Vascular Endothelial Growth Factor and Its Receptors fit and KDR in Ovarian Carcinoma. *JNCI: Journal of the National Cancer Institute*. 1995;87(7):506-516.
- Breiman RF, Cosmas L, Njuguna H, Audi A, Olack B, Ochieng JB, et al. Population-based incidence of typhoid fever in an urban informal settlement and a rural area in Kenya: implications for typhoid vaccine use in Africa. *PLoS one*. 2012;7(1):29119-22229.
- Brown JM, Wouters BG. Apoptosis, p53, and tumor cell sensitivity to anticancer agents. *Cancer research*. 1999;59(7):1391-1399.
- Brun M-P, Braud E, Angotti D, Mondésert O, Quaranta M, Montes M, et al. Design, synthesis, and biological evaluation of novel naphthoquinone derivatives with CDC25 phosphatase inhibitory activity. *Bioorganic & medicinal chemistry*. 2005;13(16):4871-4879.
- Bruyère C, Meijer L. Targeting cyclin-dependent kinases in anti-neoplastic therapy. *Current opinion in cell biology*. 2013;25(6):772-779.
- Bryden M, Evans H, Keeler R. Cyclopia in sheep caused by plant teratogens. *Journal of anatomy*. 1971;110(3):507-527.
- Buchdunger E, Zimmermann J, Mett H, Meyer T, Müller M, Druker BJ, et al. Inhibition of the Abl protein-tyrosine kinase in vitro and in vivo by a 2-phenylaminopyrimidine derivative. *Cancer research*. 1996;56(1):100-104.
- Cairns RA, Harris IS, Mak TW. Regulation of cancer cell metabolism. *Nature reviews cancer*. 2011;11(2):85-95.
- Cardoso SH, de Oliveira CR, Guimaraes AS, Nascimento J, de Oliveira Dos Santos Carmo J, de Souza Ferro JN, et al. Synthesis of newly functionalized 1,4-naphthoquinone derivatives and their effects on wound healing in alloxan-induced diabetic mice. *Chem Biol Interact*. 2018;291:55-64.
- Carter-Cooper BA, Fletcher S, Ferraris D, Choi EY, Kronfli D, Dash S, et al. Synthesis, characterization and antineoplastic activity of bis-aziridinyl dimeric naphthoquinone—A novel class of compounds with potent activity against acute myeloid leukemia cells. *Bioorganic & medicinal chemistry letters*. 2017;27(1):6-10.
- Castro MA, Gamito AM, Tangarife-Castano V, Zapata B, Miguel del Corral JM, Mesa-Arango AC, et al. Synthesis and antifungal activity of terpenyl-1,4-naphthoquinone and 1,4-anthracenedione derivatives. *European Journal of Medicinal Chemistry*. 2013;67:19-27.
- Chan MM, Lu X, Merchant FM, Iglehart JD, Miron PL. Gene expression profiling of NMU-induced rat mammary tumors: cross species comparison with human breast cancer. *Carcinogenesis*. 2005;26(8):1343-53.
- Charlier C, Hart E, Lefort A, Ribaud P, Dromer F, Denning D, et al. Fluconazole for the management of invasive candidiasis: where do we stand after 15 years. *Journal of antimicrobial chemotherapy*. 2006;57(3):384-410.
- Chung AS, Ferrara N. Developmental and pathological angiogenesis.

- Annual review of cell and developmental biology*. 2011;27:563-84.
- Chung Y, Shin YK, Zhan CG, Lee S, Cho H. Synthesis and evaluation of antitumor activity of 2- and 6-[(1,3-benzothiazol-2-yl)aminomethyl]-5,8-dimethoxy-1,4-naphthoquinone derivatives. *Archives of pharmacal research*. 2004;27(9):893-900.
 - Cooper GM, Hausman RE. *The Cell: a molecular approach*. 2000. 7th edition ASM Press, Washington DC. 2000:467-519.
 - Costa MCA, Ferreira MMC. Two-dimensional quantitative structure-activity relationship study of 1,4-naphthoquinone derivatives tested against HL-60 human promyelocytic leukaemia cells. *SAR QSAR Environmental Research*. 2017;28(4):325-339.
 - Dang CV. c-Myc target genes involved in cell growth, apoptosis, and metabolism. *Molecular and cellular biology*. 1999;19(1):1-11.
 - Danne AB, Choudhari AS, Chakraborty S, Sarkar D, Khedkar VM, Shingate BB. Triazole–diindolylmethane conjugates as new antitubercular agents: synthesis, bioevaluation, and molecular docking. *MedChemComm*. 2018;9(7):1114-1130.
 - de Rezende LCD, Fumagalli F, Bortolin MS, de Oliveira MG, de Paula MH, de Andrade-Neto VF, et al. In vivo antimalarial activity of novel 2-hydroxy-3-anilino-1, 4-naphthoquinones obtained by epoxide ring-opening reaction. *Bioorganic & medicinal chemistry letters*. 2013;23(16):4583-4586.
 - de Souza NB, de Andrade IM, Carneiro PF, Jardim GA, de Melo IM, da Silva Junior EN, et al. Blood schizonticidal activities of phenazines and naphthoquinoidal compounds against *Plasmodium falciparum* in vitro and in mice malaria studies. *Memórias do Instituto Oswaldo Cruz*. 2014;109(5):546-552.
 - Demoulin J-B, Montano-Almendras CP. Platelet-derived growth factors and their receptors in normal and malignant hematopoiesis. *American journal of blood research*. 2012;2(1):44-56.
 - Diehn M, Cho RW, Lobo NA, Kalisky T, Dorie MJ, Kulp AN, et al. Association of reactive oxygen species levels and radioresistance in cancer stem cells. *Nature*. 2009;458(7239):780-783.
 - Diogo EB, Dias GG, Rodrigues BL, Guimaraes TT, Valenca WO, Camara CA, et al. Synthesis and anti-Trypanosoma cruzi activity of naphthoquinone-containing triazoles: electrochemical studies on the effects of the quinoidal moiety. *Bioorganic & medicinal chemistry*. 2013;21(21):6337-6348.
 - Doncel G, Mauck C. Vaginal microbicides: a novel approach to preventing sexual transmission of HIV. *Current HIV/AIDS Reports*. 2004;1(1):25-32.
 - Du J-Z, Du X-J, Mao C-Q, Wang J. Tailor-made dual pH-sensitive polymer–doxorubicin nanoparticles for efficient anticancer drug delivery. *Journal of the american chemical society*. 2011;133(44):17560-17563.
 - Duronio RJ, Xiong Y. Signaling pathways that control cell proliferation. *Cold Spring Harbor perspectives in biology*. 2013;5(3):8904-8916.
 - Ece Gurdal E, Buclulgan E, Durmaz I, Cetin-Atalay R, Yarim M. Synthesis and anticancer activity evaluation of some benzothiazole-piperazine derivatives. *Anti-cancer agents in medicinal chemistry (formerly current*

- medicinal chemistry-anti-cancer agents*). 2015;15(3):382-389.
- Ehrhardt K, Davioud-Charvet E, Ke H, Vaidya AB, Lanzer M, Deponte M. The antimalarial activities of methylene blue and the 1, 4-naphthoquinone 3-[4-(trifluoromethyl) benzyl]-menadione are not due to inhibition of the mitochondrial electron transport chain. *Antimicrobial agents and chemotherapy*. 2013;57(5):2114-2120.
 - El-Din MMG, El-Gamal MI, Abdel-Maksoud MS, Yoo KH, Oh C-H. Synthesis and broad-spectrum antiproliferative activity of diarylamides and diarylureas possessing 1, 3, 4-oxadiazole derivatives. *Bioorganic & medicinal chemistry letters*. 2015;25(8):1692-1699.
 - Ellis LM, Hicklin DJ. Pathways mediating resistance to vascular endothelial growth factor-targeted therapy. *Clinical cancer research*. 2008;14(20):6371-6375.
 - Epifano F, Genovese S, Fiorito S, Mathieu V, Kiss R. Lapachol and its congeners as anticancer agents: a review. *Phytochemistry reviews*. 2014;13(1):37-49.
 - Errante G, La Motta G, Lagana C, Wittebolle V, Sarciron ME, Barret R. Synthesis and evaluation of antifungal activity of naphthoquinone derivatives. *European Journal of Medicinal Chemistry*. 2006 ;41(6):773-778.
 - Fan F, Wey JS, McCarty MF, Belcheva A, Liu W, Bauer TW, et al. Expression and function of vascular endothelial growth factor receptor-1 on human colorectal cancer cells. *Oncogene*. 2005;24(16):2647-2653.
 - Fantl WJ, Johnson DE, Williams LT. Signalling by receptor tyrosine kinases. *Annual review of biochemistry*. 1993;62(1):453-481.
 - Farsinejad S, Gheisary Z, Samani SE, Alizadeh AM. Mitochondrial targeted peptides for cancer therapy. *Tumor biology*. 2015;36(8):5715-5725.
 - Ferrara N. Vascular endothelial growth factor. *Arteriosclerosis, thrombosis, and vascular biology*. 2009;29(6):789-791.
 - Ferrara N, Carver Moore K, Chen H, Dowd M, Lu L, O'shea KS, et al. Heterozygous embryonic lethality induced by targeted inactivation of the VEGF gene. *Nature*. 1996;380(4):1-9.
 - Ferrara N, Houck K, Jakeman L, Leung DW. Molecular and biological properties of the vascular endothelial growth factor family of proteins. *Endocrine reviews*. 1992;13(1):18-32.
 - Fieser LF, M F. Organic Chemistry third edition: D.C. Heath and Company. Boston,. 1956.
 - Fons N. Focus: Drug Development: Textbook of Drug Design and Discovery. The Yale journal of biology and medicine. 2017; 5th edition 90(1);436.
 - Frank S, Hübner G, Breier G, Longaker MT, Greenhalgh DG, Werner S. Regulation of vascular endothelial growth factor expression in cultured keratinocytes. Implications for normal and impaired wound healing. *Journal of biological chemistry*. 1995;270(21):12607-12613.
 - Fry AM, Mayor T, Meraldi P, Stierhof Y-D, Tanaka K, Nigg EA. C-Nap1, a novel centrosomal coiled-coil protein and candidate substrate of the cell cycle-regulated protein kinase Nek2. *The Journal of cell biology*.

- 1998;141(7):1563-1574.
- Fuentes M, Araya-Maturana R, Palomo I, Fuentes E. Platelet mitochondrial dysfunction and mitochondria-targeted quinone-and hydroquinone-derivatives: Review on new strategy of antiplatelet activity. *Biochemical pharmacology*. 2018;156:215-222.
 - Fukasawa M, Korc M. Vascular endothelial growth factor-trap suppresses tumorigenicity of multiple pancreatic cancer cell lines. *Clinical cancer research*. 2004;10(10):3327-3332.
 - Gadbois DM, Hamaguchi JR, Swank RA, Bradbury EM. Staurosporine is a potent inhibitor of p34cdc2 and p34cdc2-like kinases. *Biochemical and biophysical research communications*. 1992;184(1):80-85.
 - Gama BE, de Oliveira NKA, de Souza JM, Santos F, de Carvalho LJ, Melo YF, et al. Brazilian Plasmodium falciparum isolates: investigation of candidate polymorphisms for artemisinin resistance before introduction of artemisinin-based combination therapy. *Malaria journal*. 2010;9(1):355-340.
 - Gautam MK, Besan M, Pandit D, Mandal S, Chadha R. Cocrystal of 5-Fluorouracil: Characterization and Evaluation of Biopharmaceutical Parameters. *AAPS PharmSciTech*. 2019;20(4):149-165.
 - Gelb MH, Hol WG. Drugs to combat tropical protozoan parasites. *Science*. 2002;297(5580):343-434.
 - Gerber H-P, Vu TH, Ryan AM, Kowalski J, Werb Z, Ferrara N. VEGF couples hypertrophic cartilage remodeling, ossification and angiogenesis during endochondral bone formation. *Nature medicine*. 1999;5(6):623-628.
 - Gokhale NH, Shirisha K, Padhye SB, Croft SL, Kendrick HD, McKee V. Metalloantimalarials: synthesis, X-ray crystal structure of potent antimalarial copper (II) complex of arylazo-4-hydroxy-1,2-naphthoquinone. *European Journal of Medicinal Chemistry lettres*. 2006;16(2):430-432.
 - Green DR, Kroemer G. The pathophysiology of mitochondrial cell death. *Science*. 2004;305(5684):626-629.
 - Guicciardi ME, Gores GJ. Life and death by death receptors. *The FASEB Journal*. 2009;23(6):1625-1637.
 - Hadden MK, Hill SA, Davenport J, Matts RL, Blagg BS. Synthesis and evaluation of Hsp90 inhibitors that contain the 1, 4-naphthoquinone scaffold. *Bioorganic & medicinal chemistry*. 2009;17(2):634-640.
 - Hajdúch M, Havlíček L, Veselý J, Novotný R, Mihál V, Strnad M. Synthetic Cyclin Dependent Kinase Inhibitors. Drug resistance in leukemia and lymphoma III: Springer; 1999;vol.457 341-53.
 - Halperin I, Ma B, Wolfson H, Nussinov R. Principles of docking: An overview of search algorithms and a guide to scoring functions. *Proteins: structure, function, and genetic*. 2002;47(4):409-443.
 - Hanahan D, Weinberg RA. Hallmarks of cancer: the next generation. *Cell*. 2011;144(5):646-674.
 - Hande KR. Clinical applications of anticancer drugs targeted to topoisomerase II. *Biochimica Biophysica Acta*. 1998;1400(1-3):173-84.
 - Haschek WM, Rousseaux CG, Wallig MA, Bolon B, Ochoa R. Haschek and Rousseaux's handbook of toxicologic pathology. 3 ed: Academic Press;

- 2013pp 4054.
- Hennessy BT, Smith DL, Ram PT, Lu Y, Mills GB. Exploiting the PI3K/AKT pathway for cancer drug discovery. *Nature reviews drug discovery*. 2005;4(12):988-1004.
 - Hirsch F, Varella-Garcia M, Cappuzzo F. Predictive value of EGFR and HER2 overexpression in advanced non-small-cell lung cancer. *Oncogene*. 2009;28(1):32-37.
 - Hu X, Wang D, Tong Y, Tong L, Wang X, Zhu L, et al. Design, Synthesis, and Evaluation of Ribose-Modified Anilino-pyrimidine Derivatives as EGFR Tyrosine Kinase Inhibitors. *Frontiers in chemistry*. 2017;5:101-110.
 - Huang G, Dong JY, Zhang QJ, Meng QQ, Zhao HR, Zhu BQ, et al. Discovery and synthesis of sulfur-containing 6-substituted 5,8-dimethoxy-1,4-naphthoquinone oxime derivatives as new and potential anti-MDR cancer agents. *European Journal of Medicinal Chemistry*. 2019;165:160-71.
 - Hubbard SR, Till JH. Protein tyrosine kinase structure and function. *Annual review of biochemistry*. 2000;69(1):373-398.
 - Hueso-Falcon I, Amesty A, Anaissi-Afonso L, Lorenzo-Castrillejo I, Machin F, Estevez-Braun A. Synthesis and biological evaluation of naphthoquinone-coumarin conjugates as topoisomerase II inhibitors. *Bioorganic & medicinal chemistry*. 2017; 27(3):484-489.
 - Hussain H, Specht S, Sarite SR, Hoerauf A, Krohn K. New quinoline-5, 8-dione and hydroxynaphthoquinone derivatives inhibit a chloroquine resistant Plasmodium falciparum strain. *European journal of medicinal chemistry*. 2012;54:936-942.
 - Iglesias VS, Giuranno L, Dubois LJ, Theys J, Vooijs M. Drug resistance in non-small cell lung cancer: a potential for NOTCH targeting. *Frontiers in oncology*. 2018;8(24):267-290.
 - Iijima K, Yoshikawa N, Connolly DT, Nakamura H. Human mesangial cells and peripheral blood mononuclear cells produce vascular permeability factor. *Kidney international*. 1993;44(5):959-966.
 - Inagaki R, Ninomiya M, Tanaka K, Koketsu M. Synthesis, characterization, and antileukemic properties of naphthoquinone derivatives of lawsone. *ChemMedChem*. 2015;10(8):1413-1423.
 - Ingham PW, McMahon AP. Hedgehog signaling in animal development: paradigms and principles. *Genes & development*. 2001;15(23):3059-3087.
 - Itakura J, Ishiwata T, Shen B, Kornmann M, Korc M. Concomitant over-expression of vascular endothelial growth factor and its receptors in pancreatic cancer. *International journal of cancer*. 2000;85(1):27-34.
 - Jain AN. Scoring functions for protein-ligand docking. *Current protein and peptide science*. 2006;7(5):407-420.
 - Johann L, Belorgey D, Huang HH, Day L, Chesse M, Becker K, et al. Synthesis and evaluation of 1,4-naphthoquinone ether derivatives as SmTGR inhibitors and new anti-schistosomal drugs. *FEBS J*. 2015;282(16):3199-3217.
 - Johnson CD, Esquela-Kerscher A, Stefani G, Byrom M, Kelnar K, Ovcharenko D, et al. The let-7 microRNA represses cell proliferation pathways in human cells. *Cancer research*. 2007;67(16):7713-7722.

- Jordan JD, Landau EM, Iyengar R. Signaling networks: the origins of cellular multitasking. *Cell*. 2000;103(2):193-200.
- Jorgensen WL. The many roles of computation in drug discovery. *Science*. 2004;303(5665):1813-1818.
- Kacmaz A, Deniz NG, Aydinli SG, Sayil C, Onay-Ucar E, Mertoglu E, et al. Synthesis and antiproliferative evaluation of some 1, 4-naphthoquinone derivatives against human cervical cancer cells. *Open Chemistry*. 17(1):337-345.
- Kano MR, Komuta Y, Iwata C, Oka M, Shirai Yt, Morishita Y, et al. Comparison of the effects of the kinase inhibitors imatinib, sorafenib, and transforming growth factor- β receptor inhibitor on extravasation of nanoparticles from neovasculature. *Cancer science*. 2009;100(1):173-180.
- Kar S, Lefterov IM, Wang M, Lazo JS, Scott CN, Wilcox CS, et al. Binding and inhibition of Cdc25 phosphatases by vitamin K analogues. *Biochemistry*. 2003;42(35):10490-10497.
- Kaur J, Tikoo K. p300/CBP dependent hyperacetylation of histone potentiates anticancer activity of gefitinib nanoparticles. *Biochimica biophysica acta (BBA)-molecular cell research*. 2013;1833(5):1028-1040.
- Kawano M, Mabuchi S, Kishimoto T, Hisamatsu T, Matsumoto Y, Sasano T, et al. Combination treatment with trabectedin and irinotecan or topotecan has synergistic effects against ovarian clear cell carcinoma cells. *International Journal of Gynecologic Cancer*. 2014;24(5):829-837.
- Ke LD, Shi Y-X, Im S-A, Chen X, Yung WA. The relevance of cell proliferation, vascular endothelial growth factor, and basic fibroblast growth factor production to angiogenesis and tumorigenicity in human glioma cell lines. *Clinical cancer research*. 2000;6(6):2562-2572.
- Kerr JF, Wyllie AH, Currie AR. Apoptosis: a basic biological phenomenon with wideranging implications in tissue kinetics. *British journal of cancer*. 1972;26(4):239-257.
- Kharb R, Shahar Yar M, C Sharma P. New insights into chemistry and anti-infective potential of triazole scaffold. *Current medicinal chemistry*. 2011;18(21):3265-97.
- Khraiwesh MH, Lee CM, Brandy Y, Akinboye ES, Berhe S, Gittens G, et al. Antitrypanosomal activities and cytotoxicity of some novel imido-substituted 1,4-naphthoquinone derivatives. *Archives of pharmacal research*. 2012;35(1):27-33.
- Kim BH, Yoo J, Park S-H, Jung J-K, Cho H, Chung Y. Synthesis and evaluation of antitumor activity of novel 1, 4-naphthoquinone derivatives (IV). *Archives of pharmacal research*. 2006;29(2):123-130.
- Kim HJ, Mun JY, Chun YJ, Choi KH, Ham SW, Kim MY. Effects of a naphthoquinone analog on tumor growth and apoptosis induction. *Archives of pharmacal research*. 2003;26(5):405-410.
- Kintzel PE, Dorr RT. Anticancer drug renal toxicity and elimination: dosing guidelines for altered renal function. *Cancer treatment reviews*. 1995;21(1):33-64.
- Koch S, Claesson-Welsh L. Signal transduction by vascular endothelial growth factor receptors. *Cold spring harbor perspectives in medicine*.

- 2012;2(7): 6502-6523.
- Kraus S, Naor Z, Seger R. Intracellular signaling pathways mediated by the gonadotropin-releasing hormone (GnRH) receptor. *Archives of medical research*. 2001;32(6):499-509.
 - Kumagai Y, Shinkai Y, Miura T, Cho AK. The chemical biology of naphthoquinones and its environmental implications. *Annual review of pharmacology and toxicology*. 2012;52:221-247.
 - Kumari S, Badana AK, Malla R. Reactive oxygen species: a key constituent in cancer survival. *Biomarker insights*. 2018;13:1-9.
 - Kuribayashi K, Mayes PA, El-Deiry WS. What are caspases 3 and 7 doing upstream of the mitochondria. *Cancer biology & therapy*. 2006;5(7):763-765.
 - Lahiani A, Yavin E, Lazarovici P. The molecular basis of toxins' interactions with intracellular signaling via discrete portals. *Toxins*. 2017;9(3):107-116.
 - Lara L, Moreira C, Calvet C, Lechuga G, Souza R, Bourguignon S, et al. Efficacy of 2-hydroxy-3-phenylsulfanylmethyl-[1,4]-naphthoquinone derivatives against different *Trypanosoma cruzi* discrete type units: Identification of a promising hit compound. *European journal of medicinal chemistry*. 2018;144:572-581.
 - Lee SH, Lee JR, Lunde CS, Kubo I. In vitro antifungal susceptibilities of *Candida albicans* and other fungal pathogens to polygodial, a sesquiterpene dialdehyde. *Planta medica*. 1999;65(03):204-208.
 - Lemmon MA, Schlessinger J. Cell signaling by receptor tyrosine kinases. *Cell*. 2010;141(7):1117-1134.
 - Lengauer T, Rarey M. Computational methods for biomolecular docking. *Current opinion in structural biology*. 1996;6(3):402-4026.
 - Leung DW, Cachianes G, Kuang W-J, Goeddel DV, Ferrara N. Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science*. 1989;246(4935):1306-1309.
 - Levitzki A, Gazit A. Tyrosine kinase inhibition: an approach to drug development. *Science*. 1995;267(5205):1782-1788.
 - Li J, Tan Jz, Chen Ll, Zhang J, Shen X, Mei Cl, et al. Design, synthesis and antitumor evaluation of a new series of N-substituted-thiourea derivatives I. *Acta pharmacologica sinica*. 2006;27(9):1259-1271.
 - Li K, Yang K, Zheng L, Li Y, Wang Q, Lin R, et al. Anti-acute myeloid leukemia activity of 2-chloro-3-alkyl-1,4-naphthoquinone derivatives through inducing mtDNA damage and GSH depletion. *Bioorganic & medicinal chemistry*. 2018;26(14):4191-200.
 - Lien J-C, Huang L-J, Teng C-M, Wang J-P, Kuo S-C. Synthesis of 2-alkoxy 1, 4-naphthoquinone derivatives as antiplatelet, antiinflammatory, and antiallergic agents. *Chemical and pharmaceutical bulletin*. 2002;50(5):672-674.
 - Lienhardt C, Vernon A, Raviglione MC. New drugs and new regimens for the treatment of tuberculosis: review of the drug development pipeline and implications for national programmes. *Current opinion in pulmonary medicine*. 2010;16(3):186-193.

- Lin L, He X-P, Xu Q, Chen G-R, Xie J. Synthesis of β -C-glycopyranosyl-1, 4-naphthoquinone derivatives and their cytotoxic activity. *Carbohydrate research*. 2008;343(4):773-779.
- Lin Y-m, Lu G-p, Cai C, Yi W-b. An odorless thia-Michael addition using Bunte salts as thiol surrogates. *RSC advances*. 2015;5(34):27107-27111.
- Liu C, Shen GN, Luo YH, Piao XJ, Jiang XY, Meng LQ, et al. Novel 1,4-naphthoquinone derivatives induce apoptosis via ROS-mediated p38/MAPK, Akt and STAT3 signaling in human hepatoma Hep3B cells. *International Journal of Biochemistry & Cell Biology*. 2018;96:9-19.
- Lockshin RA, CM W. *JInsect Physiol*. 1965;11
- Longati P, Comoglio PM, Bardelli A. Receptor tyrosine kinases as therapeutic targets the model of the MET oncogene. *Current drug targets*. 2001;2(1):41-55.
- Lu X, Altharawi A, Gut J, Rosenthal PJ, Long TE. 1, 4-Naphthoquinone cations as antiplasmodial agents: hydroxy-, acyloxy-, and alkoxy-substituted analogues. *ACS medicinal chemistry letters*. 2012;3(12):1029-1033.
- MacCallum DE, Melville J, Frame S, Watt K, Anderson S, Gianella-Borradori A, et al. Seliciclib (CYC202, R-roscovitine) induces cell death in multiple myeloma cells by inhibition of RNA polymerase II-dependent transcription and down-regulation of Mcl-1. *Cancer research*. 2005;65(12):5399-5407.
- Mandal B, Mittal NK, Balabathula P, Thoma LA, Wood GC. Development and in vitro evaluation of core-shell type lipid-polymer hybrid nanoparticles for the delivery of erlotinib in non-small cell lung cancer. *European Journal of Pharmaceutical Sciences*. 2016;81:162-171.
- Mandal S, Moudgil Mn, Mandal SK. Rational drug design. *European journal of pharmacology*. 2009;625(1-3):90-100.
- Markovic ZM, Harhaji-Trajkovic LM, Todorovic-Markovic BM, Kepić DP, Arsikin KM, Jovanović SP, et al. In vitro comparison of the photothermal anticancer activity of graphene nanoparticles and carbon nanotubes. *Biomaterials*. 2011;32(4):1121-1129.
- Mazumdar-Shaw K. Leveraging affordable innovation to tackle India's healthcare challenge. *IIMB Management review*. 2018;30(1):37-50.
- McNamara CE, Larsen L, Perry NB, Harper JL, Berridge MV, Chia EW, et al. Anti-inflammatory Sesquiterpene-quinones from the New Zealand Sponge *Dysidea* cf. *c ristagalli*. *Journal of natural products*. 2005;68(9):1431-1433.
- Mercurio AM. VEGF/Neuropilin Signaling in Cancer Stem Cells. *International journal of molecular sciences*. 2019;20(3):490-502.
- Mertenskoetter T, Kaptur P. Update on microbicide research and development-seeking new HIV prevention tools for women. *European journal of medical research*. 2011;16(1):1-6.
- Milackova I, Prnova MS, Majekova M, Sotnikova R, Stasko M, Kovacikova L, et al. 2-Chloro-1, 4-naphthoquinone derivative of quercetin as an inhibitor of aldose reductase and anti-inflammatory agent. *Journal of enzyme inhibition and medicinal chemistry*. 2015;30(1):107-113.
- Mitchell T, Cherry M. Fragment-based drug design. *Innov Pharm Technol*.

- 2005;16:34-6.
- Moghaddam SM, Amini A, Morris DL, Pourgholami MH. Significance of vascular endothelial growth factor in growth and peritoneal dissemination of ovarian cancer. *Cancer and Metastasis Reviews*. 2012;31(1-2):143-162.
 - Neet K, Hunter T. Vertebrate non-receptor protein-tyrosine kinase families. *Genes to Cells*. 1996;1(2):147-169.
 - Nussbaumer S, Bonnabry P, Veuthey J-L, Fleury-Souverain S. Analysis of anticancer drugs: a review. *Talanta*. 2011;85(5):2265-2289.
 - O'brien P. Molecular mechanisms of quinone cytotoxicity. *Chemico-biological interactions*. 1991;80(1):1-41.
 - Ol'ga AZ, Goryunov LI, Troshkova NM, Ovchinnikova LP, Shteingarts VD, Nevinsky GA. Cytotoxicity of new n-butylamino and sulfur-containing derivatives of polyfluorinated 1, 4-naphthoquinone. *European journal of medicinal chemistry*. 2010;45(1):270-274.
 - Ol'ga DZ, Ovchinnikova LP, Goryunov LI, Troshkova NM, Shteingarts VD, Nevinsky GA. Cytotoxicity of new alkylamino-and phenylamino-containing polyfluorinated derivatives of 1, 4-naphthoquinone. *European journal of medicinal chemistry*. 2010;45(6):2321-2326.
 - Ōmura S, Asami Y, Crump A. Staurosporine: New lease of life for parent compound of today's novel and highly successful anti-cancer drugs. *The Journal of antibiotics*. 2018;71(8):688-701.
 - Osada T, Hartman ZC, Wei J, Lei G, Hobeika AC, Gwin WR, et al. Polyfunctional anti-human epidermal growth factor receptor 3 (anti-HER3) antibodies induced by HER3 vaccines have multiple mechanisms of antitumor activity against therapy resistant and triple negative breast cancers. *Breast Cancer Research*. 2018;20(1):90-104.
 - Ott M, Gogvadze V, Orrenius S, Zhivotovsky B. Mitochondria, oxidative stress and cell death. *Apoptosis*. 2007;12(5):913-22.
 - Pàez-Ribes M, Allen E, Hudock J, Takeda T, Okuyama H, Viñals F, et al. Antiangiogenic therapy elicits malignant progression of tumors to increased local invasion and distant metastasis. *Cancer cell*. 2009;15(3):220-231.
 - Park M-J, Sheng R, Silkov A, Jung D-J, Wang Z-G, Xin Y, et al. SH2 domains serve as lipid-binding modules for pTyr-signaling proteins. *Molecular cell*. 2016;62(1):7-20.
 - Pawson T. Specificity in signal transduction: from phosphotyrosine-SH2 domain interactions to complex cellular systems. *Cell*. 2004;116(2):191-203.
 - Pérez-Sacau E, Díaz-Peñate RG, Estévez-Braun A, Ravelo AG, García-Castellano JM, Pardo L, et al. Synthesis and pharmacophore modeling of naphthoquinone derivatives with cytotoxic activity in human promyelocytic leukemia HL-60 cell line. *Journal of medicinal chemistry*. 2007;50(4):696-706.
 - Perfézou M, Turner A, Merkoçi A. Cancer detection using nanoparticle-based sensors. *Chemical society reviews*. 2012;41(7):2606-2622.
 - Perrot-Appianat M, Di Benedetto M. Autocrine functions of VEGF in breast tumor cells: adhesion, survival, migration and invasion. *Cell adhesion & migration*. 2012;6(6):547-553.

- Perše M, Cerar A, Injac R, Štrukelj B. N-methylnitrosourea Induced Breast Cancer in Rat, the Histopathology of the Resulting Tumours and its Drawbacks as a Model. *Pathology & Oncology Research*. 2009;15(1):115-121.
- Peter ME, Heufelder AE, Hengartner MO. Advances in apoptosis research. *Proceedings of the National Academy of Sciences*. 1997;94(24):12736-12737.
- Petrelli F, Borgonovo K, Cabiddu M, Ghilardi M, Barni S. Cetuximab and panitumumab in KRAS wild-type colorectal cancer: a meta-analysis. *International journal of colorectal disease*. 2011;26(7):823-833.
- Pieretti S, Haanstra J, Mazet M, Perozzo R, Bergamini C, Prati F, et al. Naphthoquinone derivatives exert their antitrypanosomal activity via a multi-target mechanism. *PLoS Neglected Tropical Diseases*. 2013;7:1-12.
- Pingaew R, Prachayasittikul V, Worachartcheewan A, Nantasenamat C, Prachayasittikul S, Ruchirawat S, et al. Novel 1, 4-naphthoquinone-based sulfonamides: Synthesis, QSAR, anticancer and antimalarial studies. *European journal of medicinal chemistry*. 2015;103:446-459.
- Pink R, Hudson A, Mouriès M-A, Bendig M. Opportunities and challenges in antiparasitic drug discovery. *Nature reviews drug discovery*. 2005;4(9):727-740.
- Pistrutto G, Trisciuglio D, Ceci C, Garufi A, D'Orazi G. Apoptosis as anticancer mechanism: function and dysfunction of its modulators and targeted therapeutic strategies. *Aging (Albany NY)*. 2016;8(4):603-619.
- Prachayasittikul V, Pingaew R, Worachartcheewan A, Nantasenamat C, Prachayasittikul S, Ruchirawat S, et al. Synthesis, anticancer activity and QSAR study of 1, 4-naphthoquinone derivatives. *European journal of medicinal chemistry*. 2014;84:247-263.
- Pradidphol N, Kongkathip N, Sittikul P, Boonyalai N, Kongkathip B. First synthesis and anticancer activity of novel naphthoquinone amides. *European journal of medicinal chemistry*. 2012;49:253-270.
- Rahman M, Laurent S, Tawil N, Yahia L, Mahmoudi M. Protein-nanoparticle interactions: Springer; series in biology; 2013.
- Rathish IG, Javed K, Ahamad S, Bano S, Alam MS, Akhter M, Pillai KK, Ovais S, Samim M. Synthesis and evaluation of anticancer activity of some novel 6-aryl-2-(sulfamylphenyl)-pyridazin-3(2H)one. *European journal of medicinal chemistry*. 2012;49:304-309.
- Ravichandiran P, Premnath D, Vasanthkumar S. Synthesis, molecular docking and antibacterial evaluation of new 1,4-naphthoquinone derivatives contains carbazole-6,11-dione moiety. *Journal of Chemical Biology*. 2014;7(3):93-101.
- Ravichandiran P, Subramaniyan SA, Kim SY, Kim JS, Park BH, Shim KS, et al. Synthesis and Anticancer Evaluation of 1,4-Naphthoquinone Derivatives Containing a Phenylaminosulfanyl Moiety. *ChemMedChem*. 2019;14(5):532-544.
- Rehm J, Mathers C, Popova S, Thavorncharoensap M, Teerawattananon Y, Patra J. Global burden of disease and injury and economic cost attributable to alcohol use and alcohol-use disorders. *The lancet*. 2009;373(9682):2223-

- 2233.
- Reichardt LF, Tomaselli KJ. Extracellular matrix molecules and their receptors: functions in neural development. *Annual review of neuroscience*. 1991;14(1):531-570.
 - Ren D, Tu H-C, Kim H, Wang GX, Bean GR, Takeuchi O, et al. BID, BIM, and PUMA are essential for activation of the BAX-and BAK-dependent cell death program. *Science*. 2010;330(6009):1390-1393.
 - Ren J, Jia X, Zhao Y, Shi W, Lu J, Zhang Y, et al. The RIP3-RIP1-NF- κ B signaling axis is dispensable for necroptotic cells to elicit cross-priming of CD8+ T cells. *Cellular & molecular immunology*. 2017;14(7):639-642.
 - Renehan AG, Booth C, Potten CS. What is apoptosis, and why is it important? Education and debate. *Bmj*. 2001;322(7301):1536-1538.
 - Roa-Linares VC, Miranda-Brand Y, Tangarife-Castaño V, Ochoa R, García PA, Castro M, et al. Anti-Herpetic, Anti-Dengue and Antineoplastic Activities of Simple and Heterocycle-Fused Derivatives of Terpenyl-1, 4-Naphthoquinone and 1, 4-Anthraquinone. *Molecules*. 2019;24(7):1279-1295.
 - Robertson TA, Varani G. An all-atom, distance-dependent scoring function for the prediction of protein–DNA interactions from structure. *Proteins: Structure, Function, and Bioinformatics*. 2007;66(2):359-374.
 - Roessler E, Belloni E, Gaudenz K, Jay P, Berta P, Scherer SW, et al. Mutations in the human Sonic Hedgehog gene cause holoprosencephaly. *Nature genetics*. 1996;14(3):357-360.
 - Rubin LL, de Sauvage FJ. Targeting the Hedgehog pathway in cancer. *Nature reviews Drug discovery*. 2006;5(12):1026-1033.
 - Salas C, Tapia RA, Ciudad K, Armstrong V, Orellana M, Kemmerling U, et al. Trypanosoma cruzi: Activities of lapachol and α - and β -lapachone derivatives against epimastigote and trypomastigote forms. *Bioorganic & medicinal chemistry*. 2008;16(2):668-674.
 - Salmon-Chemin L, Buisine E, Yardley V, Kohler S, Debreu M-A, Landry V, et al. 2- and 3-substituted 1, 4-naphthoquinone derivatives as subversive substrates of trypanothione reductase and lipoamide dehydrogenase from trypanosoma c ruzi: synthesis and correlation between redox cycling activities and in vitro cytotoxicity. *Journal of medicinal chemistry*. 2001;44(4):548-565.
 - Samant BS, Chakaingesu C. Novel naphthoquinone derivatives: synthesis and activity against human African trypanosomiasis. *Bioorganic & Medicinal Chemistry Letters*. 2013;23(5):1420-1423.
 - Sankar MM, Singh J, Diana SCA, Singh S. Molecular characterization of Mycobacterium tuberculosis isolates from North Indian patients with extrapulmonary tuberculosis. *Tuberculosis*. 2013;93(1):75-83.
 - Scapin G. Structural biology in drug design: selective protein kinase inhibitors. *Drug discovery today*. 2002;7(11):601-611.
 - Schieber M, Chandel NS. ROS function in redox signaling and oxidative stress. *Current biology*. 2014;24(10):R453-R62.
 - Schlessinger J. Autoinhibition control.(Signal Transduction). *Science*. 2003;300(5620):750-753.

- Schneider G, Fechner U. Computer-based de novo design of drug-like molecules. *Nature reviews Drug discovery*. 2005;4(8):649-663.
- Segurado AAC, Santi SMD, Shiroma M. In vivo and in vitro Plasmodium falciparum resistance to chloroquine, amodiaquine and quinine in the Brazilian Amazon. *Revista do instituto de medicina tropical de são paulo*. 1997;39:85-90.
- Senderowicz AM. The cell cycle as a target for cancer therapy: basic and clinical findings with the small molecule inhibitors flavopiridol and UCN-01. *The oncologist*. 2002;7(3):12-19.
- Senthilraja P, Kathiresan K. In vitro cytotoxicity MTT assay in Vero, HepG2 and MCF-7 cell lines study of Marine Yeast. *Journal of Applied Pharmaceutical Science*. 2015;5(3):80-84.
- Seshacharyulu P, Ponnusamy MP, Haridas D, Jain M, Ganti AK, Batra SK. Targeting the EGFR signaling pathway in cancer therapy. *Expert opinion on therapeutic targets*. 2012;16(1):15-31.
- Sever R, Brugge JS. Signal transduction in cancer. *Cold Spring Harbor perspectives in medicine*. 2015;5(4):6098-6119.
- Shapiro GI, Harper JW. Anticancer drug targets: cell cycle and checkpoint control. *The Journal of clinical investigation*. 1999;104(12):1645-53.
- Sharma P, Jha AB, Dubey RS, Pessarakli M. Reactive oxygen species, oxidative damage, and antioxidative defense mechanism in plants under stressful conditions. *Journal of botany*. 2012;27:1-26.
- Shoichet BK, Kuntz ID, Bodian DL. Molecular docking using shape descriptors. *Journal of computational chemistry*. 1992;13(3):380-397.
- Siwak DR, Carey M, Hennessy BT, Nguyen CT, McGahren Murray MJ, Nolden L, et al. Targeting the epidermal growth factor receptor in epithelial ovarian cancer: current knowledge and future challenges. *Journal of oncology*. 2010;2010:1-20.
- Snuderl M, Batista A, Kirkpatrick ND, de Almodovar CR, Riedemann L, Walsh EC, et al. Targeting placental growth factor/neuropilin 1 pathway inhibits growth and spread of medulloblastoma. *Cell*. 2013;152(5):1065-1076.
- Souza-Silva F, do Nascimento SB, Bourguignon SC, Pereira BAS, Carneiro PF, da Silva WS, et al. Evidences for leishmanicidal activity of the naphthoquinone derivative epoxy- α -lapachone. *Experimental parasitology*. 2014;147:81-84.
- Sreelatha T, Kandhasamy S, Dinesh R, Shruthy S, Shweta S, Mukesh D, et al. Synthesis and SAR study of novel anticancer and antimicrobial naphthoquinone amide derivatives. *Bioorganic & medicinal chemistry letters*. 2014;24(15):3647-3651.
- Srinivas G, Babykutty S, Sathiadevan PP, Srinivas P. Molecular mechanism of emodin action: transition from laxative ingredient to an antitumor agent. *Medicinal research reviews*. 2007;27(5):591-608.
- Stanton MJ, Dutta S, Zhang H, Polavaram NS, Leontovich AA, Hönscheid P, et al. Autophagy control by the VEGF-C/NRP-2 axis in cancer and its implication for treatment resistance. *Cancer research*. 2013;73(1):160-171.
- Steeg PS. Tumor metastasis: mechanistic insights and clinical challenges.

- Nature medicine. 2006;12(8):895-904.
- Stein GS, Pardee AB. Cell cycle and growth control: biomolecular regulation and cancer: John Wiley & Sons; 2004, 2nd edition.
 - Steuer CE, Khuri FR, Ramalingam SS. The next generation of epidermal growth factor receptor tyrosine kinase inhibitors in the treatment of lung cancer. *Cancer*. 2015;121(8):1-6.
 - Takahashi S. Vascular endothelial growth factor (VEGF), VEGF receptors and their inhibitors for antiangiogenic tumor therapy. *Biological and pharmaceutical bulletin*. 2011;34(12):1785-1788.
 - Tanaka N, Ishihara M, Kitagawa M, Harada H, Kimura T, Matsuyama T, et al. Cellular commitment to oncogene-induced transformation or apoptosis is dependent on the transcription factor IRF-1. *Cell*. 1994;77(6):829-839.
 - Tandon VK, Chhor RB, Singh RV, Rai S, Yadav DB. Design, synthesis and evaluation of novel 1, 4-naphthoquinone derivatives as antifungal and anticancer agents. *Bioorganic & medicinal chemistry letters*. 2004;14(5):1079-1083.
 - Tandon VK, Singh RV, Yadav DB. Synthesis and evaluation of novel 1,4-naphthoquinone derivatives as antiviral, antifungal and anticancer agents. *Bioorganic & medicinal chemistry letters*. 2004;14(11):2901-2904.
 - Tandon VK, Yadav DB, Singh RV, Vaish M, Chaturvedi AK, Shukla PK. Synthesis and biological evaluation of novel 1, 4-naphthoquinone derivatives as antibacterial and antiviral agents. *Bioorganic & medicinal chemistry letters*. 2005;15(14):3463-3466.
 - Tanrikulu Y, Schneider G. Pseudoreceptor models in drug design: bridging ligand-and receptor-based virtual screening. *Nature reviews drug discovery*. 2008;7(8):667-677.
 - Tonks NK, Neel BG. From form to function: signaling by protein tyrosine phosphatases. *Cell*. 1996;87(3):365-368.
 - Tsai C-J, Nussinov R. Emerging allosteric mechanism of EGFR activation in physiological and pathological contexts. *Biophysical Journal*. 2019;117(19):1-9.
 - Tseng C-L, Su W-Y, Yen K-C, Yang K-C, Lin F-H. The use of biotinylated-EGF-modified gelatin nanoparticle carrier to enhance cisplatin accumulation in cancerous lungs via inhalation. *Biomaterials*. 2009;30(20):3476-3485.
 - Turpin JA. Considerations and development of topical microbicides to inhibit the sexual transmission of HIV. *Expert opinion on investigational drugs*. 2002;11(8):1077-1097.
 - Varjosalo M, Taipale J. Hedgehog: functions and mechanisms. *Genes & development*. 2008;22(18):2454-2472.
 - Verheul H, Hoekman K, Luyckx-de Bakker S, Eekman CA, Folman CC, Broxterman HJ, et al. Platelet: transporter of vascular endothelial growth factor. *Clinical cancer research*. 1997;3(12):2187-2190.
 - Vulpetti A, Hommel U, Landrum G, Lewis R, Dalvit C. Design and NMR-based screening of LEF, a library of chemical fragments with different local environment of fluorine. *Journal of the american chemical society*. 2009;131(36):12949-12959.
 - Wakharde AA, Awad AH, Bhagat A, SM K. Synergistic Activation of

- Doxorubicin against Cancer: A Review. *American journal of clinical microbiology and antimicrobials*. 2018 1(2):1009-1015.
- Wallach D, Kang T-B, Dillon CP, Green DR. Programmed necrosis in inflammation: Toward identification of the effector molecules. *Science*. 2016;352(6281):2154-2163.
 - Wang B, Miao Z, Wang J, Chen R, Zhang X. Synthesis and biological evaluation of novel naphthoquinone fused cyclic aminoalkylphosphonates and aminoalkylphosphonic monoester. *Amino acids*. 2008;35(2):463-468.
 - Wang M, Ong L-LS, Dauwels J, Asada HH. Automated tracking and quantification of angiogenic vessel formation in 3D microfluidic devices. *PloS one*. 2017;12(11):0186465.
 - Wang S-H, Lo C-Y, Gwo Z-H, Lin H-J, Chen L-G, Kuo C-D, et al. Synthesis and biological evaluation of lipophilic 1, 4-naphthoquinone derivatives against human cancer cell lines. *Molecules*. 2015;20(7):11994-2015.
 - Wang X, Yan J, Wang M, Liu M, Zhang J, Chen L, et al. Synthesis and three-dimensional quantitative structure-activity relationship study of quinazoline derivatives containing a 1,3,4-oxadiazole moiety as efficient inhibitors against *Xanthomonas axonopodis* pv. *citri*. *Mol divers*. 2018;22(4):791-802.
 - Wedam SB, Low JA, Yang SX, Chow CK, Choyke P, Danforth D, et al. Antiangiogenic and antitumor effects of bevacizumab in patients with inflammatory and locally advanced breast cancer. *Journal of clinical oncology*. 2006;24(5):769-777.
 - Wilson L, Jordan M. New microtubule/tubulin-targeted anticancer drugs and novel chemotherapeutic strategies. *Journal of chemotherapy*. 2004;16:83-85.
 - Wolber G, Seidel T, Bendix F, Langer T. Molecule-pharmacophore superpositioning and pattern matching in computational drug design. *Drug discovery today*. 2008;13(1-2):23-29.
 - Wolkenberg SE, Boger DL. Mechanisms of in situ activation for DNA-targeting antitumor agents. *Chemical reviews*. 2002;102(7):2477-2496.
 - Wong RS. Apoptosis in cancer: from pathogenesis to treatment. *Journal of experimental & clinical cancer research*. 2011;30(1):87-101.
 - Wu B, Barile E, K De S, Wei J, Purves A, Pellecchia M. High-throughput screening by nuclear magnetic resonance (HTS by NMR) for the identification of PPIs antagonists. *Current topics in medicinal chemistry*. 2015;15(20):2032-2042.
 - Xie W, Chipman JG, Robertson DL, Erikson R, Simmons DL. Expression of a mitogen-responsive gene encoding prostaglandin synthase is regulated by mRNA splicing. *Proceedings of the national academy of sciences*. 1991;88(7):2692-2696.
 - Xu K, Xiao Z, Tang YB, Huang L, Chen C-H, Ohkoshi E, et al. Design and synthesis of naphthoquinone derivatives as antiproliferative agents and 20S proteasome inhibitors. *Bioorganic & medicinal chemistry letters*. 2012;22(8):2772-2774.
 - Yamashita M, Kaneko M, Iida A, Tokuda H, Nishimura K. Stereoselective

- synthesis and cytotoxicity of a cancer chemopreventive naphthoquinone from *Tabebuia avellaneda*. *Bioorganic & medicinal chemistry letters*. 2007;17(23):6417-6420.
- Yang J, Yan J, Liu B. Targeting VEGF/VEGFR to modulate antitumor immunity. *Frontiers in immunology*. 2018;9:978-988.
 - Yarden Y, Sliwkowski MX. Untangling the ErbB signalling network. *Nature reviews molecular cell biology*. 2001;2(2):127-137.
 - Yates LR, Campbell PJ. Evolution of the cancer genome. *Nature reviews genetics*. 2012;13(11):795-806.
 - Zauhar RJ, Moyna G, Tian L, Li Z, Welsh WJ. Shape signatures: a new approach to computer-aided ligand-and receptor-based drug design. *Journal of medicinal chemistry*. 2003;46(26):5674-5690.
 - Zhang Q, Dong J, Cui J, Huang G, Meng Q, Li S. Cytotoxicity of Synthesized 1,4-Naphthoquinone Oxime Derivatives on Selected Human Cancer Cell Lines. *Chem pharm bull (tokyo)*. 2018;66(6):612-619.
 - Zhang Y, Chen X, Gueydan C, Han J. Plasma membrane changes during programmed cell deaths. *Cell research*. 2018;28(1):9-21.
 - Ziemska J, Solecka J. Tyrosine kinase, aurora kinase and leucine aminopeptidase as attractive drug targets in anticancer therapy-characterisation of their inhibitors. *Roczniki Państwowego Zakładu Higieny*. 2016;67(4):329-331.