

References

- ❖ Abdolmajid, E., Kharazi, H., Chalaki, M., Khojasteh, M., Haghghat, S., Attar, F., & Falahati, M. (2019). Titanium oxide nanoparticles fabrication, hemoglobin interaction, white blood cells cytotoxicity, and antibacterial studies. *Journal of Biomolecular Structure and Dynamics*, 37(11), 3007-3017.
- ❖ Ajayan, P. M., & Zhou, O. Z. (2001). Applications of carbon nanotubes. In *Carbon nanotubes* (pp. 391-425). Springer, Berlin, Heidelberg.
- ❖ Alam, A. M., Park, B. Y., Ghouri, Z. K., Park, M., & Kim, H. Y. (2015). Synthesis of carbon quantum dots from cabbage with down-and up-conversion photoluminescence properties: excellent imaging agent for biomedical applications. *Green Chemistry*, 17(7), 3791-3797.
- ❖ Albani, D., Vilé, G., Mitchell, S., Witte, P. T., Almora-Barrios, N., Verel, R., & Pérez-Ramírez, J. (2016). Ligand ordering determines the catalytic response of hybrid palladium nanoparticles in hydrogenation. *Catalysis Science & Technology*, 6(6), 1621-1631.
- ❖ Arsiya, F., Sayadi, M. H., & Sobhani, S. (2017). Green synthesis of palladium nanoparticles using *Chlorella vulgaris*. *Materials Letters*, 186, 113-115.
- ❖ Atchudan, R., Perumal, S., Karthikeyan, D., Pandurangan, A., & Lee, Y. R. (2015). Synthesis and characterization of graphitic mesoporous carbon using metal-metal oxide by chemical vapor deposition method. *Microporous and Mesoporous Materials*, 215, 123-132.
- ❖ Baker, S. N., & Baker, G. A. (2010). Luminescent carbon nanodots: emergent nanolights. *Angewandte Chemie International Edition*, 49(38), 6726-6744.
- ❖ Bandi, R., Devulapalli, N. P., Dadigala, R., Gangapuram, B. R., & Guttena, V. (2018). Facile conversion of toxic cigarette butts to N, S-codoped carbon dots and their application in fluorescent film, security ink, bioimaging, sensing and logic gate operation. *ACS omega*, 3(10), 13454-13466.
- ❖ Bano, D., Chandra, S., Yadav, P. K., Singh, V. K., & Hasan, S. H. (2020). Off-on detection of glutathione based on the nitrogen, sulfur codoped carbon quantum dots@ MnO₂ nano-composite in human lung cancer cells and blood serum. *Journal of Photochemistry and Photobiology A: Chemistry*, 112558.
- ❖ Bano, D., Kumar, V., Singh, V. K., & Hasan, S. H. (2018). Green synthesis of fluorescent carbon quantum dots for the detection of mercury (II) and glutathione. *New Journal of Chemistry*, 42(8), 5814-5821.

- ❖ Bano, D., Kumar, V., Singh, V. K., Chandra, S., Singh, D. K., Yadav, P. K., & Hasan, S. H. (2018). A facile and simple strategy for the synthesis of label free carbon quantum dots from the latex of *Euphorbia milii* and its peroxidase-mimic activity for the naked eye detection of glutathione in a human blood serum. *ACS Sustainable Chemistry & Engineering*, 7(2), 1923-1932.
- ❖ Bao, L., Zhang, Z. L., Tian, Z. Q., Zhang, L., Liu, C., Lin, Y., & Pang, D. W. (2011). Electrochemical tuning of luminescent carbon nanodots: from preparation to luminescence mechanism. *Advanced materials*, 23(48), 5801-5806.
- ❖ Baptista, F. R., Belhout, S. A., Giordani, S., & Quinn, S. J. (2015). Recent developments in carbon nanomaterial sensors. *Chemical Society Reviews*, 44(13), 4433-4453.
- ❖ Barceloux, D. G., & Barceloux, D. (1999). Cobalt. *Journal of Toxicology: Clinical Toxicology*, 37(2), 201-216.
- ❖ Bendale, Y., Bendale, V., & Paul, S. (2017). Evaluation of cytotoxic activity of platinum nanoparticles against normal and cancer cells and its anticancer potential through induction of apoptosis. *Integrative medicine research*, 6(2), 141-148.
- ❖ Bian, W., Ma, J., Liu, Q., Wei, Y., Li, Y., Dong, C., & Shuang, S. (2014). A novel phosphorescence sensor for Co^{2+} ion based on Mn-doped ZnS quantum dots. *Luminescence*, 29(2), 151-157.
- ❖ Blagoi, G. (2004). Fluorescence resonance energy transfer (FRET) based sensors for bioanalysis.
- ❖ Bolton, O., Lee, K., Kim, H. J., Lin, K. Y., & Kim, J. (2011). Activating efficient phosphorescence from purely organic materials by crystal design. *Nature chemistry*, 3(3), 205.
- ❖ Breslow, R. (1995). Biomimetic chemistry and artificial enzymes: catalysis by design. *Accounts of Chemical Research*, 28(3), 146-153.
- ❖ Briscoe, J., Marinovic, A., Sevilla, M., Dunn, S., & Titirici, M. (2015). Biomass-derived carbon quantum dot sensitizers for solid-state nanostructured solar cells. *Angewandte Chemie International Edition*, 54(15), 4463-4468.
- ❖ Cai, Q. Y., Li, J., Ge, J., Zhang, L., Hu, Y. L., Li, Z. H., & Qu, L. B. (2015). A rapid fluorescence “switch-on” assay for glutathione detection by using carbon dots– MnO_2 nanocomposites. *Biosensors and Bioelectronics*, 72, 31-36.
- ❖ Cano, M., Núñez-Lozano, R., Lumbreras, R., González-Rodríguez, V., Delgado-García, A., Jiménez-Hoyuela, J.M. and de la Cueva-Méndez, G. (2017) Partial PEGylation of superparamagnetic iron oxide nanoparticles thinly coated with amine-

- silane as a source of ultrastable tunable nanosystems for biomedical applications. *Nanoscale* 9(2), 812-822.
- ❖ Cao, L., Wang, X., Meziani, M. J., Lu, F., Wang, H., Luo, P. G., & Xie, S. Y. (2007). Carbon dots for multiphoton bioimaging. *Journal of the American Chemical Society*, 129(37), 11318-11319.
 - ❖ Cao, L., Wang, X., Meziani, M. J., Lu, F., Wang, H., Luo, P. G., & Xie, S. Y. (2007). Carbon dots for multiphoton bioimaging. *Journal of the American Chemical Society*, 129(37), 11318-11319.
 - ❖ Cayuela, A., Benítez-Martínez, S., & Soriano, M. L. (2016). Carbon nanotools as sorbents and sensors of nanosized objects: the third way of analytical nanoscience and nanotechnology. *TrAC Trends in Analytical Chemistry*, 84, 172-180.
 - ❖ Chan, W. C., & Nie, S. (1998). Quantum dot bioconjugates for ultrasensitive nonisotopic detection. *Science*, 281(5385), 2016-2018.
 - ❖ Chandra, S., Laha, D., Pramanik, A., Ray Chowdhuri, A., Karmakar, P., & Sahu, S. K. (2016). Synthesis of highly fluorescent nitrogen and phosphorus doped carbon dots for the detection of Fe³⁺ ions in cancer cells. *Luminescence*, 31(1), 81-87.
 - ❖ Chandra, S., Patra, P., Pathan, S. H., Roy, S., Mitra, S., Layek, A., & Goswami, A. (2013). Luminescent S-doped carbon dots: an emergent architecture for multimodal applications. *Journal of Materials Chemistry B*, 1(18), 2375-2382.
 - ❖ Chandra, S., Singh, V. K., Yadav, P. K., Bano, D., Kumar, V., Pandey, V. K., & Hasan, S. H. (2019). Mustard seeds derived fluorescent carbon quantum dots and their peroxidase-like activity for colorimetric detection of H₂O₂ and ascorbic acid in a real sample. *Analytica chimica acta*, 1054, 145-156.
 - ❖ Charbgoon, F., Ahmad, M.B. and Darroudi, M. (2017) Cerium oxide nanoparticles: green synthesis and biological applications. *International journal of nanomedicine* 12, 1401.
 - ❖ Chaudhari, N. K., Song, M. Y., & Yu, J. S. (2014). Heteroatom-doped highly porous carbon from human urine. *Scientific reports*, 4, 5221.
 - ❖ Chen, D., Tang, L., & Li, J. (2010). Graphene-based materials in electrochemistry. *Chemical Society Reviews*, 39(8), 3157-3180.
 - ❖ Chen, X. and Mao, S.S. (2007) Titanium dioxide nanomaterials: synthesis, properties, modifications, and applications. *Chem. Rev* 107(7), 2891-2959.
 - ❖ Chen, Y., Shang, P., Dong, Y., & Chi, Y. (2017). Regulating the overlap between the absorption spectrum of metal ion-chromogenic agent and the emission spectrum of

- carbon-based dots to improve the sensing performance for metal ions. *Sensors and Actuators B: Chemical*, 242, 1210-1215.
- ❖ Chen, Z., Wang, Z., Chen, J., Wang, S., & Huang, X. (2012). Sensitive and selective detection of glutathione based on resonance light scattering using sensitive gold nanoparticles as colorimetric probes. *Analyst*, 137(13), 3132-3137.
 - ❖ Coates, J. P., & Shelley, P. H. (2006). Infrared spectroscopy in process analysis. *Encyclopedia of Analytical Chemistry: Applications, Theory and Instrumentation*.
 - ❖ Contino, A., Maccarrone, G., Zimbone, M., Reitano, R., Musumeci, P., Calcagno, L., & Oliveri, I. P. (2016). Tyrosine capped silver nanoparticles: a new fluorescent sensor for the quantitative determination of copper (II) and cobalt (II) ions. *Journal of colloid and interface science*, 462, 216-222.
 - ❖ Contino, A., Maccarrone, G., Zimbone, M., Reitano, R., Musumeci, P., Calcagno, L., & Oliveri, I. P. (2016). Tyrosine capped silver nanoparticles: a new fluorescent sensor for the quantitative determination of copper (II) and cobalt (II) ions. *Journal of colloid and interface science*, 462, 216-222.
 - ❖ Dahl, J.A., Maddux, B.L. and Hutchison, J.E. (2007) Toward greener nanosynthesis. *Chemical reviews* 107(6), 2228-2269
 - ❖ Daniel, M.-C. and Astruc, D. (2004) Gold nanoparticles: assembly, supramolecular chemistry, quantum-size-related properties, and applications toward biology, catalysis, and nanotechnology. *Chemical reviews* 104(1), 293-346.
 - ❖ De Caluwé, E., Halamová, K., & Van Damme, P. (2010). *Adansonia digitata* L.—A review of traditional uses, phytochemistry and pharmacology. *Afrika focus*, 23(1).
 - ❖ de Matos, R. A., da Silva Cordeiro, T., Samad, R. E., Vieira Jr, N. D., & Courrol, L. C. (2011). Green synthesis of stable silver nanoparticles using *Euphorbia milii* latex. *Colloids and Surfaces A: Physicochemical and Engineering Aspects*, 389(1-3), 134-137.
 - ❖ Demchenko, A. P., & Dekaliuk, M. O. (2013). Novel fluorescent carbonic nanomaterials for sensing and imaging. *Methods and applications in fluorescence*, 1(4), 042001.
 - ❖ Demchenko, A. P., & Dekaliuk, M. O. (2013). Novel fluorescent carbonic nanomaterials for sensing and imaging. *Methods and applications in fluorescence*, 1(4), 042001.
 - ❖ Deng, R., Xie, X., Vendrell, M., Chang, Y. T., & Liu, X. (2011). Intracellular glutathione detection using MnO₂-nanosheet-modified upconversion nanoparticles. *Journal of the American Chemical Society*, 133(50), 20168-20171.

- ❖ Deng, Y., Zhao, D., Chen, X., Wang, F., Song, H., & Shen, D. (2013). Long lifetime pure organic phosphorescence based on water soluble carbon dots. *Chemical Communications*, 49(51), 5751-5753.
- ❖ Dey, S., Chithaiah, P., Belawadi, S., Biswas, K., & Rao, C. N. R. (2014). New methods of synthesis and varied properties of carbon quantum dots with high nitrogen content. *Journal of Materials Research*, 29(3), 383-391.
- ❖ Di, W., Zhang, X., & Qin, W. (2017). Single-layer MnO₂ nanosheets for sensitive and selective detection of glutathione by a colorimetric method. *Applied Surface Science*, 400, 200-205.
- ❖ Ding, H., Wei, J. S., & Xiong, H. M. (2014). Nitrogen and sulfur co-doped carbon dots with strong blue luminescence. *Nanoscale*, 6(22), 13817-13823.
- ❖ Ding, H., Zhang, P., Wang, T. Y., Kong, J. L., & Xiong, H. M. (2014). Nitrogen-doped carbon dots derived from polyvinyl pyrrolidone and their multicolor cell imaging. *Nanotechnology*, 25(20), 205604.
- ❖ Dong, H., Gao, W., Yan, F., Ji, H., & Ju, H. (2010). Fluorescence resonance energy transfer between quantum dots and graphene oxide for sensing biomolecules. *Analytical chemistry*, 82(13), 5511-5517.
- ❖ Dong, Y., Chen, C., Zheng, X., Gao, L., Cui, Z., Yang, H., & Li, C. M. (2012). One-step and high yield simultaneous preparation of single-and multi-layer graphene quantum dots from CX-72 carbon black. *Journal of Materials Chemistry*, 22(18), 8764-8766.
- ❖ Dong, Y., Pang, H., Yang, H. B., Guo, C., Shao, J., Chi, Y., & Yu, T. (2013). Carbon-based dots co-doped with nitrogen and sulfur for high quantum yield and excitation-independent emission. *Angewandte Chemie International Edition*, 52(30), 7800-7804.
- ❖ Dong, Y., Pang, H., Yang, H. B., Guo, C., Shao, J., Chi, Y., & Yu, T. (2013). Carbon-based dots co-doped with nitrogen and sulfur for high quantum yield and excitation-independent emission. *Angewandte Chemie International Edition*, 52(30), 7800-7804.
- ❖ Dong, Y., Wang, R., Li, G., Chen, C., Chi, Y., & Chen, G. (2012). Polyamine-functionalized carbon quantum dots as fluorescent probes for selective and sensitive detection of copper ions. *Analytical chemistry*, 84(14), 6220-6224.
- ❖ Dong, Y., Wang, R., Li, G., Chen, C., Chi, Y., & Chen, G. (2012). Polyamine-functionalized carbon quantum dots as fluorescent probes for selective and sensitive detection of copper ions. *Analytical chemistry*, 84(14), 6220-6224.

-
- ❖ Dubas, S.T. and Pimpan, V. (2008) Humic acid assisted synthesis of silver nanoparticles and its application to herbicide detection. *Materials Letters* 62(17), 2661-2663.
 - ❖ Eda, G., Lin, Y. Y., Mattevi, C., Yamaguchi, H., Chen, H. A., Chen, I. S., & Chhowalla, M. (2010). Blue photoluminescence from chemically derived graphene oxide. *Advanced materials*, 22(4), 505-509.
 - ❖ Edison, T. N. J. I., Atchudan, R., Shim, J. J., Kalimuthu, S., Ahn, B. C., & Lee, Y. R. (2016). Turn-off fluorescence sensor for the detection of ferric ion in water using green synthesized N-doped carbon dots and its bio-imaging. *Journal of Photochemistry and Photobiology B: Biology*, 158, 235-242.
 - ❖ El-Naggar, M. E., Shaheen, T. I., Zaghloul, S., El-Rafie, M. H., & Hebeish, A. (2016). Antibacterial activities and UV protection of the in situ synthesized titanium oxide nanoparticles on cotton fabrics. *Industrial & Engineering Chemistry Research*, 55(10), 2661-2668.
 - ❖ Estrela, J. M., Ortega, A., & Obrador, E. (2006). Glutathione in cancer biology and therapy. *Critical reviews in clinical laboratory sciences*, 43(2), 143-181.
 - ❖ Farooq, M. U., Novosad, V., Rozhkova, E. A., Wali, H., Ali, A., Fateh, A. A., & Wang, Z. (2018). Gold nanoparticles-enabled efficient dual delivery of anticancer therapeutics to HeLa cells. *Scientific reports*, 8(1), 1-12.
 - ❖ Favier, I., Pla, D., & Gómez, M. (2019). Palladium Nanoparticles in Polyols: Synthesis, Catalytic Couplings, and Hydrogenations. *Chemical Reviews*, 120(2), 1146-1183.
 - ❖ Feng, J., Huang, P., Shi, S., Deng, K. Y., & Wu, F. Y. (2017). Colorimetric detection of glutathione in cells based on peroxidase-like activity of gold nanoclusters: a promising powerful tool for identifying cancer cells. *Analytica chimica acta*, 967, 64-69.
 - ❖ Feng, J., Huang, P., Shi, S., Deng, K. Y., & Wu, F. Y. (2017). Colorimetric detection of glutathione in cells based on peroxidase-like activity of gold nanoclusters: a promising powerful tool for identifying cancer cells. *Analytica chimica acta*, 967, 64-69.
 - ❖ Feynman, R.P. (1960) There's plenty of room at the bottom. *Engineering and science* 23(5), 22-36.
 - ❖ Filippo, E., Serra, A., Buccolieri, A. and Manno, D. (2010) Green synthesis of silver nanoparticles with sucrose and maltose: morphological and structural characterization. *Journal of Non-Crystalline Solids* 356(6), 344-350.

- ❖ Flora, S. J. S., Mittal, M., & Mehta, A. (2008). Heavy metal induced oxidative stress & its possible reversal by chelation therapy. *Indian Journal of Medical Research*, 128(4), 501.
- ❖ Gabor, G., & Walt, D. R. (1991). Sensitivity enhancement of fluorescent pH indicators by inner filter effects. *Analytical chemistry*, 63(8), 793-796.
- ❖ Gajalakshmi, S., Iswarya, V., Ashwini, R., Divya, G., Mythili, S., & Sathiavelu, A. (2012). Evaluation of heavy metals in medicinal plants growing in Vellore District. *European Journal of Experimental Biology*, 2(5), 1457-1461.
- ❖ Gao, C., Zhu, H., Chen, J., & Qiu, H. (2017). Facile synthesis of enzyme functional metal-organic framework for colorimetric detecting H₂O₂ and ascorbic acid. *Chinese Chemical Letters*, 28(5), 1006-1012.
- ❖ Ge, J., Cai, R., Chen, X., Wu, Q., Zhang, L., Jiang, Y., & Tan, W. (2019). Facile approach to prepare HSA-templated MnO₂ nanosheets as oxidase mimic for colorimetric detection of glutathione. *Talanta*, 195, 40-45.
- ❖ Geys, J., Nemmar, A., Verbeken, E., Smolders, E., Ratoi, M., Hoylaerts, M. F., & Hoet, P. H. (2008). Acute toxicity and prothrombotic effects of quantum dots: impact of surface charge. *Environmental health perspectives*, 116(12), 1607-1613.
- ❖ Gokus, T., Nair, R. R., Bonetti, A., Bohmler, M., Lombardo, A., Novoselov, K. S., & Hartschuh, A. (2009). Making graphene luminescent by oxygen plasma treatment. *ACS nano*, 3(12), 3963-3968.
- ❖ Gonçalves, H., Jorge, P. A., Fernandes, J. R. A., & da Silva, J. C. E. (2010). Hg (II) sensing based on functionalized carbon dots obtained by direct laser ablation. *Sensors and Actuators B: Chemical*, 145(2), 702-707.
- ❖ Gong, X., Lu, W., Paa, M. C., Hu, Q., Wu, X., Shuang, S., & Choi, M. M. (2015). Facile synthesis of nitrogen-doped carbon dots for Fe³⁺ sensing and cellular imaging. *Analytica chimica acta*, 861, 74-84.
- ❖ Gong, Y. J., Zhang, X. B., Chen, Z., Yuan, Y., Jin, Z., Mei, L., & Yu, R. Q. (2012). An efficient rhodamine thiospirolactam-based fluorescent probe for detection of Hg²⁺ in aqueous samples. *Analyst*, 137(4), 932-938.
- ❖ Gong, Y., Yu, B., Yang, W., & Zhang, X. (2016). Phosphorus, and nitrogen co-doped carbon dots as a fluorescent probe for real-time measurement of reactive oxygen and nitrogen species inside macrophages. *Biosensors and Bioelectronics*, 79, 822-828.
- ❖ Guo, Y., Chen, Y., Cao, F., Wang, L., Wang, Z., & Leng, Y. (2017). Hydrothermal synthesis of nitrogen and boron doped carbon quantum dots with yellow-green

- emission for sensing Cr (VI), anti-counterfeiting and cell imaging. *RSC advances*, 7(76), 48386-48393.
- ❖ Gupta, A., Verma, N. C., Khan, S., & Nandi, C. K. (2016). Carbon dots for naked eye colorimetric ultrasensitive arsenic and glutathione detection. *Biosensors and Bioelectronics*, 81, 465-472.
 - ❖ Gupta, A.K. and Gupta, M. (2005) Synthesis and surface engineering of iron oxide nanoparticles for biomedical applications. *Biomaterials* 26(18), 3995-4021.
 - ❖ Gutknecht, J. (1981). Inorganic mercury (Hg 2+) transport through lipid bilayer membranes. *The Journal of Membrane Biology*, 61(1), 61-66.
 - ❖ Han, B., Yuan, J., & Wang, E. (2009). Sensitive and selective sensor for biothiols in the cell based on the recovered fluorescence of the CdTe quantum dots– Hg (II) system. *Analytical chemistry*, 81(13), 5569-5573.
 - ❖ Han, Z., Zhang, H., He, L., Pan, S., Liu, H., & Hu, X. (2019). One-pot hydrothermal synthesis of nitrogen and sulfur co-doped carbon dots and their application for sensitive detection of curcumin and temperature. *Microchemical Journal*, 146, 300-308.
 - ❖ Handy, R.D., Owen, R. and Valsami-Jones, E. (2008) The ecotoxicology of nanoparticles and nanomaterials: current status, knowledge gaps, challenges, and future needs. *Ecotoxicology* 17(5), 315-325.
 - ❖ He, D., Zheng, C., Wang, Q., He, C., Lee, Y. I., Wu, L., & Hou, X. (2015). Dielectric barrier discharge-assisted one-pot synthesis of carbon quantum dots as fluorescent probes for selective and sensitive detection of hydrogen peroxide and glucose. *Talanta*, 142, 51-56.
 - ❖ He, H., Li, H., Mohr, G., Kovacs, B., Werner, T., & Wolfbeis, O. S. (1993). Novel type of ion-selective fluorosensor based on the inner filter effect: an optrode for potassium. *Analytical Chemistry*, 65(2), 123-127.
 - ❖ He, L., Bao, Z., Zhang, K., Yang, D., Sheng, B., Huang, R., & Zhang, C. (2018). Ratiometric determination of copper (II) using dually emitting Mn (II)-doped ZnS quantum dots as a fluorescent probe. *Microchimica Acta*, 185(11), 511.
 - ❖ Heyduk, T. (2002). Measuring protein conformational changes by FRET/LRET. *Current opinion in biotechnology*, 13(4), 292-296.
 - ❖ Hink, M. A., Bisseling, T., & Visser, A. J. (2002). Imaging protein-protein interactions in living cells. *Plant molecular biology*, 50(6), 871-883.

- ❖ Hsu, P. C., & Chang, H. T. (2012). Synthesis of high-quality carbon nanodots from hydrophilic compounds: role of functional groups. *Chemical communications*, 48(33), 3984-3986.
- ❖ Hsu, P. C., Chen, P. C., Ou, C. M., Chang, H. Y., & Chang, H. T. (2013). Extremely high inhibition activity of photoluminescent carbon nanodots toward cancer cells. *Journal of Materials Chemistry B*, 1(13), 1774-1781.
- ❖ Hu, B., Cao, X., & Zhang, P. (2013). Selective colorimetric detection of glutathione based on quasi-stable gold nanoparticles assembly. *New Journal of Chemistry*, 37(12), 3853-3856.
- ❖ Hu, C., Yu, C., Li, M., Wang, X., Yang, J., Zhao, Z., & Qiu, J. (2014). Chemically tailoring coal to fluorescent carbon dots with tuned size and their capacity for Cu (II) detection. *Small*, 10(23), 4926-4933.
- ❖ Hu, R., Liu, T., Zhang, X. B., Huan, S. Y., Wu, C., Fu, T., & Tan, W. (2014). Multicolor fluorescent biosensor for multiplexed detection of DNA. *Analytical chemistry*, 86(10), 5009-5016.
- ❖ Huang, C. C., Yang, Z., Lee, K. H., & Chang, H. T. (2007). Synthesis of highly fluorescent gold nanoparticles for sensing mercury (II). *Angewandte Chemie International Edition*, 46(36), 6824-6828.
- ❖ Huang, H., Weng, Y., Zheng, L., Yao, B., Weng, W., & Lin, X. (2017). Nitrogen-doped carbon quantum dots as fluorescent probe for “off-on” detection of mercury ions, L-cysteine and iodide ions. *Journal of colloid and interface science*, 506, 373-378.
- ❖ Huang, H., Xu, Y., Tang, C. J., Chen, J. R., Wang, A. J., & Feng, J. J. (2014). Facile and green synthesis of photoluminescent carbon nanoparticles for cellular imaging. *New Journal of Chemistry*, 38(2), 784-789.
- ❖ Huang, Q., Hu, S., Zhang, H., Chen, J., He, Y., Li, F., & Lin, Y. (2013). Carbon dots and chitosan composite film based biosensor for the sensitive and selective determination of dopamine. *Analyst*, 138(18), 5417-5423.
- ❖ Ismail, E., Khenfouch, M., Dhlamini, M., Dube, S. and Maaza, M. (2017) Green palladium and palladium oxide nanoparticles synthesized via *Aspalathus linearis* natural extract. *Journal of Alloys and Compounds* 695, 3632-3638.
- ❖ Jeong, C. J., Roy, A. K., Kim, S. H., Lee, J. E., Jeong, J. H., In, I., & Park, S. Y. (2014). Fluorescent carbon nanoparticles derived from natural materials of mango fruit for bio-imaging probes. *Nanoscale*, 6(24), 15196-15202.

-
- ❖ Jiang, C., Guan, Z., Lim, S. Y. R., Polavarapu, L., & Xu, Q. H. (2011). Two-photon ratiometric sensing of Hg²⁺ by using cysteine functionalized Ag nanoparticles. *Nanoscale*, 3(8), 3316-3320.
 - ❖ Jiang, H., Chen, F., Lagally, M. G., & Denes, F. S. (2010). New strategy for synthesis and functionalization of carbon nanoparticles. *Langmuir*, 26(3), 1991-1995.
 - ❖ Jiang, Y., Song, N., Wang, C., Pinna, N., & Lu, X. (2017). A facile synthesis of Fe₃O₄/nitrogen-doped carbon hybrid nanofibers as a robust peroxidase-like catalyst for the sensitive colorimetric detection of ascorbic acid. *Journal of Materials Chemistry B*, 5(27), 5499-5505.
 - ❖ Jing, N., Tian, M., Wang, Y., & Zhang, Y. (2019). Nitrogen-doped carbon dots synthesized from acrylic acid and ethylenediamine for simple and selective determination of cobalt ions in aqueous media. *Journal of Luminescence*, 206, 169-175.
 - ❖ Ju, J., Zhang, R., & Chen, W. (2016). Photochemical deposition of surface-clean silver nanoparticles on nitrogen-doped graphene quantum dots for sensitive colorimetric detection of glutathione. *Sensors and Actuators B: Chemical*, 228, 66-73.
 - ❖ Jv, Y., Li, B., & Cao, R. (2010). Positively-charged gold nanoparticles as peroxidase mimic and their application in hydrogen peroxide and glucose detection. *Chemical Communications*, 46(42), 8017-8019.
 - ❖ Kai, K., Yoshida, Y., Kageyama, H., Saito, G., Ishigaki, T., Furukawa, Y., & Kawamata, J. (2008). Room-temperature synthesis of manganese oxide monosheets. *Journal of the American Chemical Society*, 130(47), 15938-15943.
 - ❖ Kailasa, S. K., Hasan, N., & Wu, H. F. (2012). Identification of multiply charged proteins and amino acid clusters by liquid nitrogen assisted spray ionization mass spectrometry. *Talanta*, 97, 539-549.
 - ❖ Kamidate, T., & Watanabe, H. (1996). Peroxidase-catalysed luminol chemiluminescence method for the determination of glutathione. *Talanta*, 43(10), 1733-1738.
 - ❖ Kaur, N., Mehta, A., Mishra, A., Chaudhary, S., Rawat, M., & Basu, S. (2019). Amphiphilic carbon dots derived by cationic surfactant for selective and sensitive detection of metal ions. *Materials Science and Engineering: C*, 95, 72-77.
 - ❖ Khan, W. U., Wang, D., & Wang, Y. (2018). Highly green emissive nitrogen-doped carbon dots with excellent thermal stability for bioimaging and solid-state LED. *Inorganic chemistry*, 57(24), 15229-15239.

- ❖ Kim, J. H., Gibb, H. J., Howe, P. D., & World Health Organization. (2006). Cobalt and inorganic cobalt compounds.
- ❖ Kim, J., Park, J., Kim, H., Singha, K., & Kim, W. J. (2013). Transfection and intracellular trafficking properties of carbon dot-gold nanoparticle molecular assembly conjugated with PEI-pDNA. *Biomaterials*, *34*(29), 7168-7180.
- ❖ Koch, C., Ovid'ko, I., Seal, S. and Veprek, S. (2007) Structural nanocrystalline materials: fundamentals and applications, Cambridge University Press.
- ❖ Kong, D., Yan, F., Han, Z., Xu, J., Guo, X., & Chen, L. (2016). Cobalt (II) ions detection using carbon dots as an sensitive and selective fluorescent probe. *RSC advances*, *6*(72), 67481-67487.
- ❖ Kumar, R., da Silva, E. T., Singh, R. K., Savu, R., Alaferdov, A. V., Fonseca, L. C., & Alves, O. L. (2018). Microwave-assisted synthesis of palladium nanoparticles intercalated nitrogen doped reduced graphene oxide and their electrocatalytic activity for direct-ethanol fuel cells. *Journal of colloid and interface science*, *515*, 160-171.
- ❖ Kumar-Krishnan, S., Hernandez-Rangel, A., Pal, U., Ceballos-Sanchez, O., Flores-Ruiz, F. J., Prokhorov, E., & Meyyappan, M. (2016). Surface functionalized halloysite nanotubes decorated with silver nanoparticles for enzyme immobilization and biosensing. *Journal of Materials Chemistry B*, *4*(15), 2553-2560.
- ❖ Lakowicz, J. R. (Ed.). (2013). *Principles of fluorescence spectroscopy*. Springer science & business media.
- ❖ Lambert, M., Leven, B. A., & Green, R. M. (2000). New methods of cleaning up heavy metal in soils and water. *Environmental science and technology briefs for citizens*. Kansas State University, Manhattan, KS.
- ❖ Larson, D. R., Zipfel, W. R., Williams, R. M., Clark, S. W., Bruchez, M. P., Wise, F. W., & Webb, W. W. (2003). Water-soluble quantum dots for multiphoton fluorescence imaging in vivo. *Science*, *300*(5624), 1434-1436.
- ❖ Laurent, S., Forge, D., Port, M., Roch, A., Robic, C., Vander Elst, L. and Muller, R.N. (2008) Magnetic iron oxide nanoparticles: synthesis, stabilization, vectorization, physicochemical characterizations, and biological applications. *Chemical reviews* *108*(6), 2064-2110.
- ❖ Lee, P. T., Goncalves, L. M., & Compton, R. G. (2015). Electrochemical determination of free and total glutathione in human saliva samples. *Sensors and Actuators B: Chemical*, *221*, 962-968.
- ❖ Li, C. L., Huang, C. C., Periasamy, A. P., Roy, P., Wu, W. C., Hsu, C. L., & Chang, H. T. (2015). Synthesis of photoluminescent carbon dots for the detection of cobalt ions. *RSC Advances*, *5*(3), 2285-2291.

- ❖ Li, H., He, X., Liu, Y., Yu, H., Kang, Z., & Lee, S. T. (2011). Synthesis of fluorescent carbon nanoparticles directly from active carbon via a one-step ultrasonic treatment. *Materials Research Bulletin*, 46(1), 147-151.
- ❖ Li, H., He, X., Liu, Y., Yu, H., Kang, Z., & Lee, S. T. (2011). Synthesis of fluorescent carbon nanoparticles directly from active carbon via a one-step ultrasonic treatment. *Materials Research Bulletin*, 46(1), 147-151.
- ❖ Li, H., Kang, Z., Liu, Y., & Lee, S. T. (2012). Carbon nanodots: synthesis, properties and applications. *Journal of materials chemistry*, 22(46), 24230-24253.
- ❖ Li, H., Sun, C., Vijayaraghavan, R., Zhou, F., Zhang, X., & MacFarlane, D. R. (2016). Long lifetime photoluminescence in N, S co-doped carbon quantum dots from an ionic liquid and their applications in ultrasensitive detection of pesticides. *Carbon*, 104, 33-39.
- ❖ Li, H., Zhang, Y., Wang, L., Tian, J., & Sun, X. (2011). Nucleic acid detection using carbon nanoparticles as a fluorescent sensing platform. *Chemical Communications*, 47(3), 961-963.
- ❖ Li, J., Wu, Y., Song, F., Wei, G., Cheng, Y., & Zhu, C. (2012). A highly selective and sensitive polymer-based OFF-ON fluorescent sensor for Hg²⁺ detection incorporating salen and perylenyl moieties. *Journal of Materials Chemistry*, 22(2), 478-482.
- ❖ Li, L., Yu, B., & You, T. (2015). Nitrogen and sulfur co-doped carbon dots for highly selective and sensitive detection of Hg (II) ions. *Biosensors and Bioelectronics*, 74, 263-269.
- ❖ Li, M., Wu, X., Wang, Y., Li, Y., Zhu, W., & James, T. D. (2014). A near-infrared colorimetric fluorescent chemodosimeter for the detection of glutathione in living cells. *Chemical Communications*, 50(14), 1751-1753.
- ❖ Li, T., Zhou, Y., Sun, J., Tang, D., Guo, S., & Ding, X. (2011). Ultrasensitive detection of mercury (II) ion using CdTe quantum dots in sol-gel-derived silica spheres coated with calix [6] arene as fluorescent probes. *Microchimica Acta*, 175(1-2), 113.
- ❖ Li, W., Yue, Z., Wang, C., Zhang, W., & Liu, G. (2013). An absolutely green approach to fabricate carbon nanodots from soya bean grounds. *RSC advances*, 3(43), 20662-20665.
- ❖ Li, W., Zhang, Z., Kong, B., Feng, S., Wang, J., Wang, L., & Zhao, D. (2013). Simple and green synthesis of nitrogen-doped photoluminescent carbonaceous nanospheres for bioimaging. *Angewandte Chemie International Edition*, 52(31), 8151-8155.

- ❖ Li, X., Lau, S. P., Tang, L., Ji, R., & Yang, P. (2014). Sulphur doping: a facile approach to tune the electronic structure and optical properties of graphene quantum dots. *Nanoscale*, 6(10), 5323-5328.
- ❖ Li, X., Wang, H., Shimizu, Y., Pyatenko, A., Kawaguchi, K., & Koshizaki, N. (2010). Preparation of carbon quantum dots with tunable photoluminescence by rapid laser passivation in ordinary organic solvents. *Chemical Communications*, 47(3), 932-934.
- ❖ Li, X., Wang, H., Shimizu, Y., Pyatenko, A., Kawaguchi, K., & Koshizaki, N. (2010). Preparation of carbon quantum dots with tunable photoluminescence by rapid laser passivation in ordinary organic solvents. *Chemical Communications*, 47(3), 932-934.
- ❖ Li, Y., Liu, Y., Shang, X., Chao, D., Zhou, L., & Zhang, H. (2018). Highly sensitive and selective detection of Fe³⁺ by utilizing carbon quantum dots as fluorescent probes. *Chemical Physics Letters*, 705, 1-6.
- ❖ Li, Y., Wang, Z., Li, X., Yin, T., Bian, K., Gao, F. and Gao, D. (2017) Facile synthesis of bacitracin-templated palladium nanoparticles with superior electrocatalytic activity. *Journal of Power Sources* 341, 183-191.
- ❖ Li, Y., Zhao, Y., Cheng, H., Hu, Y., Shi, G., Dai, L., & Qu, L. (2012). Nitrogen-doped graphene quantum dots with oxygen-rich functional groups. *Journal of the American Chemical Society*, 134(1), 15-18.
- ❖ Liao, S., Zhu, F., Zhao, X., Yang, H., & Chen, X. (2018). A reusable P, N-doped carbon quantum dot fluorescent sensor for cobalt ion. *Sensors and Actuators B: Chemical*, 260, 156-164.
- ❖ Lim, S. Y., Shen, W., & Gao, Z. (2015). Carbon quantum dots and their applications. *Chemical Society Reviews*, 44(1), 362-381.
- ❖ Lin, L., & Zhang, S. (2012). Creating high yield water soluble luminescent graphene quantum dots via exfoliating and disintegrating carbon nanotubes and graphite flakes. *Chemical communications*, 48(82), 10177-10179.
- ❖ Lin, L., Rong, M., Lu, S., Song, X., Zhong, Y., Yan, J., & Chen, X. (2015). A facile synthesis of highly luminescent nitrogen-doped graphene quantum dots for the detection of 2, 4, 6-trinitrophenol in aqueous solution. *Nanoscale*, 7(5), 1872-1878.
- ❖ Lin, P., Chen, J. W., Chang, L. W., Wu, J. P., Redding, L., Chang, H., & Kuo, Y. C. (2008). Computational and ultrastructural toxicology of a nanoparticle, Quantum Dot 705, in mice. *Environmental science & technology*, 42(16), 6264-6270.
- ❖ Lin, T., Zhong, L., Guo, L., Fu, F., & Chen, G. (2014). Seeing diabetes: visual detection of glucose based on the intrinsic peroxidase-like activity of MoS₂ nanosheets. *Nanoscale*, 6(20), 11856-11862.

- ❖ Lin, Y. H., & Tseng, W. L. (2010). Ultrasensitive sensing of Hg²⁺ and CH₃Hg⁺ based on the fluorescence quenching of lysozyme type VI-stabilized gold nanoclusters. *Analytical chemistry*, 82(22), 9194-9200.
- ❖ Lison, D. (1996). Human toxicity of cobalt-containing dust and experimental studies on the mechanism of interstitial lung disease (hard metal disease). *Critical reviews in toxicology*, 26(6), 585-616.
- ❖ Liu, H., He, Z., Jiang, L. P., & Zhu, J. J. (2015). Microwave-assisted synthesis of wavelength-tunable photoluminescent carbon nanodots and their potential applications. *ACS applied materials & interfaces*, 7(8), 4913-4920.
- ❖ Liu, H., Ye, T., & Mao, C. (2007). Fluorescent carbon nanoparticles derived from candle soot. *Angewandte Chemie International Edition*, 46(34), 6473-6475.
- ❖ Liu, H., Ye, T., & Mao, C. (2007). Fluorescent carbon nanoparticles derived from candle soot. *Angewandte chemie*, 119(34), 6593-6595.
- ❖ Liu, J., Bao, C., Zhong, X., Zhao, C., & Zhu, L. (2010). Highly selective detection of glutathione using a quantum-dot-based OFF-ON fluorescent probe. *Chemical communications*, 46(17), 2971-2973.
- ❖ Liu, J., Li, J., Xu, L., Qiao, Y., & Chen, J. (2017). Facile synthesis of N, B-doped carbon dots and their application for multisensor and cellular imaging. *Industrial & Engineering Chemistry Research*, 56(14), 3905-3912.
- ❖ Liu, R., Wu, D., Liu, S., Koynov, K., Knoll, W., & Li, Q. (2009). An aqueous route to multicolor photoluminescent carbon dots using silica spheres as carriers. *Angewandte Chemie International Edition*, 48(25), 4598-4601.
- ❖ Liu, S., Tian, J., Wang, L., Luo, Y., & Sun, X. (2012). A general strategy for the production of photoluminescent carbon nitride dots from organic amines and their application as novel peroxidase-like catalysts for colorimetric detection of H₂O₂ and glucose. *Rsc Advances*, 2(2), 411-413.
- ❖ Liu, S., Tian, J., Wang, L., Zhang, Y., Qin, X., Luo, Y., & Sun, X. (2012). Hydrothermal treatment of grass: a low-cost, green route to nitrogen-doped, carbon-rich, photoluminescent polymer nanodots as an effective fluorescent sensing platform for label-free detection of Cu (II) ions. *Advanced materials*, 24(15), 2037-2041.
- ❖ Liu, X., Wang, Q., Zhang, Y., Zhang, L., Su, Y., & Lv, Y. (2013). Colorimetric detection of glutathione in human blood serum based on the reduction of oxidized TMB. *New Journal of Chemistry*, 37(7), 2174-2178.

- ❖ Liu, X., Wei, S., Diao, Q., Ma, P., Xu, L., Xu, S., & Wang, X. (2017). Hydrothermal synthesis of N-doped carbon dots for selective fluorescent sensing and cellular imaging of cobalt (II). *Microchimica Acta*, 184(10), 3825-3831.
- ❖ Long, Y., Jiang, D., Zhu, X., Wang, J., & Zhou, F. (2009). Trace Hg²⁺ analysis via quenching of the fluorescence of a CdS-encapsulated DNA nanocomposite. *Analytical chemistry*, 81(7), 2652-2657.
- ❖ Lu, L., Sun, R., Chen, R., Hui, C.-K., Ho, C.-M., Luk, J.M., Lau, G. and Che, C.-M. (2008) Silver nanoparticles inhibit hepatitis B virus replication. *Antiviral therapy* 13(2), 253.
- ❖ Lu, P.-J., Huang, S.-C., Chen, Y.-P., Chiueh, L.-C. and Shih, D.Y.-C. (2015) Analysis of titanium dioxide and zinc oxide nanoparticles in cosmetics. *journal of food and drug analysis* 23(3), 587-594.
- ❖ Lu, Q., Wu, C., Liu, D., Wang, H., Su, W., Li, H., & Yao, S. (2017). A facile and simple method for synthesis of graphene oxide quantum dots from black carbon. *Green Chemistry*, 19(4), 900-904.
- ❖ Lu, Q., Wu, C., Liu, D., Wang, H., Su, W., Li, H., & Yao, S. (2017). A facile and simple method for synthesis of graphene oxide quantum dots from black carbon. *Green Chemistry*, 19(4), 900-904.
- ❖ Lu, Q., Wu, C., Liu, D., Wang, H., Su, W., Li, H., & Yao, S. (2017). A facile and simple method for synthesis of graphene oxide quantum dots from black carbon. *Green Chemistry*, 19(4), 900-904.
- ❖ Lu, S. C. (2009). Regulation of glutathione synthesis. *Molecular aspects of medicine*, 30(1-2), 42-59.
- ❖ Lu, S., Guo, S., Xu, P., Li, X., Zhao, Y., Gu, W., & Xue, M. (2016). Hydrothermal synthesis of nitrogen-doped carbon dots with real-time live-cell imaging and blood–brain barrier penetration capabilities. *International journal of nanomedicine*, 11, 6325.
- ❖ Lu, W., Qin, X., Asiri, A. M., Al-Youbi, A. O., & Sun, X. (2013). Green synthesis of carbon nanodots as an effective fluorescent probe for sensitive and selective detection of mercury (II) ions. *Journal of nanoparticle research*, 15(1), 1344.
- ❖ Lu, W., Qin, X., Liu, S., Chang, G., Zhang, Y., Luo, Y., & Sun, X. (2012). Economical, green synthesis of fluorescent carbon nanoparticles and their use as probes for sensitive and selective detection of mercury (II) ions. *Analytical chemistry*, 84(12), 5351-5357.

- ❖ Lynn, K. R., & Clevette-Radford, N. A. (1987). Biochemical properties of latices from the Euphorbiaceae. *Phytochemistry*, 26(4), 939-944.
- ❖ Ma, X., Dong, Y., Sun, H., & Chen, N. (2017). Highly fluorescent carbon dots from peanut shells as potential probes for copper ion: The optimization and analysis of the synthetic process. *Materials Today Chemistry*, 5, 1-10.
- ❖ Ma, Y., Zhang, Z., Ren, C., Liu, G., & Chen, X. (2012). A novel colorimetric determination of reduced glutathione in A549 cells based on Fe₃O₄ magnetic nanoparticles as peroxidase mimetics. *Analyst*, 137(2), 485-489.
- ❖ Mao, Y., Bao, Y., Han, D., Li, F., & Niu, L. (2012). Efficient one-pot synthesis of molecularly imprinted silica nanospheres embedded carbon dots for fluorescent dopamine optosensing. *Biosensors and Bioelectronics*, 38(1), 55-60.
- ❖ Mauter, M. S., & Elimelech, M. (2008). Environmental applications of carbon-based nanomaterials. *Environmental Science & Technology*, 42(16), 5843-5859.
- ❖ Mehta, V. N., Mungara, A. K., & Kailasa, S. K. (2013). Dopamine dithiocarbamate functionalized silver nanoparticles as colorimetric sensors for the detection of cobalt ion. *Analytical Methods*, 5(7), 1818-1822.
- ❖ Mei, Q., Chen, J., Zhao, J., Yang, L., Liu, B., Liu, R., & Zhang, Z. (2016). Atomic oxygen tailored graphene oxide nanosheets emissions for multicolor cellular imaging. *ACS applied materials & interfaces*, 8(11), 7390-7395.
- ❖ Ming, H., Ma, Z., Liu, Y., Pan, K., Yu, H., Wang, F., & Kang, Z. (2012). Large scale electrochemical synthesis of high quality carbon nanodots and their photocatalytic property. *Dalton Transactions*, 41(31), 9526-9531.
- ❖ Mobed, A., Hasanzadeh, M., Shadjou, N., Hassanpour, S., Saadati, A., & Agazadeh, M. (2020). Immobilization of ssDNA on the surface of silver nanoparticles-graphene quantum dots modified by gold nanoparticles towards biosensing of microorganism. *Microchemical Journal*, 152, 104286.
- ❖ Moon, B. J., Jang, D., Yi, Y., Lee, H., Kim, S. J., Oh, Y., & Bae, S. (2017). Multi-functional nitrogen self-doped graphene quantum dots for boosting the photovoltaic performance of BHJ solar cells. *Nano Energy*, 34, 36-46.
- ❖ Naganuma, T. (2017) Shape design of cerium oxide nanoparticles for enhancement of enzyme mimetic activity in therapeutic applications. *Nano Research* 10(1), 199-217.
- ❖ Ngamchuea, K., Batchelor-McAuley, C., & Compton, R. G. (2016). The Copper (II)-Catalyzed Oxidation of Glutathione. *Chemistry—A European Journal*, 22(44), 15937-15944.

- ❖ Ni, P., Sun, Y., Dai, H., Hu, J., Jiang, S., Wang, Y., & Li, Z. (2015). Highly sensitive and selective colorimetric detection of glutathione based on Ag [I] ion–3, 3', 5, 5'-tetramethylbenzidine (TMB). *Biosensors and Bioelectronics*, 63, 47-52.
- ❖ Niu, J., & Gao, H. (2014). Synthesis and drug detection performance of nitrogen-doped carbon dots. *Journal of Luminescence*, 149, 159-162.
- ❖ Oh, E., Hong, M. Y., Lee, D., Nam, S. H., Yoon, H. C., & Kim, H. S. (2005). Inhibition assay of biomolecules based on fluorescence resonance energy transfer (FRET) between quantum dots and gold nanoparticles. *Journal of the American Chemical Society*, 127(10), 3270-3271.
- ❖ Omer, K. M., Tofiq, D. I., & Ghafoor, D. D. (2019). Highly photoluminescent label free probe for Chromium (II) ions using carbon quantum dots co-doped with nitrogen and phosphorous. *Journal of Luminescence*, 206, 540-546.
- ❖ Ou, G., Li, Z., Li, D., Cheng, L., Liu, Z., & Wu, H. (2016). Photothermal therapy by using titanium oxide nanoparticles. *Nano Research*, 9(5), 1236-1243.
- ❖ Ouyang, L., Zhu, L., Jiang, J., & Tang, H. (2014). A surface-enhanced Raman scattering method for detection of trace glutathione on the basis of immobilized silver nanoparticles and crystal violet probe. *Analytica chimica acta*, 816, 41-49.
- ❖ Pakrashi, S., Jain, N., Dalai, S., Jayakumar, J., Chandrasekaran, P.T., Raichur, A.M., Chandrasekaran, N. and Mukherjee, A. (2014) In vivo genotoxicity assessment of titanium dioxide nanoparticles by Allium cepa root tip assay at high exposure concentrations. *PLoS One* 9(2), e87789.
- ❖ Pandey, S., Mewada, A., Thakur, M., Pillai, S., Dharmatti, R., Phadke, C., & Sharon, M. (2014). Synthesis of mesoporous silica oxide/C-dot complex (meso-SiO₂/C-dots) using pyrolysed rice husk and its application in bioimaging. *RSC Advances*, 4(3), 1174-1179.
- ❖ Pandey, S., Thakur, M., Mewada, A., Anjarlekar, D., Mishra, N., & Sharon, M. (2013). Carbon dots functionalized gold nanorod mediated delivery of doxorubicin: tri-functional nano-worms for drug delivery, photothermal therapy and bioimaging. *Journal of Materials Chemistry B*, 1(38), 4972-4982.
- ❖ Pardo, J., Peng, Z., & Leblanc, R. M. (2018). Cancer targeting and drug delivery using carbon-based quantum dots and nanotubes. *Molecules*, 23(2), 378.
- ❖ Park, K. S., Kim, M. I., Woo, M. A., & Park, H. G. (2013). A label-free method for detecting biological thiols based on blocking of Hg²⁺-quenching of fluorescent gold nanoclusters. *Biosensors and Bioelectronics*, 45, 65-69.
- ❖ Park, S. Y., Lee, H. U., Park, E. S., Lee, S. C., Lee, J. W., Jeong, S. W., & Lee, J. (2014). Photoluminescent green carbon nanodots from food-waste-derived sources:

- large-scale synthesis, properties, and biomedical applications. *ACS applied materials & interfaces*, 6(5), 3365-3370.
- ❖ Parsons, M., Vojnovic, B., & Ameer-Beg, S. (2004). Imaging protein–protein interactions in cell motility using fluorescence resonance energy transfer (FRET).
 - ❖ Pavase, T. R., Lin, H., Hussain, S., Li, Z., Ahmed, I., Lv, L., & Kalhor, M. T. (2018). Recent advances of conjugated polymer (CP) nanocomposite-based chemical sensors and their applications in food spoilage detection: A comprehensive review. *Sensors and Actuators B: Chemical*, 273, 1113-1138.
 - ❖ Prasannan, A., & Imae, T. (2013). One-pot synthesis of fluorescent carbon dots from orange waste peels. *Industrial & Engineering Chemistry Research*, 52(44), 15673-15678.
 - ❖ Qiao, R., Yang, C. and Gao, M. (2009) Superparamagnetic iron oxide nanoparticles: from preparations to in vivo MRI applications. *Journal of Materials Chemistry* 19(35), 6274-6293.
 - ❖ Qu, D., Zheng, M., Du, P., Zhou, Y., Zhang, L., Li, D., & Sun, Z. (2013). Highly luminescent S, N co-doped graphene quantum dots with broad visible absorption bands for visible light photocatalysts. *Nanoscale*, 5(24), 12272-12277.
 - ❖ Rahban, M., Divsalar, A., Saboury, A.A. and Golestani, A. (2010) Nanotoxicity and spectroscopy studies of silver nanoparticle: Calf thymus DNA and K562 as targets. *The Journal of Physical Chemistry C* 114(13), 5798-5803.
 - ❖ Ramachandran, R., Krishnaraj, C., Sivakumar, A.S., Prasannakumar, P., Kumar, V.A., Shim, K.S., Song, C.-G. and Yun, S.-I. (2017) Anticancer activity of biologically synthesized silver and gold nanoparticles on mouse myoblast cancer cells and their toxicity against embryonic zebrafish. *Materials Science and Engineering: C* 73, 674-683.
 - ❖ Ramalingam, V., Sundaramahalingam, S., & Rajaram, R. (2019). Size-dependent antimycobacterial activity of titanium oxide nanoparticles against Mycobacterium tuberculosis. *Journal of Materials Chemistry B*, 7(27), 4338-4346.
 - ❖ Rao, C. N. R., Venkataraghavan, R., & Kasturi, T. R. (1964). Contribution to the infrared spectra of organosulphur compounds. *Canadian journal of chemistry*, 42(1), 36-42.
 - ❖ Ribeiro, M., Ferraz, M.P., Monteiro, F.J., Fernandes, M.H., Beppu, M.M., Mantione, D. and Sardon, H. (2017) Antibacterial silk fibroin/nanohydroxyapatite hydrogels with silver and gold nanoparticles for bone regeneration. *Nanomedicine: Nanotechnology, Biology and Medicine* 13(1), 231-239.
 - ❖ Rosi, N. L., & Mirkin, C. A. (2005). Nanostructures in biodiagnostics. *Chemical reviews*, 105(4), 1547-1562.

- ❖ Sachdev, A., & Gopinath, P. (2015). Green synthesis of multifunctional carbon dots from coriander leaves and their potential application as antioxidants, sensors and bioimaging agents. *Analyst*, 140(12), 4260-4269.
- ❖ Sahu, S., Behera, B., Maiti, T. K., & Mohapatra, S. (2012). Simple one-step synthesis of highly luminescent carbon dots from orange juice: application as excellent bio-imaging agents. *Chemical communications*, 48(70), 8835-8837.
- ❖ Sakamoto-Arnold, C. M., & Johnson, K. S. (1987). Determination of picomolar levels of cobalt in seawater by flow injection analysis with chemiluminescence detection. *Analytical Chemistry*, 59(14), 1789-1794.
- ❖ Santos, L., Silveira, C.M., Elangovan, E., Neto, J.P., Nunes, D., Pereira, L., Martins, R., Viegas, J., Moura, J.J. and Todorovic, S. (2016) Synthesis of WO₃ nanoparticles for biosensing applications. *Sensors and Actuators B: Chemical* 223, 186-194.
- ❖ Santra, S., Yang, H., Stanley, J. T., Holloway, P. H., Moudgil, B. M., Walter, G., & Mericle, R. A. (2005). Rapid and effective labeling of brain tissue using TAT-conjugated CdS: Mn/ZnS quantum dots. *Chemical communications*, (25), 3144-3146.
- ❖ Saxena, A., Tripathi, R., Zafar, F. and Singh, P. (2012) Green synthesis of silver nanoparticles using aqueous solution of *Ficus benghalensis* leaf extract and characterization of their antibacterial activity. *Materials Letters* 67(1), 91-94.
- ❖ Schmidt, T., Gasteiger, H. and Behm, R. (1999) Rotating disk electrode measurements on the CO tolerance of a high-surface area Pt/vulcan carbon fuel cell catalyst. *Journal of The Electrochemical Society* 146(4), 1296-1304.
- ❖ Scida, K., Stege, P. W., Haby, G., Messina, G. A., & García, C. D. (2011). Recent applications of carbon-based nanomaterials in analytical chemistry: critical review. *Analytica chimica acta*, 691(1-2), 6-17.
- ❖ Selvin, P. R. (2000). The renaissance of fluorescence resonance energy transfer. *Nature structural biology*, 7(9), 730-734.
- ❖ Shamsipur, M., Safavi, A., & Mohammadpour, Z. (2014). Indirect colorimetric detection of glutathione based on its radical restoration ability using carbon nanodots as nanozymes. *Sensors and Actuators B: Chemical*, 199, 463-469.
- ❖ Shan, X., Chai, L., Ma, J., Qian, Z., Chen, J., & Feng, H. (2014). B-doped carbon quantum dots as a sensitive fluorescence probe for hydrogen peroxide and glucose detection. *Analyst*, 139(10), 2322-2325.

- ❖ Shao, N., Zhang, Y., Cheung, S., Yang, R., Chan, W., Mo, T., & Liu, F. (2005). Copper ion-selective fluorescent sensor based on the inner filter effect using a spiropyran derivative. *Analytical Chemistry*, 77(22), 7294-7303.
- ❖ Sharma, K.D. (2017) Antibacterial Activity of Biogenic Platinum Nanoparticles: An invitro Study. *Int. J. Curr. Microbiol. App. Sci* 6(2), 801-808.
- ❖ Shen, C., Wang, J., Cao, Y., & Lu, Y. (2015). Facile access to B-doped solid-state fluorescent carbon dots toward light emitting devices and cell imaging agents. *Journal of Materials Chemistry C*, 3(26), 6668-6675.
- ❖ Shen, J., Zhu, Y., Chen, C., Yang, X., & Li, C. (2011). Facile preparation and upconversion luminescence of graphene quantum dots. *Chemical communications*, 47(9), 2580-2582.
- ❖ Shen, J., Zhu, Y., Yang, X., & Li, C. (2012). Graphene quantum dots: emergent nanolights for bioimaging, sensors, catalysis and photovoltaic devices. *Chemical communications*, 48(31), 3686-3699.
- ❖ Shen, W., Qu, Y., Pei, X., Li, S., You, S., Wang, J., Zhang, Z. and Zhou, J. (2017) Catalytic reduction of 4-nitrophenol using gold nanoparticles biosynthesized by cell-free extracts of *Aspergillus sp.* WL-Au. *Journal of Hazardous Materials* 321, 299-306.
- ❖ Shi, C., Qi, H., Ma, R., Sun, Z., Xiao, L., Wei, G., & Fan, J. (2019). N, S-self-doped carbon quantum dots from fungus fibers for sensing tetracyclines and for bioimaging cancer cells. *Materials Science and Engineering: C*, 105, 110132.
- ❖ Shi, J., Lu, C., Yan, D., & Ma, L. (2013). High selectivity sensing of cobalt in HepG2 cells based on necklace model microenvironment-modulated carbon dot-improved chemiluminescence in Fenton-like system. *Biosensors and Bioelectronics*, 45, 58-64.
- ❖ Shi, W., Guo, F., Han, M., Yuan, S., Guan, W., Li, H., & Kang, Z. (2017). N, S co-doped carbon dots as a stable bio-imaging probe for detection of intracellular temperature and tetracycline. *Journal of Materials Chemistry B*, 5(18), 3293-3299.
- ❖ Shi, W., Li, X., & Ma, H. (2012). A tunable ratiometric pH sensor based on carbon nanodots for the quantitative measurement of the intracellular pH of whole cells. *Angewandte Chemie International Edition*, 51(26), 6432-6435.
- ❖ Shi, Y., Pan, Y., Zhang, H., Zhang, Z., Li, M. J., Yi, C., & Yang, M. (2014). A dual-mode nanosensor based on carbon quantum dots and gold nanoparticles for discriminative detection of glutathione in human plasma. *Biosensors and Bioelectronics*, 56, 39-45.

- ❖ Sinduja, B., & John, S. A. (2016). Sensitive determination of tannic acid using blue luminescent graphene quantum dots as fluorophore. *RSC advances*, 6(65), 59900-59906.
- ❖ Singh, V. K., Singh, V., Yadav, P. K., Chandra, S., Bano, D., Koch, B., & Hasan, S. H. (2019). Nitrogen doped fluorescent carbon quantum dots for on-off-on detection of Hg²⁺ and glutathione in aqueous medium: Live cell imaging and IMPLICATION logic gate operation. *Journal of Photochemistry and Photobiology A: Chemistry*, 384, 112042.
- ❖ Singh, V. K., Singh, V., Yadav, P. K., Chandra, S., Bano, D., Kumar, V., & Hasan, S. H. (2018). Bright-blue-emission nitrogen and phosphorus-doped carbon quantum dots as a promising nanoprobe for detection of Cr (VI) and ascorbic acid in pure aqueous solution and in living cells. *New Journal of Chemistry*, 42(15), 12990-12997.
- ❖ Srivastava, S., Awasthi, R., Tripathi, D., Rai, M. K., Agarwal, V., Agrawal, V., & Gupta, R. K. (2012). Magnetic-nanoparticle-doped carbogenic nanocomposite: an effective magnetic resonance/fluorescence multimodal imaging probe. *Small*, 8(7), 1099-1109.
- ❖ Stroh, M., Zimmer, J. P., Duda, D. G., Levchenko, T. S., Cohen, K. S., Brown, E. B., & Jain, R. K. (2005). Quantum dots spectrally distinguish multiple species within the tumor milieu in vivo. *Nature medicine*, 11(6), 678.
- ❖ Sun, H., Wu, L., Wei, W., & Qu, X. (2013). Recent advances in graphene quantum dots for sensing. *Materials today*, 16(11), 433-442.
- ❖ Sun, Y. P., Zhou, B., Lin, Y., Wang, W., Fernando, K. S., Pathak, P., & Luo, P. G. (2006). Quantum-sized carbon dots for bright and colorful photoluminescence. *Journal of the American Chemical Society*, 128(24), 7756-7757.
- ❖ Sun, Y. P., Zhou, B., Lin, Y., Wang, W., Fernando, K. S., Pathak, P., & Luo, P. G. (2006). Quantum-sized carbon dots for bright and colorful photoluminescence. *Journal of the American Chemical Society*, 128(24), 7756-7757.
- ❖ Sung, H. K., Oh, S. Y., Park, C., & Kim, Y. (2013). Colorimetric detection of Co²⁺ ion using silver nanoparticles with spherical, plate, and rod shapes. *Langmuir*, 29(28), 8978-8982.
- ❖ Taghavi, S., Abnous, K., Taghdisi, S. M., Ramezani, M., & Alibolandi, M. (2020). Hybrid carbon-based materials for gene delivery in cancer therapy. *Journal of Controlled Release*, 318, 158-175.
- ❖ Tahir, K., Nazir, S., Ahmad, A., Li, B., Khan, A.U., Khan, Z.U.H., Khan, F.U., Khan, Q.U., Khan, A. and Rahman, A.U. (2017) Facile and green synthesis of phytochemicals capped platinum nanoparticles and in vitro their superior antibacterial activity. *Journal of Photochemistry and Photobiology B: Biology* 166, 246-251.

-
- ❖ Tan, X., Li, Y., Li, X., Zhou, S., Fan, L., & Yang, S. (2015). Electrochemical synthesis of small-sized red fluorescent graphene quantum dots as a bioimaging platform. *Chemical Communications*, 51(13), 2544-2546.
 - ❖ Tang, L., Ji, R., Li, X., Teng, K. S., & Lau, S. P. (2013). Energy-level structure of nitrogen-doped graphene quantum dots. *Journal of Materials Chemistry C*, 1(32), 4908-4915.
 - ❖ Tang, X., Zeng, X., Liu, H., Yang, Y., Zhou, H., & Cai, H. (2019). A nanohybrid composed of MoS₂ quantum dots and MnO₂ nanosheets with dual-emission and peroxidase mimicking properties for use in ratiometric fluorometric detection and cellular imaging of glutathione. *Microchimica Acta*, 186(8), 572.
 - ❖ Taniguchi, N. (1974) On the basic concept of nano-technology Proceedings of the International Conference on Production Engineering Tokyo Part II *Japan Society of Precision Engineering*.
 - ❖ Thakur, M., Pandey, S., Mewada, A., Patil, V., Khade, M., Goshi, E., & Sharon, M. (2014). Antibiotic conjugated fluorescent carbon dots as a theranostic agent for controlled drug release, bioimaging, and enhanced antimicrobial activity. *Journal of drug delivery*, 2014.
 - ❖ Thirumurugan, A., Aswitha, P., Kiruthika, C., Nagarajan, S., & Christy, A. N. (2016). Green synthesis of platinum nanoparticles using *Azadirachta indica*—An eco-friendly approach. *Materials Letters*, 170, 175-178.
 - ❖ Tian, D., Qian, Z., Xia, Y., & Zhu, C. (2012). Gold nanocluster-based fluorescent probes for near-infrared and turn-on sensing of glutathione in living cells. *Langmuir*, 28(8), 3945-3951.
 - ❖ Tian, L., Ghosh, D., Chen, W., Pradhan, S., Chang, X., & Chen, S. (2009). Nanosized carbon particles from natural gas soot. *Chemistry of materials*, 21(13), 2803-2809.
 - ❖ Townsend, D. M., Tew, K. D., & Tapiero, H. (2003). The importance of glutathione in human disease. *Biomedicine & Pharmacotherapy*, 57(3-4), 145-155.
 - ❖ Travlou, N. A., Secor, J., & Bandosz, T. J. (2017). Highly luminescent S-doped carbon dots for the selective detection of ammonia. *Carbon*, 114, 544-556.
 - ❖ Tsikas, D., Raida, M., Sandmann, J., Rossa, S., Forssmann, W. G., & Frölich, J. C. (2000). Electrospray ionization mass spectrometry of low-molecular-mass S-nitroso compounds and their thiols. *Journal of Chromatography B: Biomedical Sciences and Applications*, 742(1), 99-108.
 - ❖ Varun, S., Daniel, S.K. and Gorthi, S.S. (2017) Rapid sensing of melamine in milk by interference green synthesis of silver nanoparticles. *Materials Science and Engineering: C* 74, 253-258.

- ❖ Vivek, R., Thangam, R., Muthuchelian, K., Gunasekaran, P., Kaveri, K. and Kannan, S. (2012) Green biosynthesis of silver nanoparticles from *Annona squamosa* leaf extract and its in vitro cytotoxic effect on MCF-7 cells. *Process Biochemistry* 47(12), 2405-2410.
- ❖ Wang, C. I., Wu, W. C., Periasamy, A. P., & Chang, H. T. (2014). Electrochemical synthesis of photoluminescent carbon nanodots from glycine for highly sensitive detection of hemoglobin. *Green Chemistry*, 16(5), 2509-2514.
- ❖ Wang, C., Wang, Y., Shi, H., Yan, Y., Liu, E., Hu, X., & Fan, J. (2019). A strong blue fluorescent nanoprobe for highly sensitive and selective detection of mercury (II) based on sulfur doped carbon quantum dots. *Materials Chemistry and Physics*, 232, 145-151.
- ❖ Wang, H., Liang, S. C., Zhang, Z. M., & Zhang, H. S. (2004). 3-Iodoacetylaminobenzanthrone as a fluorescent derivatizing reagent for thiols in high-performance liquid chromatography. *Analytica chimica acta*, 512(2), 281-286.
- ❖ Wang, H., Lu, Q., Hou, Y., Liu, Y., & Zhang, Y. (2016). High fluorescence S, N co-doped carbon dots as an ultra-sensitive fluorescent probe for the determination of uric acid. *Talanta*, 155, 62-69.
- ❖ Wang, H., Sun, P., Cong, S., Wu, J., Gao, L., Wang, Y., & Zou, G. (2016). Nitrogen-doped carbon dots for “green” quantum dot solar cells. *Nanoscale research letters*, 11(1), 1-6.
- ❖ Wang, J., Zhang, P., Huang, C., Liu, G., Leung, K. C. F., & Wang, Y. X. J. (2015). High performance photoluminescent carbon dots for in vitro and in vivo bioimaging: effect of nitrogen doping ratios. *Langmuir*, 31(29), 8063-8073.
- ❖ Wang, L., Ruan, F., Lv, T., Liu, Y., Deng, D., Zhao, S., & Xu, S. (2015). One step synthesis of Al/N co-doped carbon nanoparticles with enhanced photoluminescence. *Journal of Luminescence*, 158, 1-5.
- ❖ Wang, L., Wu, T., Du, S., Pei, M., Guo, W., & Wei, S. (2016). High performance supercapacitors based on ternary graphene/Au/polyaniline (PANI) hierarchical nanocomposites. *RSC advances*, 6(2), 1004-1011.
- ❖ Wang, Q., Yang, Z., Zhang, X., Xiao, X., Chang, C. K., & Xu, B. (2007). A supramolecular-hydrogel-encapsulated hemin as an artificial enzyme to mimic peroxidase. *Angewandte Chemie International Edition*, 46(23), 4285-4289.
- ❖ Wang, Q., Zheng, H., Long, Y., Zhang, L., Gao, M., & Bai, W. (2011). Microwave-hydrothermal synthesis of fluorescent carbon dots from graphite oxide. *Carbon*, 49(9), 3134-3140.

- ❖ Wang, W., Li, Y., Cheng, L., Cao, Z., & Liu, W. (2013). Water-soluble and phosphorus-containing carbon dots with strong green fluorescence for cell labeling. *Journal of Materials Chemistry B*, 2(1), 46-48.
- ❖ Wang, Y., & Hu, A. (2014). Carbon quantum dots: synthesis, properties and applications. *Journal of Materials Chemistry C*, 2(34), 6921-6939.
- ❖ Wardle, B. (2009). *Principles and applications of photochemistry*. John Wiley & Sons.
- ❖ Wei, H., Bruns, O. T., Kaul, M. G., Hansen, E. C., Barch, M., Wiśniowska, A., & Cordero, J. M. (2017). Exceedingly small iron oxide nanoparticles as positive MRI contrast agents. *Proceedings of the national academy of sciences*, 114(9), 2325-2330.
- ❖ Wei, J., Wang, H., Zhang, Q., & Li, Y. (2015). One-pot hydrothermal synthesis of N-doped carbon quantum dots using the waste of shrimp for hydrogen evolution from formic acid. *Chemistry Letters*, 44(3), 241-243.
- ❖ Wei, W., Xu, C., Ren, J., Xu, B., & Qu, X. (2012). Sensing metal ions with ion selectivity of a crown ether and fluorescence resonance energy transfer between carbon dots and graphene. *Chemical Communications*, 48(9), 1284-1286.
- ❖ Wen, X., Shi, L., Wen, G., Li, Y., Dong, C., Yang, J., & Shuang, S. (2016). Green and facile synthesis of nitrogen-doped carbon nanodots for multicolor cellular imaging and Co²⁺ sensing in living cells. *Sensors and Actuators B: Chemical*, 235, 179-187.
- ❖ Wen, X., Shi, L., Wen, G., Li, Y., Dong, C., Yang, J., & Shuang, S. (2015). Green synthesis of carbon nanodots from cotton for multicolor imaging, patterning, and sensing. *Sensors and Actuators B: Chemical*, 221, 769-776.
- ❖ Wu, P., Li, W., Wu, Q., Liu, Y., & Liu, S. (2017). Hydrothermal synthesis of nitrogen-doped carbon quantum dots from microcrystalline cellulose for the detection of Fe³⁺ ions in an acidic environment. *RSC advances*, 7(70), 44144-44153.
- ❖ Wu, X., Song, Y., Yan, X., Zhu, C., Ma, Y., Du, D., & Lin, Y. (2017). Carbon quantum dots as fluorescence resonance energy transfer sensors for organophosphate pesticides determination. *Biosensors and Bioelectronics*, 94, 292-297.
- ❖ Wu, Z. L., Zhang, P., Gao, M. X., Liu, C. F., Wang, W., Leng, F., & Huang, C. Z. (2013). One-pot hydrothermal synthesis of highly luminescent nitrogen-doped amphoteric carbon dots for bioimaging from Bombyx mori silk–natural proteins. *Journal of Materials Chemistry B*, 1(22), 2868-2873.
- ❖ Xu, H., & Hepel, M. (2011). “Molecular beacon”-based fluorescent assay for selective detection of glutathione and cysteine. *Analytical Chemistry*, 83(3), 813-819.

- ❖ Xu, J., Li, D., Chen, Y., Tan, L., Kou, B., Wan, F., & Li, F. (2017). Constructing sheet-on-sheet structured graphitic carbon nitride/reduced graphene oxide/layered MnO₂ ternary nanocomposite with outstanding catalytic properties on thermal decomposition of ammonium perchlorate. *Nanomaterials*, 7(12), 450.
- ❖ Xu, L., Fan, H., Huang, L., Xia, J., Huang, J., Li, M., & Li, S. (2017). Eosinophilic nitrogen-doped carbon dots derived from tribute chrysanthemum for label-free detection of Fe³⁺ ions and hydrazine. *Journal of the Taiwan Institute of Chemical Engineers*, 78, 247-253.
- ❖ Xu, X., Ray, R., Gu, Y., Ploehn, H. J., Gearheart, L., Raker, K., & Scrivens, W. A. (2004). Electrophoretic analysis and purification of fluorescent single-walled carbon nanotube fragments. *Journal of the American Chemical Society*, 126(40), 12736-12737.
- ❖ Xue, M., Zhang, L., Zhan, Z., Zou, M., Huang, Y., & Zhao, S. (2016). Sulfur and nitrogen binary doped carbon dots derived from ammonium thiocyanate for selective probing doxycycline in living cells and multicolor cell imaging. *Talanta*, 150, 324-330.
- ❖ Xue, M., Zou, M., Zhao, J., Zhan, Z., & Zhao, S. (2015). Green preparation of fluorescent carbon dots from lychee seeds and their application for the selective detection of methylene blue and imaging in living cells. *Journal of Materials Chemistry B*, 3(33), 6783-6789.
- ❖ Yadav, P. K., Singh, V. K., Kumar, C., Chandra, S., Jit, S., Singh, S. K., & Hasan, S. H. (2019). A Facile Synthesis of Green-Blue Carbon Dots from Artocarpus lakoocha Seeds and Their Application for the Detection of Iron (III) in Biological Fluids and Cellular Imaging. *ChemistrySelect*, 4(42), 12252-12259.
- ❖ Yan, F., Zou, Y., Wang, M., Mu, X., Yang, N., & Chen, L. (2014). Highly photoluminescent carbon dots-based fluorescent chemosensors for sensitive and selective detection of mercury ions and application of imaging in living cells. *Sensors and Actuators B: Chemical*, 192, 488-495.
- ❖ Yan, X., Song, Y., Zhu, C., Song, J., Du, D., Su, X., & Lin, Y. (2016). Graphene quantum dot-MnO₂ nanosheet based optical sensing platform: a sensitive fluorescence “turn off-on” nanosensor for glutathione detection and intracellular imaging. *ACS applied materials & interfaces*, 8(34), 21990-21996.
- ❖ Yang, F., He, X., Wang, C., Cao, Y., Li, Y., Yan, L., & Li, Y. (2018). Controllable and eco-friendly synthesis of P-rich carbon quantum dots and its application for copper (II) ion sensing. *Applied Surface Science*, 448, 589-598.
- ❖ Yang, H., Li, S., Feng, F., Ou, S., Li, F., Yang, M., & Ma, J. (2019). Palladium nanoparticles with surface enrichment of palladium oxide species immobilized on the

- aniline-functionalized graphene as an advanced electrocatalyst of ethanol oxidation. *ACS Sustainable Chemistry & Engineering*, 7(17), 14621-14628.
- ❖ Yang, P., Zhao, J., Wang, J., Cao, B., Li, L., & Zhu, Z. (2014). Light-induced synthesis of photoluminescent carbon nanoparticles for Fe³⁺ sensing and photocatalytic hydrogen evolution. *Journal of Materials Chemistry A*, 3(1), 136-138.
 - ❖ Yang, P., Zhao, J., Zhang, L., Li, L., & Zhu, Z. (2015). Intramolecular hydrogen bonds quench photoluminescence and enhance photocatalytic activity of carbon nanodots. *Chemistry—A European Journal*, 21(23), 8561-8568.
 - ❖ Yang, S. T., Cao, L., Luo, P. G., Lu, F., Wang, X., Wang, H., & Sun, Y. P. (2009). Carbon dots for optical imaging in vivo. *Journal of the American Chemical Society*, 131(32), 11308-11309.
 - ❖ Yang, S. T., Wang, X., Wang, H., Lu, F., Luo, P. G., Cao, L., & Huang, Y. (2009). Carbon dots as nontoxic and high-performance fluorescence imaging agents. *The Journal of Physical Chemistry C*, 113(42), 18110-18114.
 - ❖ Yang, S.J. and Park, C.R. (2007) Facile preparation of monodisperse ZnO quantum dots with high quality photoluminescence characteristics. *Nanotechnology* 19(3), 035609.
 - ❖ Yang, X., Wang, K., & Guo, C. (2000). A fluorescent optode for sodium ion based on the inner filter effect. *Analytica chimica acta*, 407(1-2), 45-52.
 - ❖ Yang, Y., Huo, D., Wu, H., Wang, X., Yang, J., Bian, M., & Hou, C. (2018). N, P-doped carbon quantum dots as a fluorescent sensing platform for carbendazim detection based on fluorescence resonance energy transfer. *Sensors and Actuators B: Chemical*, 274, 296-303.
 - ❖ Yao, Y., Tian, D., & Li, H. (2010). Cooperative binding of bifunctionalized and click-synthesized silver nanoparticles for colorimetric Co²⁺ sensing. *ACS applied materials & interfaces*, 2(3), 684-690.
 - ❖ Yu, J., Xu, C., Tian, Z., Lin, Y., & Shi, Z. (2016). Facilely synthesized N-doped carbon quantum dots with high fluorescent yield for sensing Fe³⁺. *New Journal of Chemistry*, 40(3), 2083-2088.
 - ❖ Yuan, C., Liu, B., Liu, F., Han, M. Y., & Zhang, Z. (2014). Fluorescence “turn on” detection of mercuric ion based on bis (dithiocarbamate) copper (II) complex functionalized carbon nanodots. *Analytical chemistry*, 86(2), 1123-1130.
 - ❖ Yuan, P., & Walt, D. R. (1987). Calculation for fluorescence modulation by absorbing species and its application to measurements using optical fibers. *Analytical Chemistry*, 59(19), 2391-2394.

- ❖ Zhai, X., Zhang, P., Liu, C., Bai, T., Li, W., Dai, L., & Liu, W. (2012). Highly luminescent carbon nanodots by microwave-assisted pyrolysis. *Chemical Communications*, 48(64), 7955-7957.
- ❖ Zhang, H., Chen, Y., Liang, M., Xu, L., Qi, S., Chen, H., & Chen, X. (2014). Solid-phase synthesis of highly fluorescent nitrogen-doped carbon dots for sensitive and selective probing ferric ions in living cells. *Analytical chemistry*, 86(19), 9846-9852.
- ❖ Zhang, J., Tian, Z., Liang, L., Subirade, M., & Chen, L. (2013). Binding interactions of β -conglycinin and glycinin with vitamin B12. *The Journal of Physical Chemistry B*, 117(45), 14018-14028.
- ❖ Zhang, M., Liu, Y. Q., & Ye, B. C. (2012). Colorimetric assay for parallel detection of Cd²⁺, Ni²⁺ and Co²⁺ using peptide-modified gold nanoparticles. *Analyst*, 137(3), 601-607.
- ❖ Zhang, N., Qu, F., Luo, H. Q., & Li, N. B. (2013). Sensitive and selective detection of biothiols based on target-induced agglomeration of silver nanoclusters. *Biosensors and Bioelectronics*, 42, 214-218.
- ❖ Zhang, R., & Chen, W. (2014). Nitrogen-doped carbon quantum dots: Facile synthesis and application as a “turn-off” fluorescent probe for detection of Hg²⁺ ions. *Biosensors and Bioelectronics*, 55, 83-90.
- ❖ Zhang, S., Sun, W. L., Xian, Y. Z., Zhang, W., Jin, L. T., Yamamoto, K., & Jin, J. (1999). Multichannel amperometric detection system for liquid chromatography to assay the thiols in human whole blood using the platinum microelectrodes chemically modified by copper tetraaminophthalocyanine. *Analytica chimica acta*, 399(3), 213-221.
- ❖ Zhang, W., Nefedov, A., Naboka, M., Cao, L., & Wöll, C. (2012). Molecular orientation of terephthalic acid assembly on epitaxial graphene: NEXAFS and XPS study. *Physical chemistry chemical physics*, 14(29), 10125-10131.
- ❖ Zhang, X. L., Zheng, C., Guo, S. S., Li, J., Yang, H. H., & Chen, G. (2014). Turn-on fluorescence sensor for intracellular imaging of glutathione using g-C₃N₄ nanosheet–MnO₂ sandwich nanocomposite. *Analytical chemistry*, 86(7), 3426-3434.
- ❖ Zhang, Y. Q., Ma, D. K., Zhang, Y. G., Chen, W., & Huang, S. M. (2013). N-doped carbon quantum dots for TiO₂-based photocatalysts and dye-sensitized solar cells. *Nano Energy*, 2(5), 545-552.
- ❖ Zhao, H. X., Liu, L. Q., De Liu, Z., Wang, Y., Zhao, X. J., & Huang, C. Z. (2011). Highly selective detection of phosphate in very complicated matrixes with an off–on fluorescent probe of europium-adjusted carbon dots. *Chemical Communications*, 47(9), 2604-2606.

- ❖ Zhao, L., Li, H., Liu, H., Liu, M., Huang, N., He, Z., & Ding, L. (2019). Microwave-assisted facile synthesis of polymer dots as a fluorescent probe for detection of cobalt (II) and manganese (II). *Analytical and bioanalytical chemistry*, 411(11), 2373-2381.
- ❖ Zhao, Q. L., Zhang, Z. L., Huang, B. H., Peng, J., Zhang, M., & Pang, D. W. (2008). Facile preparation of low cytotoxicity fluorescent carbon nanocrystals by electrooxidation of graphite. *Chemical Communications*, (41), 5116-5118.
- ❖ Zhao, S., Lan, M., Zhu, X., Xue, H., Ng, T. W., Meng, X., & Zhang, W. (2015). Green synthesis of bifunctional fluorescent carbon dots from garlic for cellular imaging and free radical scavenging. *ACS applied materials & interfaces*, 7(31), 17054-17060.
- ❖ Zheng, L., Chi, Y., Dong, Y., Lin, J., & Wang, B. (2009). Electrochemiluminescence of water-soluble carbon nanocrystals released electrochemically from graphite. *Journal of the American Chemical Society*, 131(13), 4564-4565.
- ❖ Zheng, L., Chi, Y., Dong, Y., Lin, J., & Wang, B. (2009). Electrochemiluminescence of water-soluble carbon nanocrystals released electrochemically from graphite. *Journal of the American Chemical Society*, 131(13), 4564-4565.
- ❖ Zheng, M., Liu, S., Li, J., Qu, D., Zhao, H., Guan, X., & Sun, Z. (2014). Integrating oxaliplatin with highly luminescent carbon dots: an unprecedented theranostic agent for personalized medicine. *Advanced Materials*, 26(21), 3554-3560.
- ❖ Zheng, M., Xie, Z., Qu, D., Li, D., Du, P., Jing, X., & Sun, Z. (2013). On-off-on fluorescent carbon dot nanosensor for recognition of chromium (VI) and ascorbic acid based on the inner filter effect. *ACS applied materials & interfaces*, 5(24), 13242-13247.
- ❖ Zheng, X. T., Ananthanarayanan, A., Luo, K. Q., & Chen, P. (2015). Glowing graphene quantum dots and carbon dots: properties, syntheses, and biological applications. *Small*, 11(14), 1620-1636.
- ❖ Zhong, W., Liang, J., & Yu, J. (2009). Systematic study of the interaction of cobalt ions with different-sized CdTe quantum dots. *Spectrochimica Acta Part A: Molecular and Biomolecular Spectroscopy*, 74(3), 603-606.
- ❖ Zhou, C., He, X., Ya, D., Zhong, J., & Deng, B. (2017). One step hydrothermal synthesis of nitrogen-doped graphitic quantum dots as a fluorescent sensing strategy for highly sensitive detection of metacycline in mice plasma. *Sensors and Actuators B: Chemical*, 249, 256-264.
- ❖ Zhou, G., Paek, E., Hwang, G. S., & Manthiram, A. (2015). Long-life Li/polysulphide batteries with high sulphur loading enabled by lightweight three-

- dimensional nitrogen/sulphur-codoped graphene sponge. *Nature communications*, 6(1), 1-11.
- ❖ Zhou, J., Booker, C., Li, R., Zhou, X., Sham, T. K., Sun, X., & Ding, Z. (2007). An electrochemical avenue to blue luminescent nanocrystals from multiwalled carbon nanotubes (MWCNTs). *Journal of the American Chemical Society*, 129(4), 744-745.
 - ❖ Zhou, J., Booker, C., Li, R., Zhou, X., Sham, T. K., Sun, X., & Ding, Z. (2007). An electrochemical avenue to blue luminescent nanocrystals from multiwalled carbon nanotubes (MWCNTs). *Journal of the American Chemical Society*, 129(4), 744-745.
 - ❖ Zhou, J., Shan, X., Ma, J., Gu, Y., Qian, Z., Chen, J., & Feng, H. (2014). Facile synthesis of P-doped carbon quantum dots with highly efficient photoluminescence. *Rsc Advances*, 4(11), 5465-5468.
 - ❖ Zhou, J., Zhou, H., Tang, J., Deng, S., Yan, F., Li, W., & Qu, M. (2017). Carbon dots doped with heteroatoms for fluorescent bioimaging: a review. *Microchimica Acta*, 184(2), 343-368.
 - ❖ Zhou, L., Lin, Y., Huang, Z., Ren, J., & Qu, X. (2012). Carbon nanodots as fluorescence probes for rapid, sensitive, and label-free detection of Hg 2+ and biothiols in complex matrices. *Chemical communications*, 48(8), 1147-1149.
 - ❖ Zhou, L., Lin, Y., Huang, Z., Ren, J., & Qu, X. (2012). Carbon nanodots as fluorescence probes for rapid, sensitive, and label-free detection of Hg 2+ and biothiols in complex matrices. *Chemical communications*, 48(8), 1147-1149.
 - ❖ Zhou, W., Zhuang, J., Li, W., Hu, C., Lei, B., & Liu, Y. (2017). Towards efficient dual-emissive carbon dots through sulfur and nitrogen co-doped. *Journal of Materials Chemistry C*, 5(32), 8014-8021.
 - ❖ Zhu, C., Zhai, J., & Dong, S. (2012). Bifunctional fluorescent carbon nanodots: green synthesis via soy milk and application as metal-free electrocatalysts for oxygen reduction. *Chemical communications*, 48(75), 9367-9369.
 - ❖ Zhu, H., Wang, X., Li, Y., Wang, Z., Yang, F., & Yang, X. (2009). Microwave synthesis of fluorescent carbon nanoparticles with electrochemiluminescence properties. *Chemical Communications*, (34), 5118-5120.
 - ❖ Zhu, H., Wang, X., Li, Y., Wang, Z., Yang, F., & Yang, X. (2009). Microwave synthesis of fluorescent carbon nanoparticles with electrochemiluminescence properties. *Chemical Communications*, (34), 5118-5120.
 - ❖ Zhu, N., Ji, H., Yu, P., Niu, J., Farooq, M. U., Akram, M. W., & Niu, X. (2018). Surface modification of magnetic iron oxide nanoparticles. *Nanomaterials*, 8(10), 810.

-
- ❖ Zhu, S., Meng, Q., Wang, L., Zhang, J., Song, Y., Jin, H., & Yang, B. (2013). Highly photoluminescent carbon dots for multicolor patterning, sensors, and bioimaging. *Angewandte Chemie International Edition*, 52(14), 3953-3957.
 - ❖ Zhu, S., Song, Y., Zhao, X., Shao, J., Zhang, J., & Yang, B. (2015). The photoluminescence mechanism in carbon dots (graphene quantum dots, carbon nanodots, and polymer dots): current state and future perspective. *Nano research*, 8(2), 355-381.
 - ❖ Zhu, S., Wang, S., Xia, M., Wang, B., Huang, Y., Zhang, D., & Wang, G. (2019). Intracellular Imaging of Glutathione with MnO₂ Nanosheet@ Ru (bpy) 3²⁺-UiO-66 Nanocomposites. *ACS applied materials & interfaces*, 11(35), 31693-31699.
 - ❖ Zhu, S., Zhang, J., Tang, S., Qiao, C., Wang, L., Wang, H., & Wang, X. (2012). Surface chemistry routes to modulate the photoluminescence of graphene quantum dots: from fluorescence mechanism to up-conversion bioimaging applications. *Advanced Functional Materials*, 22(22), 4732-4740.
 - ❖ Zhu, S., Zhang, J., Tang, S., Qiao, C., Wang, L., Wang, H., & Wang, X. (2012). Surface chemistry routes to modulate the photoluminescence of graphene quantum dots: from fluorescence mechanism to up-conversion bioimaging applications. *Advanced Functional Materials*, 22(22), 4732-4740.
 - ❖ Zhuo, S., Shao, M., & Lee, S. T. (2013). ACS Nano, 2012, 6, 1059; H. Choi, SJ Ko, Y. Choi, P. Joo, T. Kim, BR Lee, JW Jung, HJ Choi, M. Cha, JR Jeong, IW Hwang, MH Song, BS Kim and JY Kim, *Nat. Photonics*.
 - ❖ Zi, L., Huang, Y., Yan, Z., & Liao, S. (2014). Thioglycolic acid-capped CuInS₂/ZnS quantum dots as fluorescent probe for cobalt ion detection. *Journal of luminescence*, 148, 359-363.
 - ❖ Zuo, P., Liu, J., Guo, H., Wang, C., Liu, H., Zhang, Z., & Liu, Q. (2019). Multifunctional N, S co-doped carbon dots for sensitive probing of temperature, ferric ion, and methotrexate. *Analytical and bioanalytical chemistry*, 411(8), 1647-1657.
 - ❖ Zuo, P., Lu, X., Sun, Z., Guo, Y., & He, H. (2016). A review on syntheses, properties, characterization and bioanalytical applications of fluorescent carbon dots. *Microchimica Acta*, 183(2), 519-542.