

Contents

PREFACE	iii
List of Tables	xiii
List of Figures	xv
List of Abbreviations	xxiii
1 General Introduction	1
1.1 CRYSTAL STRUCTURE OF PRUSSIAN BLUE	2
1.2 COLOUR OF PB	3
1.3 PROPERTIES OF PB.....	4
1.3.1 Electrochemical.....	4
1.3.2 Electrochromic	7
1.3.3 Photo-physical.....	8
1.3.4 Magnetic.....	9
1.3.5 Charging and Discharging Characteristics.....	10
1.4 APPLICATION OF PB AND ITS ANALOGUES MODIFIED ELECTRODES IN CHEMICAL SENSING	11
1.4.1 Electrochemical Sensor for Oxidizable Compounds.....	11
1.4.2 Electrochemical Sensor for Non-electroactive Cations.....	13
1.4.3 Advanced Transducer for Hydrogen peroxide (H ₂ O ₂)	14
1.4.4 Electrochemical Biosensors based on PB and PBAs	15
1.4.5 Photo-Fenton Catalysis based on PBN.....	16
1.5 THE ARTIFICIAL PEROXIDASE ACTIVITY OF NANOMATERIALS: APPLICATION IN H ₂ O ₂ AND GLUCOSE SENSING.....	18

1.6	CHALLENGES IN THE SYNTHESIS OF PB AND ITS ANALYTICAL CHEMISTRY	19
1.7	ORIGIN OF THE PRESENT RESEARCH PROBLEMS	21
1.8	OBJECTIVES OF THE PRESENT INVESTIGATION.....	22
1.9	WORK PLAN OF THE THESIS	23
2	Functional Alkoxysilane Mediated Controlled Synthesis of Prussian Blue Nanoparticles, Enabling Silica-Alginate Bead Development; Nanomaterial for Selective Electrochemical Sensing	27
2.1	INTRODUCTION.....	27
2.2	EXPERIMENTAL	30
2.2.1	Materials	30
2.2.2	EETMSi mediated synthesis of PBN	30
2.2.3	Fabrication of PBN encapsulated silica-alginate beads.....	30
2.2.4	Instrumentation.....	30
2.2.5	Kinetic assay	31
2.2.6	Catalytic activity of PBN as peroxidase.....	31
2.2.7	Electrochemical sensing of H ₂ O ₂	32
2.2.8	Electrochemical oxidation of pyrogallol through PBN encapsulated silica-alginate beads	32
2.3	RESULT AND DISCUSSION.....	32
2.3.1	Effect of alkoxysilane on PBN synthesis.....	32
2.3.2	Effect of EETMSi over charge transfer spectra of PBN	38
2.3.3	IR spectrum of PBN	39
2.3.4	Chemical approach of EETMSi and THF-HPO during PBN fabrication	39
2.3.5	Stability and solubility of nanoparticle	41
2.3.6	Electrochemical behavior of PBN	41
2.3.7	Electrochemical sensing of H ₂ O ₂	42
2.3.8	Peroxidase mimetic activity of PBN	45
2.3.9	Development of PBN encapsulated silica-alginate beads.....	45

2.3.10	Electrochemical sensing of pyrogallol using PBN encapsulated silica-alginate beads	48
2.4	CONCLUSION	52
3	Synthesis and Application of Functional Prussian Blue Nanoparticles; A Photo-Fenton Catalyst for Dye	55
3.1	INTRODUCTION	55
3.2	EXPERIMENTAL	57
3.2.1	Materials.....	58
3.2.2	Instrumentation.....	58
3.2.3	Experimental setup and measurement of photon flux	58
3.2.4	Photocatalytic assay of RhB using PBN	59
3.2.5	EPR analysis of generated hydroxyl radical.....	59
3.2.6	Preparation of PBN.....	60
3.2.7	Characterization and analysis of photo-Fenton catalyst.....	60
3.3	RESULT AND DISCUSSION.....	73
3.3.1	Photowork for RhB degradation	73
3.3.2	Kinetic study of photo-Fenton degradation process.....	81
3.3.3	Rhodamine B photodegradation	87
3.3.4	BET surface area analysis of unused and recycled catalyst	94
3.3.5	Anticipated degradation mechanism of photo-Fenton process	94
3.3.6	Assessment of photo-Fenton process over polluted river water	96
3.3.7	Study of recyclability of catalyst after dye degradation	97
3.3.8	Characterization of catalyst after post-degradation.....	97
3.4	CONCLUSION	99
4	Synthesis of Prussian Blue Nanoparticles-Embedded Heterogeneous Catalyst for Hydrogen Peroxide Detection	101
4.1	INTRODUCTION	101
4.2	EXPERIMENTAL	103
4.2.1	Materials.....	103
4.2.2	EETMSi mediated synthesis of PBN and PBN@SiO ₂	103

4.2.3	Instrumentation.....	104
4.3	RESULT AND DISCUSSION.....	104
4.3.1	Study of the electrochemical behavior of PBN through cyclic voltammetry.....	104
4.3.2	Electrochemical sensing of H ₂ O ₂ through PBN.....	105
4.3.3	SEM analysis of fabricated PBN@SiO ₂	107
4.3.4	Mapping analysis.....	108
4.3.5	FTIR analysis.....	108
4.3.6	XRD analysis.....	109
4.3.7	XPS analysis.....	110
4.3.8	Peroxidase mimetic activity of PBN@SiO ₂	112
4.4	CONCLUSION.....	114

5 Palladium-Prussian Blue Nanoparticles; As Homogeneous and Heterogeneous Electrocatalysts **115**

5.1	INTRODUCTION.....	115
5.2	EXPERIMENTAL.....	116
5.2.1	Materials.....	116
5.2.2	Synthesis of PBN and its nanodispersion with PdNP.....	117
5.2.3	Electrochemical setup.....	118
5.2.4	Instrumentation and kinetic assay.....	118
5.2.5	Hydrogen peroxide detection using GOx as homogeneous catalyst.....	118
5.3	RESULT AND DISCUSSION.....	119
5.3.1	Investigation of nanoparticles synthesis through UVVis spectroscopy..	119
5.3.2	TEM analysis.....	119
5.3.3	Electrocatalytic behavior of PB-PdNP immobilized on carbon electrode.....	123
5.3.4	Electrocatalytic reduction of H ₂ O ₂ at modified electrode.....	124
5.3.5	Glucose detection using GOx in homogeneous system.....	129
5.3.6	Glucose sensing based on GOx and PB-PdNP ₂ modified electrode....	129
5.3.7	Stability and reproducibility of modified electrodes.....	131
5.4	CONCLUSION.....	133

SUMMARY	135
FUTURE PROJECTION	139
Bibliography	143
List of Publication	179

List of Tables

1.1	Electrochromic behaviour of different MHCFs.....	8
2.1	IR data listed with assigned possible functional groups.....	41
2.2	Kinetic parameters (K_m and V_{max}) justifying peroxidase like activity of various Catalysts.....	45
3.1	Comparative study of RhB degradation under different conditions.	77
3.2	Kinetic analysis of different degradation system.....	85
3.3	Data listed here displaying the amount of H_2O_2 concentration used in degradation process based on the coloured complex formation.	87
3.4	Organic end product identified from LC-MS spectrum after photodegradation of RhB.....	91
3.5	Band gap analysis of all synthesized Nanoparticles.....	96
3.6	Comparative study of previously reported system with this catalyst.....	99
5.1	Comparison of different modified electrodes in the determination of H_2O_2 ...	128
5.2	Comparison of the calculated kinetic parameters for PB-PdNP with earlier reported materials.	130

List of Figures

1.1	3-D framework of soluble (A) and insoluble PB (B); demonstrated with Fe(II) [red], Fe(III) [yellow], C [golden], N [grey], K [purple], and H ₂ O molecule [green colour]	2
1.2	Electronic arrangement of valence electrons of Fe(II) and Fe(III) in octahedral environment of carbon and nitrogen hole respectively	4
1.3	Cyclic voltammogram of a PB modified electrode showing oxidation and reduction peak [Ricci and Palleschi, (2050)].	5
2.1	UV-Vis spectra of; (a) 0.1 M potassium ferricyanide (b) reaction mixture containing 0.1 M ferricyanide, and 1 M THF-HPO (c) reaction mixture containing 0.1 M ferricyanide, 1 M THF-HPO, and 0.03 M EETMSi after incubation at 60°C for 5 h. The inset shows the visual images (a,b,and c) of correspondings.....	33
2.2	PBN formation, based on THF-HPO concentration variations; (a) 1 M (b) 2 M (c) 3 M (d) 4 M, at constant ferricyanide (0.1 M) and EETMSi (0.03 M) content.	34
2.3	UV-Vis spectra of PBN synthesis, based on concentration variation of ferricyanide (a) 0.05 M (b) 0.1 M (c) 0.5 M, at constant THF-HPO (1 M) and EETMSi (0.03 M).....	35
2.4	Study of PBN formation, based on concentration variation of EETMSi; (a) 0.0 M (b) 0.03 M (c) 0.06 M and (d) 0.12 M, at constant THF-HPO (1 M) and ferricyanide (0.1 M).....	35
2.5	TEM and visual image of PBN-1 (A), SAED pattern (B) of corresponding nanoparticles (PBN-1).	36

2.6	TEM and visual image of PBN-2 (A), SAED pattern (B) of corresponding nanoparticles (PBN-2).....	36
2.7	TEM and visual image of PBN-3 (A), SAED pattern (B) of corresponding nanoparticles (PBN-3). Inset (a) displays the nanoparticle at high magnification (scale bar of 50 nm).	37
2.8	Bar histograms showing the particle size distribution of PBN-1 (A), PBN-2 (B), and PBN-3 (C).	37
2.9	Dependence of zeta potential on particle diameter of PBN (A). Relation of nanoparticle diameter with EETMSi /Fe ⁺³ stoichiometric ratio (B).....	38
2.10	FTIR analysis of as-synthesized PBN (A) with THF-HPO (B), and EETMSi (C) through ATR.	40
2.11	ATR spectrum of as-synthesized PBN (A), THF-HPO (B), and EETMSi (C).....	40
2.12	Dispersibility of as-synthesized PBN in different solvents.....	42
2.13	Study of electrochemical nature of PBN-1 (A), PBN-2 (B), and PBN-3 (C) modified electrodes in 0.1 M phosphate buffer (pH 7.0) at the scan rate of 5 mV/s.....	43
2.14	Cyclic voltammograms of PBN-1 (A), PBN-2 (B), and PBN-3 (C) modified systems in the absence (i) and presence (ii) of 1 mM H ₂ O ₂ in 0.1 M phosphate buffer (pH-7.0) containing 0.5 M KCl.....	44
2.15	UV-Vis spectra of o-dianisidine-H ₂ O ₂ -PBN system displaying change in absorbance as a function of H ₂ O ₂ concentration variations catalyzed by PBN-1 (A), PBN-2 (B), and PBN-3 (C).	46
2.16	Time dependent absorbance changes at 430 nm in the presence of different concentration (0.05 to 25 mM) of H ₂ O ₂ and fixed concentration of o-dianisidine (50 μM) catalyzed by PBN-1 (A), PBN-2 (B), and PBN-3 (C) respectively	46
2.17	Kinetic analysis of PBN-1 (A), PBN-2 (B), and PBN-3 (C) with H ₂ O ₂ as a substrate.....	47
2.18	SEM images of calcium-alginate bead (A), silica-alginate bead (B), PBN-encapsulated silica-alginate bead (C), and PBN-3 (D).....	49

2.19	EDX spectra of PBN-encapsulated silica-alginate beads identify the presence of all elemental species.....	49
2.20	Differential pulse voltammogram justifying auto-oxidation of pyrogallol individually (A) in 0.1 M phosphate buffer (pH-7.0) containing 0.5 M KCl. UV-Vis spectrum of the resultant system (B).....	50
2.21	Differential pulse voltammogram justifying auto-oxidation of pyrogallol, in the presence of H ₂ O ₂ (A) in 0.1 M phosphate buffer (pH-7.0) containing 0.5 M KCl. And UV-Vis spectrum of the resultant system (B).	50
2.22	Differential pulse voltammogram justifying auto-oxidation of pyrogallol, in the presence of H ₂ O ₂ and PB-encapsulated silica-alginate beads (A) in 0.1 M phosphate buffer (pH-7.0) containing 0.5 M KCl. The UV-Vis spectrum of resultant system displaying increase in absorption maxima as function of pyrogallol concentration (B).	51
2.23	Plausible mechanism of Pyrogallol oxidation to purpurogallin via PB-encapsulated silica-alginate beads with H ₂ O ₂	51
2.24	Catalytic oxidation of pyrogallol to purpurogallin through PBN doped silica-alginate bead.	53
3.1	Recorded UV-Vis spectra of potassium ferricyanide (a) and synthesized PBN (b).....	60
3.2	Recorded UV-Vis spectra of PBN synthesis (A) at different concentrations of 100 μ l of ferricyanide; (a) 0.05 M (b) 0.04 M (c) 0.03 M (d) 0.02 M and (e) 0.01 M with constant EETMSi (20 μ l, 0.05 M) and cyclohexanone (20 μ l, 9.8 M) contents, and (B) via varying cyclohexanone amount; (a) 25 μ l, (b) 20 μ l, (c) 15 μ l and (d) 10 μ l; (e) 5 μ l at constant EETMSi (20 μ l, 0.05 M) and ferricyanide content (100 μ l, 0.03 M).....	62
3.3	Spectrum recorded at different concentrations of 20 μ l of EETMSi; (a) 0.5 M (b) 0.1 M (c) 0.05 M (d) 0.01 M with a fixed ferricyanide content (100 μ l, 0.03 M) and cyclohexanone (20 μ l, 9.8 M).....	63
3.4	P-XRD of as prepared PBN ₁ (A), PBN ₂ (B), PBN ₃ (C), and PBN ₄ (D).....	64
3.5	FTIR spectrum of as-synthesized PBN indicating functional species.....	65
3.6	Elemental confirmation was characterized by EDX.....	66

3.7	TEM images of PBN ₁ (A,B), and the image shown in inset of (A) displayed lattice fringes of the corresponding nanoparticles.	67
3.8	SAED pattern of PBN ₁ (A), and bar histogram displaying the particle size distribution of corresponding nanoparticles (B).	67
3.9	TEM images of PBN ₂ (A,B), and the image shown in inset of (A) displayed lattice fringes of the corresponding nanoparticles.	68
3.10	SAED pattern of PBN ₂ (A), and bar histogram displaying the particle size distribution of corresponding nanoparticles (B).	68
3.11	TEM images of PBN ₃ (A,B), and the image shown in inset of (A) displayed lattice fringes of the corresponding nanoparticles.	69
3.12	SAED pattern of PBN ₃ (A), and bar histogram displaying the particle size distribution of corresponding nanoparticles (B).	69
3.13	TEM images of PBN ₄ (A,B), and the image shown in inset of (A) displayed lattice fringes of the corresponding nanoparticles.	70
3.14	SAED pattern of PBN ₄ (A), and bar histogram displaying the particle size distribution of corresponding nanoparticles (B).	70
3.15	XPS analysis of PBN exhibiting presence of Fe ⁺² (BE of 721.87 eV and 708.64 eV) and Fe ⁺³ (B.E. of 712.82 eV) chemical species in PBN.....	71
3.16	Measured zeta potentials of differently sized nanoparticles; PBN ₁ (A), PBN ₂ (B), PBN ₃ (C), and PBN ₄ (D) via DLS techniques.	72
3.17	UV-Vis spectrum of aqueous RhB solution (10 ppm) degradation using PBN/H ₂ O ₂ (0.5 mM/10 mM) in different condition; RT (A), oven (B), UV-lamp (C), and solar light (D).	74
3.18	Recorded UV-Vis spectrum (with magnified view) of RhB during photodegradation under solar light in different systems; RhB (10 ppm) (A, A-I), RhB (10 ppm)-H ₂ O ₂ (1.0 mM) (B, B-I), RhB (10 ppm)-PBN (0.5 mM) (C, C-I), and RhB (10 ppm)-H ₂ O ₂ (10 mM)-PBN (0.5 mM) (D, D-I).	75
3.19	Effect of the irradiation time on photocatalytic degradation process recorded in different reactant conditions.	76

3.20	Effect of PBN concentration variation (A = 0.5 mM, B = 1.0 mM, C = 1.5 mM) with continual concentration of H ₂ O ₂ (10 mM), and effect of H ₂ O ₂ concentration variation (D = 10 mM, E = 20 mM, F = 30 mM) with continual concentration of PBN (0.5 mM) on solar light assisted photo-Fenton degradation of RhB (10ppm) in RhB-PBN-H ₂ O ₂ system as recorded in UV-Vis spectrum.....	78
3.21	Concentration variation influence of catalyst PBN (0.5 mM-1.5 mM) with continual H ₂ O ₂ (10 mM) and RhB (10 ppm) (A), and H ₂ O ₂ (10 mM-30 mM) with continual PBN (0.5 mM) and RhB (10 ppm) (B) on the irradiation time of photocatalytic degradation process in RhB-H ₂ O ₂ -PBN system.	79
3.22	Size influence of photo-Fenton catalyst PBN ₁ (A), PBN ₂ (B), PBN ₃ (C), and PBN ₄ (D) over RhB (10ppm) degradation with H ₂ O ₂ (10 mM) in RhB-H ₂ O ₂ -PBN system.	80
3.23	Relationship between unreacted fraction of RhB of different concentration (10-50 ppm) vs. irradiation time using constant H ₂ O ₂ (10 mM) (A). Linear relation between Ln(α) vs. degradation time for different concentration of RhB with H ₂ O ₂ (B).	81
3.24	Relationship between unreacted fraction of RhB (10 ppm) vs. irradiation time with varying H ₂ O ₂ (10mM-30mM) concentrations (A). Linear relationship between Ln(α) vs. degradation time for different concentration of H ₂ O ₂ (B). And plot of Ln (k) vs. Ln (H ₂ O ₂) for RhB photodegradation (C).	83
3.25	Relationship between unreacted fraction of RhB (10 ppm) vs. irradiation time using constant H ₂ O ₂ (10 mM) and different catalyst (0.5 mM-1.5 mM) concentrations (A). Linear relationship between ln(α) vs. degradation time in different concentration of PBN (0.5 mM-1.5 mM) with constant H ₂ O ₂ (10 mM) and RhB (10 ppm) (B). Plot of Ln (k) vs. Ln (PBN) for RhB photodegradation (C).....	84

3.26	Recorded absorbance maxima ($\lambda_{\text{max}} = 480 \text{ nm}$) as a function of H_2O_2 (1mM-30mM) concentrations correspond to the oxidized product of o-dianisidine dye (25 μM) in the presence of PBN (1 mM) (A). Calibration curve displaying H_2O_2 concentration (1 mM-30 mM) dependent change in the absorbance maxima at 480 nm (B). And change in absorption for o-dianisidine-PBN- H_2O_2 complex with respect to residual amount of H_2O_2 (10 mM, 20 mM, 30 mM) after photodegradation process (C).	86
3.27	Hypsochromic (blue shift) and hypochromic shift in absorption maxima peak of RhB during the photodegradation process in acidic (A), and alkaline (B) medium	88
3.28	LC-MS spectra correspond to RhB (A), post-degraded end product of RhB after degradation in acidic (B), and alkaline medium (C).	89
3.29	Photodegradation pathways of RhB in the presence of catalyst PBN and H_2O_2 via photo-Fenton process.....	90
3.30	EPR spectra of DMPO-OH adduct of intensity 1:2:2:1 (A), and TEMPO- O_2 adduct of intensity 1:1:1 (B) correspond to generated hydroxyl and superoxide radicals throughout photo-Fenton process.	92
3.31	N_2 adsorption-desorption isotherms of unused (A), and recycled PBN catalyst (B). And BJH plot displaying pore diameter distribution curve for unused (A-I) and recycled PBN catalyst (B-I).....	93
3.32	Study of recyclability of photo-Fenton catalyst via five subsequent repetitive cycles of RhB (10ppm) degradation.	97
3.33	Comparative study of regenerated catalyst with unused one via ATR technique, after repetitive (five times) photodegradation process.....	98
3.34	TEM image (A) and corresponding SAED (B) of reprocessed PBN_4 after repetitive (five times) photodegradation process. The histogram (in the inset of A) revealed the size of recycled nanoparticles.....	98
4.1	Recorded cyclic voltammogram of PBN in 0.1 M KNO_3 solution.	105
4.2	Study of electrochemical response of PBN in the presence of H_2O_2 (0-10mM) by DPV (A). Concentration dependent calibration curve for H_2O_2 (B).....	106

4.3	HR SEM images of PBN@SiO ₂ at different magnification displayed the successful formation of PBN over SiO ₂ (A-C), the visual images; (I) and (II) in inset of Fig. (A), corresponds to SiO ₂ and PBN@SiO ₂ respectively. The EDX spectrum revealed the presence of all mandatory elements (D). Bar histogram displaying the particle size distribution curve of corresponding nanoparticles (E).....	107
4.4	Mapping analysis of selected area in PBN@SiO ₂ (A) identified to the presence of all elemental species; carbon (B), nitrogen (C), oxygen (D,) iron (E), and silicon (F) with their relative quantity	108
4.5	FTIR spectrum of as-synthesized PBN@SiO ₂ and heterogeneous support SiO ₂	109
4.6	P-XRD of SiO ₂ and as-synthesized PBN@SiO ₂ unveiling the presence of planes.....	110
4.7	XPS analysis of PBN@SiO ₂ exhibiting complete survey scan with all recognized species (A), Fe ⁺² and Fe ⁺³ species in PBN@SiO ₂ (B), and identified chemical states of Si (IV) in PBN@SiO ₂ (C).....	111
4.8	UV-Vis spectra exhibiting peroxidase like activity of PBN@SiO ₂ towards H ₂ O ₂ by oxidizing chromogenic substrate (o-dianisidine).	113
4.9	Recorded UV-Vis spectra displaying change in the absorption maxima correspond to o-dianisidine oxidation with respect to H ₂ O ₂ concentration variation in the presence of PBN@SiO ₂ catalyst.....	113
5.1	Absorption spectra of PBN (1), PdNP (2), and PB-PdNP (3); the visual images (i, ii and iii) of corresponding nanoparticles.	120
5.2	TEM and SAED pattern of PdNP ₁ (A) and PdNP ₂ (B).....	120
5.3	TEM images of sequentially synthesized PB-PdNP ₁ (A,B,C) and PB-PdNP ₂ (D,E,F) nanoparticles.	121
5.4	Mapping analysis and EDX spectrum of PB-PdNP ₁ (A), and PB-PdNP ₂ (B) shows the presence of Fe, Pd and Si elements.....	122
5.5	Cyclic voltammograms of PBN (A), PB-PdNP ₁ (B), and PB-PdNP ₂ (C) at various scan rate of 0.01 to 0.3 Vs ⁻¹ in 0.1 M KNO ₃	123

5.6	Recorded cyclic voltammograms of PBN (A), PB-PdNP ₁ (B), and PB-PdNP ₂ (C) modified systems in the absence (1) and presence (2) of 1 mM H ₂ O ₂ in 0.1 M phosphate buffer.	125
5.7	Amperometric response of PBN (1), PB-PdNP ₁ (2), and PB-PdNP ₂ (3) modified systems, on the addition of varying concentrations of H ₂ O ₂ (0.01 μM to 5 mM) at 0.0 V vs. Ag/AgCl, in 0.1 M phosphate buffer (pH-7.0) containing 0.5 M KCl. Inset shows the corresponding calibration curves for H ₂ O ₂	126
5.8	Time dependent absorbance variation at 430 nm, depending on H ₂ O ₂ (0.03 to 25 mM) catalytic oxidation by PB-PdNP ₁ (A), and PB-PdNP ₂ (B). Kinetic analysis of PB-PdNP ₁ (A-I), and PB-PdNP ₂ (B-I) with H ₂ O ₂ as a substrate in phosphate buffer (pH-7.0).....	127
5.9	Concentration-response curve for glucose detection using GOx enzyme and o-dianisidine in the presence of PB-PdNP ₂ (A). The linear calibration curve of glucose (A-I).	131
5.10	Cyclic voltammetry of PBN-GOx (A) and PB-PdNP ₂ -GOx modified enzyme electrode (B) in absence (1) and the presence (2) of 200 mM glucose in 0.1 M phosphate buffer (pH 7.0) at the scan rate of 10 mV/s. Amperometric responses recorded upon addition of increasing concentrations of glucose for PBN (C) and PB-PdNP ₂ (D) modified enzyme electrodes at constant potential of +0.6 V vs. Ag/AgCl in 0.1 M phosphate buffer (pH 7.0).....	132