

Table of contents

Acknowledgement	xiii
Abstract	xvi
List of figures	xxi
List of tables	xxv
List of Symbols	xxx
1 Introduction	1
1.1 A very brief history of the Universe	2
1.1.1 21 cm signal emission from Neutral Hydrogen	5
1.1.2 Detection of neutral hydrogen at high redshift	8
1.2 Gravitational Lensing	10
1.2.1 Lensing Basic	11
1.2.2 H I observation with gravitational lensing	13
1.3 Plan of the Thesis	15
2 Post-Reionization H I power spectrum and detection challenges	17
2.1 Dark Matter power spectrum	17
2.2 H I power spectrum	19

2.2.1	Modelling redshifted 21 cm power spectrum	20
2.2.2	21 cm Angular power spectrum	22
2.3	Radio interferometer as a probe of 21 cm power spectrum	24
2.4	Effect of foreground	24
2.5	Discussion	27
3	Using strong gravitational lensing to probe the post reionization H I power spectrum	29
3.1	Formalism	30
3.1.1	Visibility function of lensed H I emission	30
3.1.2	Power spectrum for the lensed H I emission	32
3.2	Simulating lensed 21 cm power spectrum	33
3.2.1	Model for Lensing Potential	33
3.2.2	Point spread function of the gravitational lens	36
3.3	Results	37
3.4	Discussion and Conclusion	40
4	Measuring redshifted 21 cm signal using Combine lense estimator	43
4.1	Combined Lensed Estimator	44
4.1.1	Effect of Strong Lensing	44
4.1.2	Power Spectrum Estimator	45
4.2	Implementation of the Estimator	48
4.2.1	Calculation of S_L in the observed baseline positions	48
4.2.2	Gridding the observed lens visibilities	49
4.2.3	Estimates of C_l and its variance	49
4.3	Simulations and Results	50
4.3.1	Lens Model	50

Table of contents	xix
4.3.2 Results	53
4.3.3 Effect of Foreground	58
4.4 Discussion and Conclusion	59
5 Summary and conclusion	61
5.1 Limitations and Future Scope	62
References	65