

List of Figures

Figure 1.1 Crystal structure of perovskite ABO_3	2
Figure 1.2 Representation of the d orbitals.	4
Figure 1.3 d orbitals splitting in octahedral crystal field in perovskite.	5
Figure 1.4 Jahn-Teller distortion in Mn^{3+} ions in rare earth manganites [6].	6
Figure 1.5 Schematic diagram of magnetic dipole moments of ferromagnetic material aligned in single direction.	8
Figure 1.6 Curie – Weiss Law for ferromagnetic materials.	9
Figure 1.7 Schematic plot of magnetic hysteresis loop for ferromagnetic material [9].	10
Figure 1.8 Schematic diagram of magnetic dipole moment of anti-ferromagnetic material align in opposite direction.	11
Figure 1.9 Different types of antiferromagnetism.	12
Figure 1.10 Curie –Weiss law for antiferromagnetic materials [10].	12
Figure 1.11 Bethe–Slater curve for ferromagnetic and antiferromagnetic materials [8].	15
Figure 1.12 Schematic representation of super-exchange interaction in magnetic oxides. The p-orbital of anion interact with d orbital of transition metal cations [11].	16
Figure 1.13 Schematic diagram for double exchange interaction.	17
Figure 1.14 Crystal structure of $GdMnO_3$ [29].	21
Figure 1.15 Crystal structure of $LaMnO_3$ [61].	26
Figure 1.16 Schematic diagram of negative magnetization [68].	28
Figure 2.1 Flow chart for the synthesis of pure $GdMnO_3$ using sol-gel technique.	34
Figure 2.2 Process flow of co-precipitation method for the synthesis of pure $LaMnO_3$	35
Figure 3.1 Powder X-ray diffraction patterns of $GdMn_{1-x}Fe_xO_3$ ($x = 0, 0.1$ and 0.2).	49

Figure 3.2 Rietveld refinement of X-ray diffraction (XRD) patterns of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ (a) $x = 0$, (b) $x = 0.1$ and (c) $x = 0.2$	49
Figure 3.3 Crystal structure obtained from Rietveld refinement drawn using Vesta software.	52
Figure 3.4 Mn-O bond lengths of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2) obtained from Rietveld refinement.	53
Figure 3.5 Raman spectra of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2).	56
Figure 3.6 (a) X-ray photoelectron spectroscopy for Mn 2p and (b) O1s of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2).	58
Figure 3.7 (a) Field emission scanning electron micrographs of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2), (b) shows the corresponding particle size distribution histogram.	58
Figure 3.8 (a) Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 0.050 kOe for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2) (b) Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 10 kOe for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2).	61
Figure 3.9 The inverse susceptibility, $1/\chi$ plots of ZFC and FC in the presence of 0.050 kOe of the $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2).	62
Figure 3.10 Fitting of $1/\chi$ vs T with Curie- Weiss law of the $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2).	64
Figure 3.11 Magnetic field-dependent magnetization at 20 and 4 K of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2) with zoomed view.	65
Figure 3.12 Temperature dependence of the real part of ac susceptibility at various frequencies at the 3Oe ac field (a) $x= 0$ (b) $x= 0.1$ and (c) $x= 0.2$. The right panel of the respective figure shows the expanded view of the peak shift with increasing frequency....	69
Figure 3.13 Variation of relaxation time (τ) with T plotted with the Neel–Arrhenius fitting equation for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2). The solid line represents the fitting.	70
Figure 3.14 Variation of relaxation time (τ) with T plotted with vogel-fulcher (equation 3.4) and the power law (equation 3.5) for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x= 0, 0.1$ and 0.2). The solid line represents the fitting with experimental data point.	71
Figure 4.1 Field emission scanning electron micrographs of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	74

Figure 4.2 X-ray diffraction (XRD) patterns of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5) fitted using Rietveld refinement of Fullprof programe.....	75
Figure 4.3 Crystal structure obtained from Rietveld refinement drawn using Vesta software of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).....	78
Figure 4.4 Bond- lengths obtained from Rietveld refinement drawn using Vesta software of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).....	79
Figure 4.5 Raman spectra of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	80
Figure 4.6 Room temperature emission spectra of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0, 0.3$ and 0.5) at an excitation wavelength 200 nm.	81
Figure 4.7 Room temperature excitation spectra of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0, 0.3$ and 0.5) at an emission wavelength 466 nm.	84
Figure 4.8 Energy band diagram of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	85
Figure 4.9 X-ray photoelectron spectroscopy for Mn 2p of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).....	87
Figure 4.10 X-ray photoelectron spectroscopy for Fe 2p of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).....	88
Figure 4.11 X-ray photoelectron spectroscopy for O1s of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	89
Figure 4.12 Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 0.050 kOe for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	91
Figure 4.13 Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 10 kOe for $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5).	92
Figure 4.14 The inverse susceptibility, $1/\chi$ plots of ZFC and FC in the presence of 0.050 kOe of $x = 0.3$ and 0.5	93
Figure 4.15 Irreversibility magnetization vs temperature for $x = 0.3$ and 0.5	94
Figure 4.16 Magnetic field-dependent magnetization at 4 K and 20 K of $\text{GdMn}_{1-x}\text{Fe}_x\text{O}_3$ ($x = 0.3$ and 0.5) and zoom view of Magnetic field-dependent magnetization.	96
Figure 4.17 Temperature dependence of the real part of ac susceptibility at various frequencies at the 3 Oe ac field of $x = 0.3$ and $x = 0.5$	98

Figure 4.18 Magnetic relaxation after field cooling in the presence of 50 Oe. The red line shows the fitting to the power law function.	99
Figure 5.1 Rietveld refinement of X-ray diffraction (XRD) patterns of $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x=0.1, 0.2, 0.3$ and 0.4).	102
Figure 5.2 Variations in lattice parameters and volume with Cr concentration.	104
Figure 5.3 Crystal structure obtained from Rietveld refinement drawn using Vesta software.	105
Figure 5.4 Raman spectra of $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x= 0.1, 0.2, 0.3$ and 0.4).	107
Figure 5.5 The emission spectra of $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x=0, 0.3$ and 0.4) under the excitation wavelength of 200 nm. (Inset depicts the deconvoluted peaks).	108
Figure 5.6 The emission spectra of $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x=0, 0.1, 0.2, 0.3$ and 0.4) at (a) $\lambda_{\text{ex}}=460$ nm, (b) $\lambda_{\text{ex}}= 582$ nm, (c) $\lambda_{\text{ex}}= 625$ nm and (d) the excitation spectra after monitoring $\lambda_{\text{em}}= 692$ nm.	110
Figure 5.7 The proposed energy band diagram for Cr doped GdMnO_3	111
Figure 5.8 Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 500 for $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x=0.1- 0.4$).	114
Figure 5.9 Zoom part of Temperature dependent magnetization under zero (ZFC) and field cooling (FC) measured at 500 Oe and $(M_{\text{ZFC}}-M_{\text{FC}})/M_{\text{ZFC}}$ vs T for $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x= 0.3$).	115
Figure 5.10 Schematic diagrams of the Cr^{3+} moment (M_{Cr}) and Gd^{3+} moments under different temperature range.	117
Figure 5.11 Temperature dependent magnetization under field cooling (FC) measured at different fields for $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x= 0.3$ and 0.4).	117
Figure 5.12 Magnetization curves at 100 Oe applied field in FC mode (the red line shows fitting).	118
Figure 5.13 Mirror like behavior in FC mode for $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x= 0.3$ and 0.4).	120
Figure 5.14 The inverse susceptibility, $1/\chi$ plots of ZFC and FC in the presence of 500 Oe of the $\text{GdMn}_{1-x}\text{Cr}_x\text{O}_3$ ($x= 0.1- 0.4$).	121

Figure 5.15 Curie- Weiss law fitting in the presence of 500 Oe of the $GdMn_{1-x}Cr_xO_3$ ($x= 0.1- 0.4$).....	122
Figure 5.16 Magnetic field dependent magnetization at 4 and 2 K of $GdMn_{1-x}Cr_xO_3$ ($x= 0.1-0.4$) with zoomed view.	123
Figure 5.17 Magnetic relaxation after field cooling in 100 Oe. The red line shows the fitting of experimental data with power law.....	125
Figure 5.18 Magnetic switching behavior of $GdMn_{1-x}Cr_xO_3$ ($x= 0.3$ and 0.4) at different field.....	126
Figure 6.1 Rietveld refinement of X-ray diffraction (XRD) patterns (a) LaMnO1 and (b) LamnO2 calcined at $900^\circ C$	130
Figure 6.2 Crystal structure obtained from Rietveld refinement drawn using Vesta software (a) LaMnO1 and (b) LaMnO2.	132
Figure 6.3 (a) Field emission scanning electron micrographs of LaMnO1 and LaMnO2 (b) shows the corresponding particle size distribution histogram.	134
Figure 6.4 (a) X-ray photoelectron spectroscopy for Mn 2p and (b) O1s of of LaMnO1 and LaMnO2.....	135
Figure 6.5 (a) Temperature dependent magnetization under zero field cooling (ZFC) and field cooling (FC) measured at 100 Oe and at 10 kOe for LaMnO1and LaMnO2.....	137
Figure 6.6 Curie- Weiss law fitting of the LaMnO1 and LaMnO2.....	138
Figure 6.7 Magnetic field dependent magnetization at 300 K, 120 K, 100 K and 5 K of LaMnO1 and LaMnO2.	141
Figure 6.8 Temperature dependence of the real and imaginary part of χ susceptibility at various frequencies at the 3Oe ac field for LaMnO1 and LaMnO2. The right panel of the respective figure shows the expanded view of the peak shift with increasing frequency. .	143
Figure 6.9 Variation of relaxation time (τ) with T plotted with Neel-Arrhenius, Vogel-Fulcher and the Power law fitting for LaMnO1 and LaMnO2. The solid line represents the fitting.	145