

# Table of contents

<b>List of figures</b>	<b>xvii</b>
<b>1 Introduction</b>	<b>1</b>
1.1 Statistical mechanics:macro to micro . . . . .	1
1.2 Ensembles, phase space and thermodynamic averages . . . . .	2
1.3 Active matter systems: A class of nonequilibrium systems . . . . .	5
1.4 Classification of active matter system . . . . .	8
1.4.1 Polar, apolar and active Brownian particles . . . . .	8
1.4.2 Dry and wet active system . . . . .	10
1.5 Methodology . . . . .	11
1.5.1 Agent-based simulation . . . . .	11
1.5.2 Phenomenological approach : hydrodynamic equations of motion	17
1.6 Coarsening . . . . .	21
1.6.1 Coarsening in systems with nonconserved kinetics (Model A) . .	21
1.6.2 Coarsening in systems with conserved kinetics (Model B) . . . .	23
1.6.3 Correlation function and structure factor . . . . .	25
1.6.4 Coarsening in active systems . . . . .	26
1.7 Organisation of the thesis . . . . .	27
1.8 Technical details . . . . .	29

---

<b>2</b>	<b>Activity Driven Phase Separation and Ordering Kinetics Of Passive Particles</b>	<b>31</b>
2.1	Introduction . . . . .	31
2.2	Model . . . . .	34
2.3	Results . . . . .	37
2.4	Characteristics of steady state . . . . .	38
2.4.1	Radial distribution function ( <i>RDF</i> ) $g_{pp}(r)$ . . . . .	38
2.4.2	Cluster size distribution ( <i>CSD</i> ) . . . . .	40
2.5	Growth kinetics . . . . .	41
2.6	Conclusion . . . . .	44
<b>3</b>	<b>Obstacle induced phase separation of micro swimmers</b>	<b>47</b>
3.1	Introduction . . . . .	47
3.2	Model . . . . .	50
3.3	Results . . . . .	52
3.3.1	Phase Separation . . . . .	52
3.3.2	Steady state phase diagram . . . . .	55
3.4	Conclusion . . . . .	56
<b>4</b>	<b>Role of thermal noise for the growth kinetics of active model B (Critical mixture)</b>	<b>59</b>
4.1	Introduction . . . . .	59
4.2	Model . . . . .	61
4.3	Results . . . . .	64
4.3.1	Scaling Theory . . . . .	72
4.4	Discussion . . . . .	75
<b>5</b>	<b>Role of thermal noise for the growth kinetics of active model B (off-critical mixture)</b>	<b>77</b>

Table of contents	xv
<hr/>	
5.1 Introduction . . . . .	77
5.2 Results . . . . .	80
5.2.1 Slightly off-critical composition (same sign, $\lambda \psi_0 > 0$ ) . . . . .	80
5.2.2 Slightly off-critical composition (opposite sign, $\lambda \psi_0 < 0$ ) . . . . .	84
5.2.3 Deep off-critical composition . . . . .	85
5.3 Conclusion . . . . .	87
<b>6 Ordering kinetics in active polar fluid</b>	<b>89</b>
6.1 Introduction . . . . .	89
6.2 Model . . . . .	92
6.3 Results . . . . .	98
6.4 Conclusions . . . . .	103
<b>7 Summary and conclusion</b>	<b>105</b>
7.1 Summary . . . . .	105
7.2 Future aspects . . . . .	107
<b>References</b>	<b>109</b>
<b>List of Publications</b>	<b>i</b>



# List of figures

1.1	(color online) Schematic diagram shows the polar, apolar and active Brownian particle and the different ordering exhibited by them. Arrow shows their direction of self propulsion. Head and tail symmetry is depicted by double arrows. . . . .	9
2.1	(color online) Schematic diagram of the model. It shows the initial homogeneous state of the system. Here the colours and sizes represent two types of particles. Bigger particles (red) are active particles and smaller particles (grey) are passive particles. Small white dot on red particles denote the instantaneous orientation direction $\theta$ of ABPs. . . . .	36
2.2	(color online) The real space snapshots of local density of passive particles at different times. (a-c) are for size ratio 10 and $\bar{v}= 100, 300$ and 600 respectively. (d-f) are for size ratio 6 and $\bar{v}= 100, 300$ and 600 respectively. From bottom to top panels are for times, $t = 0.05, t = 0.5$ and $t = 30.0$ . The color bar shows the local density of passive particles.(g) The lower color plot shows the phase separation order parameter ( <i>PSOP</i> ) in the plane of size ratio and activity $(S, \bar{v})$ .(h-i) Probability distribution function (PDF) of local density of passive particles $P(\rho_p)$ is plotted for two different size ratios 10 and 6 ((h) and (i)) respectively and for three $\bar{v}= 100, 300$ and 600.	38

- 2.3 (color online) Passive-passive radial distribution function (*RDF*),  $g_{pp}(r)$  is plotted for different parameters. (a-c) is for fixed activity and varying size ratio.  $\bar{v}$  is varied for three different values, 100, 300 and 600. (a) is for  $\bar{v} = 100$  (b) is for  $\bar{v} = 300$  and (c) is for  $\bar{v} = 600$  and size ratio  $S$  is taken 6, 8 and 10 for black  $\circ$ , red  $\square$  and green  $\diamond$  respectively. inset figures are zoomed for the same parameters as mentioned in (a), (b) and (c). At the bottom (d), (e) and (f) are real space (zoomed) snapshots to support the cluster formation by fixing the activity  $\bar{v} = 300$  and  $t = 30$  for size ratios 6, 8 and 10 respectively. . . . . 40
- 2.4 (color online) Cluster size distribution (*CSD*) and real space snapshots of part of the system are plotted for different parameters. Different panels (a-d) are for fixed  $\bar{v} = 100$ , (e-h) are for  $\bar{v} = 300$  and (i-l) are for  $\bar{v} = 600$  and black, red and green colors are for size ratios 6, 8 and 10 respectively. *CSD* is shown in (a),(e) and (i) for  $\bar{v} = 100, 300$  and 600 respectively and for size ratio 6, 8 and 10. Symbols have the same meaning as in Fig. 2.3. . . . . 42
- 2.5 (color online) Mean mass of the largest cluster  $\langle m(t) \rangle$  is plotted for activities  $\bar{v} = 100, 300$  and 600. Left panel is for size ratio 10 and right panel is for size ratio 6. Black circles ( $\bullet$ ); red squares ( $\blacksquare$ ); green diamonds ( $\blacklozenge$ ) show the activities 100, 300 and 600 respectively. Straight line is the line with slope 1/3. (c),(d) and (e), (f) are the zoomed plot for intermediate ( $t \simeq 0.01 - 10$ ) and late time ( $t \simeq 10 - 100$ ) region for the same parameters. 43

3.1 (color online) Time evolution of the coarse grained local density,  $\phi_{loc}$  of ABPs is shown for  $\phi_a = 0.35$  for system with no obstacle, and obstacle with size ratio  $S = 2, 6$  and  $10$  from left (a) – (d) at fixed  $Pe = 100$ . The snapshots from bottom to top panel is in increasing time direction for times  $10^{-2}v_r^{-1}, 10^{-1}v_r^{-1}$  and  $v_r^{-1}$  respectively. Color bar shows the local density in area of  $12r_a \times 12r_a$ . . . . . 52

3.2 (color online) Time evolution of the coarse grained local density of ABPs is shown for  $\phi_a = 0.35$  for system with no obstacle  $e$  and  $f$ , and obstacle of size ratio  $10, g$  and  $h$  for  $Pe = 30$  and  $50$  respectively. The snapshots from bottom to top panel is for times  $10^{-2}v_r^{-1}, 10^{-1}v_r^{-1}$  and  $v_r^{-1}$  respectively. Color bar shows the local density in area of  $12r_a \times 12r_a$ . . . . . 53

3.3 (color online) The probability distribution function (*PDF*) of local density  $P(\phi_{loc})$ , are shown for three packing fractions  $\phi_a = 0.3, 0.35$  and  $0.4$  as a function of size ratio,  $S$  and Peclet number,  $Pe$ . (a) – (c) are for the system with no obstacle and (d) – (f) are when there is obstacle present in the system. . . . . 54

3.4 (color online) shows the phase coexistence curves for three  $\phi_a = 0.3, 0.35$  and  $0.4$  for system with no obstacle (black  $\circ$ ) and for system with obstacle of size  $10$  times larger than ABP (red  $\square$ ). The number on Y-axis shows the location of peaks of *PDF* of  $\phi_{loc}$ . X-axis is the Peclet number. . . . . 55

3.5 (color online) shows the phase diagram in the plane of  $\phi_a$  and  $Pe$  for the system with no obstacle and with obstacle of size  $6$  and  $10$  times larger than ABP. The range of  $\phi_a$  varies from  $0.1$  to  $0.4$  and for Peclet number it goes from  $30$  to  $150$ . Color bar shows the values of the differences in location of low and high density peaks. . . . . 56

- 
- 3.6 (color online) Real space snapshots of clean system of ABPs in steady state in plane of  $\phi_a$  and  $Pe$ .  $Pe$  varies from 30 – 150 (left to right) and  $\phi_a$  goes from 0.1 – 0.4 (bottom to top). . . . . 57
- 3.7 (color online) Real space snapshots of the system of ABPs in presence of single quenched obstacle in steady state in plane of  $\phi_a$  and  $Pe$ .  $Pe$  varies from 30 – 150 (left to right) and  $\phi_a$  goes from 0.1 – 0.4 (bottom to top) . 58
- 4.1 (color online) Time evolution of the system for different values of noise strengths  $\eta_a$ , keeping the activity,  $\lambda$  fixed to 1. (a),(b),(c) and (d) are for noise strengths 0 (AMB), 0.5, 0.8 and 1.0 respectively. The average order parameter value is  $\psi_0 = 0$ , corresponding to critical composition. The color bar shows the range of  $\psi$  value. . . . . 65
- 4.2 (color online) (a) and (b) show the dynamical scaling of correlation function for two different values of the noise strengths 0.5 and 1.0 respectively and compared them with the passive binary mixture. The dashed lines indicate different time for each noise strength. The value of the activity is 1 for all three cases. (c) shows the static scaling for different noise strengths, 0.5, 0.75 and 1.0 at  $t=50000$ . (d) and (e) show the dynamical scaling for structure factor for two different values of the noise strengths 0.5 and 1.0 respectively and compared them with the passive binary mixture. (f) shows the scaled static structure factor for same set of parameters. Black solid line represents the PMB. Magenta dashed line shows the line of slope -3 . 66

4.3 (color online) Log-Log plot of  $L(t)$  vs.  $t$  is shown in (a) for different noise strengths,  $\eta_a$  and compared with PMB and AMB. Black and red dashed lines are the lines of slope  $1/3$  and  $1/4$  that show the power law of growth as  $L(t) \sim t^{1/3}$  and  $L(t) \sim t^{1/4}$  respectively. (b) – (g) shows the effective exponent  $1/z$  vs.  $t$  for same set of parameters as in (a). In each frame, (b) – (g), the horizontal black and red dashed lines are drawn at  $1/z = 1/3$  and  $1/z = 1/4$  respectively. . . . . 68

4.4 (color online) Time evolution of the system is shown for different activities left to right panel and for fixed moderate strength of noise  $\eta_a = 0.5$ . (a) – (f) are for activities  $\lambda = 0$  (PMB), 1, 4, 5, 6 and 8 respectively. The average order parameter value is  $\psi_0 = 0$ , corresponding to critical composition. The color bar shows the range of  $\psi$  value. . . . . 69

4.5 (color online) (a) – (e) show the effective exponent for activity  $\lambda = 1, 4, 5, 6$  and 8 for fixed strength of noise 0.5. Black and red horizontal dashed lines are drawn at  $1/3$  and  $1/4$  respectively. Scaled Correlation function and structure factor are shown in (f) and g is for activity 8. Strength of noise is fixed to 0.5. All plots are shown for three different time regimes, starting from very early time. Black solid line is for PMB. Magenta dashed line is line of slope -3. . . . . 71

5.1 (color online) Time evolution of the system for different values of noise strengths  $\eta_a$ , keeping the activity,  $\lambda$  fixed to 0.5. (a),(b),(c) and (d) are for noise strengths 0.2, 0.5, 0.8 and 1.0 respectively. The average order parameter value is  $\psi_0 = 0.05$ , that corresponds to slightly off-critical composition. The color bar shows the range of  $\psi$  value. . . . . 78

- 5.2 (color online) shows the (a) static scaling for different strengths of noise  $\eta_a = 0.5, 0.75, 0.1$  at fixed time  $t = 50000$  for activity,  $\lambda = 0.5$  and (b) dynamic scaling for noise strength 1.0. Black solid line shows the correlation for PMB. . . . . 79
- 5.3 (color online) (a) Domain length is shown for noise strengths  $\eta_a = 0.5, 0.75$  and 1.0 is shown here. Black solid line shows the PMB. Black and red dashed lines are lines of slope  $1/3$  and  $1/4$  respectively. (b) shows the effective growth exponent for noise strengths  $\eta_a = 0.5, 0.65, 0.75, 0.85$  and 1.0. Black and red dashed horizontal lines are to show the value 0.33 and 0.25 respectively . . . . . 79
- 5.4 (color online) Time evolution of the system for different values of noise strengths  $\eta_a$ , keeping the activity,  $\lambda$  fixed to 0.5. (a), (b), (c) and (d) are for noise strengths 0.2, 0.5, 0.8 and 1.0 respectively. The average order parameter value is  $\psi_0 = -0.05$ . The color bar shows the range of  $\psi$  value. 81
- 5.5 (color online) Snapshots at late times are shown to compare the two cases for same and opposite sign of activity,  $\lambda$  and off-criticality  $\psi_0$ . We have taken  $\lambda = 0.5$  and  $\psi_0 = \pm 0.05$ . Left column is for  $\lambda \psi_0 > 0$  and right column is shown for  $\lambda \psi_0 < 0$  at time,  $t = 50000$ . (a),(b),(c) and (d) are for noise strengths 0.2, 0.5, 0.8 and 1.0 respectively. The color bar shows the range of  $\psi$  value. . . . . 82
- 5.6 (color online) (a) Characteristic length is shown for  $\psi_0 = -0.05$  for two different strengths of noise  $\eta_a = 0.5$  and 1.0. Activity,  $\lambda$  is fixed to 0.5. Green and blue dashed lines are lines of slopes  $1/3$  and  $1/4$  respectively. (b) Shows the effective growth exponent for  $\psi_0 = -0.05$  for noise strengths  $\eta_a = 0.5$  and 1.0. Activity,  $\lambda$  is fixed to 0.5. Black and red dashed lines show the value 0.25 and 0.33 respectively . . . . . 83

- 5.7 (color online) Time evolution of the system for different values of noise strengths  $\eta_a$ , keeping the activity,  $\lambda$  fixed to 0.5. (a),(b),(c) and (d) are for noise strengths 0.2, 0.5, 0.8 and 1.0 respectively. The average order parameter value is  $\psi_0 = 0.4$ . The color bar shows the range of  $\psi$  value. . . . . 84
- 5.8 (color online) Time evolution of the system for different values of noise strengths  $\eta_a$ , keeping the activity,  $\lambda$  fixed to 0.5. (a),(b),(c) and (d) are for noise strengths 0.2, 0.5, 0.8 and 1.0 respectively. The average order parameter value is  $\psi_0 = -0.4$ . The color bar shows the range of  $\psi$  value. . . . . 85
- 5.9 (color online) Characteristic length is shown for  $\psi_0 = 0.4$  for two different strengths of noise  $\eta_a = 0.5$  and 1.0. Activity,  $\lambda$  is fixed to 1. Black and red dashed lines are lines of slopes 1/4 and 1/3 respectively. . . . . 86
- 5.10 (color online) Shows the effective growth exponent for  $\psi_0 = 0.4$  for PMB (a) and for AMBN with noise strengths  $\eta_a = 0.5$  and 1.0, (b) and (c) respectively. Activity,  $\lambda$  is fixed to 1.0. Black and red dashed lines show the value 0.25 and 0.33 respectively. . . . . 86
- 6.1 (color online)(a)-(e) are plot of left hand size of equation 6.13 for different activities  $-3, -1, 0, 1$  and  $3$  respectively. Upper to bottom panels are for different times 80, 800 and 8000. . . . . 97
- 6.2 (color online) (f)-(j) and (k-o) are corresponding angle  $\theta(\mathbf{r}, t)$  and magnitude of fluid velocity  $|\mathbf{v}(\mathbf{r}, t)|$  respectively for the same set of activities as in (a-e) and at late time 8000. Color bar shows their respective values. In (f)-(j) the circles and squares represent the location of some of the vortices and antivortices with winding number  $k = +1$  and  $-1$  respectively. . . . . 98
- 6.3 (color online) Number of defects vs. time is plotted for three different activities  $-3, 0, 3$ . Black solid line of slope  $-1$  is drawn. . . . . 100

- 6.4 (color online) Two-point correlation function  $C(r,t)$  vs. distance  $r$  at different times for two different  $\xi = 3$  and  $-3$  for system size  $L = 1024$  in (a) and (b) respectively. The inset figures show the scaled two-point correlation  $C(r/L(t))$  vs. scaled distance  $r/L(t)$ . . . . . 101
- 6.5 (color online)(a) Shows the plot of characteristic lengths  $L(t)$  vs. time  $t$  on linear scale for positive  $\xi = 0, 1, 3, 4$  and  $5$  and for negative  $\xi = -3$  and  $-4$ . (b) shows the same plot on  $\log - \log$  scale. The two dashed lines are lines of slope  $1/2$ . (c) The plot of  $\frac{t}{(L(t))^2}$  vs.  $\ln(t)$  for the  $\xi$ 's as in (a) and (b). 101
- 6.6 (color online) Static scaled two-point correlation  $C(r/L(t))$  vs. scaled distance  $r/L(t)$  for  $\xi = 1, 2, 3$  and  $5$  in (a) and  $-1, -2$  and  $-3$  in (b) for system size  $L = 1024$  at fixed  $t = 1500$ . (c) Plot of  $(1 - C(r/L(t)))$  vs.  $r/L(t)$  on  $\log - \log$  scale for different  $\xi$ . The dashed line has slope  $1.6$  . . . 103