

**Phytochemistry, Medicinal Chemistry & *In-vitro*
Cytotoxicity of *Ipomoea nil* and
*Gloriosa superba***



Thesis submitted in partial fulfillment for the
award of degree

Doctor of Philosophy

By

Bharat Goel

PHARMACEUTICAL ENGINEERING & TECHNOLOGY
INDIAN INSTITUTE OF TECHNOLOGY
(BANARAS HINDU UNIVERSITY)
VARANASI - 221005
INDIA

Roll No. 18161004

2022

Shreyans K. Jain, PhD, Assistant Professor

CERTIFICATE

It is certified that the work contained in the thesis titled “**Phytochemistry, Medicinal Chemistry & In-vitro Cytotoxicity of Ipomoea nil and Gloriosa superba**” by **Mr. Bharat Goel** has been carried out under my supervision and that this work has not been submitted elsewhere for a degree.

It is further certified that the student has fulfilled all the requirements of Comprehensive Examination, Candidacy, and SOTA for the award of Ph.D. Degree.

Date: 27.6.2022

Place: IIT (BHU), Varanasi



Dr. Shreyans K. Jain
(Supervisor)


डा० श्रेयांस कु० जैन / Dr. Shreyans K. Jain
सहायक आचार्य / Assistant Professor
भेषजकीय अभियांत्रिकी एवं प्रौद्योगिकी विभाग /
Department of Pharmaceutical Engineering & Technology
भारतीय प्रौद्योगिकी संस्थान / Indian Institute of Technology
(काशी हिन्दू विश्वविद्यालय) / (Banaras Hindu University)
वाराणसी-221005 / Varanasi-221005

DECLARATION BY THE CANDIDATE

I, **Bharat Goel** certify that the work embodied in this PhD thesis is my own bona fide work and carried out by me under the supervision of **Shreyans K. Jain**, from July 2018 to June 2022, at the Department of Pharmaceutical Engineering and Technology, Indian Institute of Technology (Banaras Hindu University), Varanasi. The matter embodied in this thesis has not been submitted for the award of any other degree/diploma. I declare that I have faithfully acknowledged and given credits to the research workers wherever their works have been cited in my work in this thesis. I further declare that I have not wilfully copied any other's work, paragraphs, text, data, results, *etc.*, reported in journals, books, magazines, dissertations, theses, *etc.*, or available at websites and have not included them in this PhD thesis and have not cited as my own work.


Date: 27.06.2022

Place: IIT (BHU), Varanasi


Signature of the Student
(Bharat Goel)

CERTIFICATE BY THE SUPERVISOR

It is certified that the above statement made by the student is correct to the best of my knowledge.


Dr. Shreyans K. Jain

(Supervisor)

डा० श्रेयांस कु० जैन / Dr. Shreyans K. Jain
सहायक आचार्य / Assistant Professor
भैषजकीय अभियांत्रिकी एवं प्रौद्योगिकी विभाग /
Department of Pharmaceutical Engineering & Technology
भारतीय प्रौद्योगिकी संस्थान / Indian Institute of Technology
(काशी हिन्दू विश्वविद्यालय) / (Banaras Hindu University)
वाराणसी-221005 / Varanasi-221005


Prof. S. Hemalatha

(Head of the Department)

विभागाध्यक्ष / Head

भैषजकीय अभियांत्रिकी एवं प्रौद्योगिकी विभाग /
Department of Pharmaceutical Engineering & Technology
भारतीय प्रौद्योगिकी संस्थान / INDIAN INSTITUTE OF TECHNOLOGY
(बनारस हिन्दू विश्वविद्यालय) / (BANARAS HINDU UNIVERSITY)
वाराणसी-221005 / Varanasi-221005

COPYRIGHT TRANSFER CERTIFICATE

Title of the Thesis: Phytochemistry, Medicinal Chemistry & *In-vitro*
Cytotoxicity of *Ipomoea nil* and *Gloriosa superba*

Name of the Student: Bharat Goel

Copyright Transfer

The undersigned hereby assigns to the Indian Institute of Technology (Banaras Hindu University) Varanasi all rights under copyright that may exist in and for the above thesis submitted for the award of the "**Doctor of Philosophy**"

Date: 27.06.2022

Place: IIT (BHU), Varanasi



Signature of the Student
(Bharat Goel)

Note: However, the author may reproduce or authorize others to reproduce material extracted verbatim from the thesis or derivative of the thesis for author's personal use provided that the source and the Institute's copyright notice are indicated.

Acknowledgments

At the foremost, special reverence to ***Baba Vishwanath*** and ***the Holy river Ganga***. This work would not have been possible without their blessings. At this moment of accomplishment, I would like to pay the homage to the founder of Banaras Hindu University, ***Pandit Madan Mohan Malviya ji***, who made this glorious temple to realize spiritual, technical and scientific knowledge about this vast existing universe.

I embrace the opportunity to express my deep sense of gratitude to my supervisor ***Dr. Shreyans Kumar Jain***, for his constant guidance, valuable suggestions and kind encouragement during my association with his research group. His encouragement, constant support, intellectual stimulation, perceptive guidance, immensely valuable ideas, and suggestions from the initial to the final level enabled me to develop an understanding of the subject. His scholarly suggestions, prudent admonitions, immense interest, constant help, and affectionate behavior have been a source of inspiration for me. His suggestions will remain with me as an inexhaustible source of scientific learning throughout my life.

I am also grateful to ***Prof. Brahmeshwar Mishra, Prof. Sushil Kumar Singh, Prof. Sanjay Singh, & Prof. Sushant Kumar Shrivastava***, Former Heads of department, for their constant support, which led to the successful completion of this work.

I also take this opportunity to express my deepest gratitude to ***Prof. Siva Hemalatha***, Head, Department of Pharmaceutical Engineering and Technology, IIT (BHU), Varanasi, for allowing me to use the departmental facilities to carry out my research work.

I owe my gratitude to all my respected Research Progress and Evaluation Committee (RPEC) members ***Dr. Gyan Prakash Modi*** and ***Dr. Marshal*** for providing valuable suggestions and kind help in my research work. I express my thanks to ***Dr. Santosh Kumar Guru***, NIPER Hyderabad, for helping me in carrying out the *in-vitro* cytotoxicity studies. I am thankful to ***Dr. Arindam Indra***, Dept. Of Chemistry, IIT (BHU), for helping me in new synthetic methodology development. I

would also like to thank **Prof. Sairam Krishnamurthy, Dr. Ashok Kumar, Dr. Senthil Raja A., Dr. Alakh Niranjana Sahu, Dr. Ruchi Chawla, Dr. M. S. Muthu, Dr. Prasanta Kumar Nayak, Dr. Sunil Kumar Mishra, Dr. Vinod Tiwari, Dr. Ashish Kumar Agrawal, Dr. Rajnish, Dr. Deepak Kumar, Dr. Dinesh Kumar, Dr. Jairam Meena,** and **Dr. Arun Khattri** for their support and encouragement.

I am thankful to **Prof. N. K. Dubey** (BHU), and **Dr. Bikarma Singh** (CSIR-NBRI) for the authentication of plant materials. I am also thankful to **Prof. Amrita Cheema**, Georgetown University, USA for the LCMS analysis. I acknowledge **Mr. Amit Gautam**, CSIR-IIIM, Jammu for the HRMS analysis.

I pay my sincere thanks to **all the non-teaching staff** of the department for their timely assistance and co-operation during my Ph.D. tenure.

I am indeed thankful to my labmates, batchmates, seniors, and juniors **Ms. Nancy Tripathi, Ms. Nivedita Bhardwaj, Dr. Yashpal Singh, Mr. Himashu Rai, Mr. Bharat Sahu, Mr. Gourav Singh, Mr. Gauri Shankar, Ms. Shreyasi Majumdar, Ms. Vishnu Priya, Mr. Vikas, Mr. Sanjay Kumar, Mr. Anurag Saini, Mr. Rakesh Sahu,** and **Mr. Sunil Kumar** for their support and co-operation.

I would particularly like to acknowledge my friends **Dr. Krishna Kumar Patel, Dr. Gopichand Gutti, Mr. Tarkeshwar Dubey** for their support. They have always been there to motivate me during the hour of hardship and helped me unconditionally.

A special thanks to my family. Words can not express how grateful I am to **my family members** for all the sacrifices that they have made on my behalf and continuous support they provided to me. They have been a constant source of love, moral support, and strength all these years. I owe this achievement to them.

I gratefully acknowledge the MHRD, Govt. of India for providing me the necessary funding and fellowship to pursue research work.

Date: 27.06.2022

Place: IIT (BHU), Varanasi



(Bharat Goel)

DEDICATED TO MY BELOVED FAMILY

TABLE OF CONTENTS

CONTENTS	Page No.
Certificates	ii
Acknowledgments	v
Table of contents	viii
List of figures	xi
List of tables	xiv
List of abbreviations	xv
Preface	xvii
Chapter 1. Introduction	
1. Introduction	1
1.1. Natural products as potential sources of drugs	1
1.2. Semi-synthesis as an important tool in natural products-based drug discovery	3
1.3. Total synthesis	5
1.4. Diverted total synthesis (DTS)	7
1.5. Function-oriented synthesis (FOS)	7
1.6. Biology-oriented synthesis (BIOS)	8
1.7. Biosynthesis inspired synthesis	8
1.8. Aims and objectives	9
1.9. Plan of work	10
1.9.1. Isolation, total synthesis, and cytotoxic activity of compounds of <i>Ipomoea nil</i> seeds	10
1.9.2. Isolation, semi-synthetic modification, and cytotoxic evaluation of compounds of <i>Gloriosa superba</i> roots	11
Chapter 2. Isolation, total synthesis, and cytotoxic activity of compounds of <i>Ipomoea nil</i> seeds	
2. Isolation, total synthesis, and cytotoxic activity of compounds of <i>Ipomoea nil</i> seeds	13
2.1. Literature review	13
2.2. Results and discussion	16
2.2.1. Phytochemical investigation of <i>I. nil</i> seeds	16
2.2.2. Structure elucidation of new compound 2.11	18
2.2.3. Ipomone (2.11): An unusual bicyclo[3.2.1]octanone, perhaps a process generated product	22
2.2.4. <i>In-vitro</i> cytotoxic activity of ipomone (2.11)	23
2.2.4.1. Cytotoxicity of ipomone (2.11) against cancer cells	23
2.2.4.2. Ipomone (2.11) causes A549 cell death by apoptosis	24
2.2.4.3. Ipomone (2.11) induces autophagy	25
2.2.4.4. Immunoblot analysis	25
2.2.4.5. Molecular docking studies with caspase-3, p110 α , and PARP-1 proteins	27

2.2.5. Total synthesis of ipomone (2.11)	28
2.2.5.1. Mechanism of formation of ipomone from gibberellic acid	30
2.3. Experimental section	35
2.3.1. General experimental procedures	35
2.3.2. Plant material	36
2.3.3. Extraction and isolation	36
2.3.4. Cell line and cell culture	38
2.3.5. Cell viability assay	38
2.3.6. Fluorescence microscopy	39
2.3.7. Acridine orange staining	39
2.3.8. Preparation of whole-cell lysates for immunoblotting	39
2.3.9. Molecular docking	40
2.3.10. Procedure for the synthesis of ipomone (2.11)	41
2.3.11. Procedure for the synthesis of compounds 2.19 and 2.20	41
Chapter 3. Isolation, semi-synthetic modification, and cytotoxic evaluation of compounds of <i>Gloriosa superba</i> roots	
3. Isolation, semi-synthetic modification, and cytotoxic evaluation of compounds of <i>Gloriosa superba</i> roots	43
3.1. Literature review	43
3.1.1. SAR of colchinoids	45
3.2. Results and discussion	48
3.2.1. Phytochemistry of <i>Gloriosa superba</i> roots	48
3.2.2. Structure elucidation of new compound (3.19)	55
3.2.3. Biological activity of gloriosine	58
3.2.3.1. In-vitro cytotoxicity screening of gloriosine	58
3.2.3.2. Nuclear staining and fluorescence microscopy	59
3.2.3.3. Scratch assay for cell migration study	61
3.2.4. <i>In-silico</i> investigation of binding modes of gloriosine against tubulin protein	62
3.2.4.1. Molecular docking studies of gloriosine with tubulin protein	62
3.2.4.2. Molecular dynamic simulation of gloriosine with tubulin protein	63
3.2.5. Synthetic modifications of gloriosine	65
3.2.5.1. Series 1: Synthesis of C-10 amine derivatives of gloriosine	66
3.2.5.2. Series 2: Synthesis of C-10 amide derivatives of gloriosine	68
3.3. Experimental section	75
3.3.1. General experimental procedures	75
3.3.2. Plant material	76
3.3.3. Extraction and isolation	76
3.3.4. Synthetic modifications on gloriosine and new synthetic	81

method	
3.3.4.1. Synthesis of C-10 amino derivatives of gloriosine	81
3.3.4.2. Synthesis of amides by new synthetic methodology	87
3.3.4.3. Synthesis of C-10 amide derivatives of gloriosine	92
3.3.5. <i>In-vitro</i> cytotoxicity studies	93
3.3.5.1. General procedure for cytotoxicity (MTT) assay	94
3.3.5.2. Nuclear staining and fluorescence microscopy	94
3.3.5.3. Scratch assay for migration studies	95
3.3.6. Molecular docking studies	95
3.3.7. Molecular dynamics simulation of gloriosine-tubulin complex	96
Chapter 4. Conclusion and future perspectives	98
References	101
Appendix	118
List of publications	120

LIST OF FIGURES

	Figure Captions	Page No.
Figure 1.1	Chemical structures of plant- and microbial-derived natural products used as drugs	2
Figure 1.2	Typical examples of semisynthetic anticancer drugs	4
Figure 1.3	Examples of total synthesis	6
Figure 1.4	Example of diverted total synthesis (DTS)	7
Figure 1.5	Example of function-oriented synthesis (FOS)	8
Figure 1.6	Biosynthesis inspired synthesis	9
Figure 1.7	Flow chart showing plan of work of <i>Ipomoea nil</i>	11
Figure 1.8	Flow chart showing plan of work of <i>Gloriosa superba</i>	12
Figure 2.1	Chemical structures of compounds reported from <i>I. nil</i> (2.1-2.9) and semi-synthetic derivative GA-13315 (2.10)	14
Figure 2.2	Schematic diagram of isolation of compounds 2.11-2.18 from <i>I. nil</i> seeds	17
Figure 2.3	Chemical structures of compounds isolated from <i>I. nil</i> seeds	17
Figure 2.4	¹ H NMR spectrum of compound 2.11 in CDCl ₃ (500 MHz)	18
Figure 2.5	¹³ C NMR and DEPT-135 spectra of compound 2.11 in CDCl ₃ (125 MHz)	19
Figure 2.6	(a) ECD spectrum and (b) key NOESY correlations of compound 2.11	21
Figure 2.7	A plausible mechanism of 1,2 alkyl shift; a sort of pinacol-pinacolone rearrangement to produce 2.11	23
Figure 2.8	Dose and time dependent cytotoxicity of ipomone (2.11) against A549 cells	24
Figure 2.9	Fluorescence micrographs (DAPI staining) of A549 cells untreated and treated with ipomone for 24 hours	25
Figure 2.10	Morphology of untreated A549 cells (control) and those treated with 10, 30, and 50 μM of ipomone (2.11)	26
Figure 2.11	Immunoblot analysis using antibodies against the p110α, PARP-1, caspase-3, and LC 3B proteins in A549 cells after treatment with ipomone (2.11) (0-50 μM) for 24 hours	26
Figure 2.12	The protein-ligand interaction diagrams of ipomone with (A) caspase-3 (PDB id. 3KJF), (B) p110α (PDB id. 5DXT), and (C) PARP (PDB id. 6I8T)	27
Figure 2.13	Chemical reaction of gibberellic acid (2.1) with iodine to synthesize ipomone (2.11)	29
Figure 2.14	(a) ¹ H NMR spectra of ipomone and compound 2.19 in CDCl ₃ (500 MHz); (b) ¹³ C spectra of ipomone and	31

	compound 2.19 in CDCl ₃ (125 MHz); (c) DEPT-135 NMR spectrum of compound 2.19 in CDCl ₃ (125 MHz)	
Figure 2.15	A plausible mechanism involved in the synthesis of ipomone (2.11). (E ⁺ for electrophile)	33
Figure 2.16	¹³ C NMR spectra of intermediates (a) 2.11a and (b) 2.11b in CDCl ₃ (125 MHz)	33
Figure 2.17	Plausible mechanism for the aromatization of ring A	34
Figure 3.1	Reported compounds from <i>Gloriosa superba</i> roots	44
Figure 3.2	Colchicine at the colchicine-binding site located between α - and β -subunits of the tubulin protein	45
Figure 3.3	Structure-Activity Relationship (SAR) studies of colchinoids	47
Figure 3.4	Isolated compounds from <i>G. superba</i> roots	49
Figure 3.5	NOESY NMR spectra of β -lumicolchicine (3.10) and γ -lumicolchicine (3.18)	51
Figure 3.6	¹ H NMR spectra of (a) colchicine (3.2), and (b) gloriosine (3.3) in CDCl ₃ (500 MHz)	52
Figure 3.7	¹³ C NMR spectra of (a) colchicine (3.2), and (b) gloriosine (3.3) in CDCl ₃ (125 MHz)	53
Figure 3.8	DEPT-135 and HSQC NMR spectra of gloriosine (3.3)	54
Figure 3.9	Chemical structure of new compound, gloriogerine (3.19)	55
Figure 3.10	¹ H, ¹³ C, and DEPT-135 NMR spectra of new compound 3.19 in CDCl ₃	57
Figure 3.11	HSQC and HMBC spectra of new compound 3.19	58
Figure 3.12	Nuclear staining after treatment with different concentrations of gloriosine (50, 100, and 500 nM) and colchicine (500 nM) on A549 cells for 24 h revealed the formation of apoptotic bodies in a dose-dependent manner.	60
Figure 3.13	Effect of gloriosine (3.3) and colchicine (3.2) on <i>in-vitro</i> cell migration in A549 cells after 24 h	61
Figure 3.14	The protein-ligand interaction diagram of compounds A) Gloriosine and B) Colchicine against tubulin protein (PDB ID: 4O2B)	62
Figure 3.15	MD simulation of gloriosine complexed with tubulin	64
Figure 3.16	Different series of gloriosine derivatives	65
Figure 3.17	Synthesis of 10-amine derivatives of gloriosine (3.3)	66
Figure 3.18	¹ H NMR spectrum of compound 3.20h in CDCl ₃ (500 MHz)	67
Figure 3.19	¹³ C NMR spectrum of compound 3.20h in CDCl ₃ (125 MHz)	67

Figure 3.20	HMBC NMR spectrum of compound 3.20h	68
Figure 3.21	Synthesis of C-10 benzamide derivative of colchicine <i>via</i> EDC coupling reaction	69
Figure 3.22	LCMS analysis of EDC coupling reaction mass	69
Figure 3.23	Reaction of C-10 benzamide derivative of colchicine <i>via</i> benzoylation reaction	70
Figure 3.24	Synthesis of C-10 amide derivative of colchicine <i>via</i> oxidative amidation reaction	73
Figure 3.25	LCMS analysis of oxidative amidation reaction mass	74
Figure 3.26	Synthesis of C-10 amide derivatives of gloriosine	75
Figure 3.27	¹³ C NMR spectrum of compound 3.26a in CDCl ₃ (125 MHz)	75

LIST OF TABLES

	Table Legends	Page No.
Table 2.1	^1H and ^{13}C NMR spectroscopic data for new compound (2.11) in CDCl_3	21
Table 2.2	Cytotoxicity of ipomone (2.11) against cell lines tested	24
Table 2.3	Treatment of gibberellic acid (2.1) under different reaction conditions	29
Table 2.4	^1H and ^{13}C NMR spectroscopic data for compound 2.19 in CDCl_3	32
Table 3.1	NMR spectroscopic data for compound 3.19 in CDCl_3	56
Table 3.2	IC_{50} values (nM) of gloriosine (3.3) and colchicine (3.2) in various human cancer cell lines and normal cell line after 48 h	59
Table 3.3	Optimization of reaction conditions	71
Table 3.4	Scope of the amidation reaction	72

LIST OF ABBREVIATIONS

2D NMR	Two dimensional NMR
Å	Angstrom
ACN	Acetonitrile
ADMET	Absorption, distribution, metabolism, excretion, and toxicity
AO	Acridine orange
BIOS	Biology-oriented synthesis
Calcd	Calculated
CBS	Colchicine binding site
CC	Column chromatography
CD ₃ OD	Methanol-d ₄
CDCl ₃	Deuterated chloroform
CHCl ₃	Chloroform
COSY	Correlation spectroscopy
d, dd	Double, Double doublet
DAPI	(4',6-Diamidino-2-phenylindole)
DCC	N,N'-Dicyclohexylcarbodiimide
DCM	Dichloromethane
DEPT	Distortionless enhancement by polarization transfer
DMSO	Dimethylsulphoxide
DNA	Deoxyribonucleic acid
DNP	Dictionary of natural products
DTS	Diverted total synthesis
ECD	Electronic circular dichroism
EDC	1-Ethyl-3-(3-dimethylaminopropyl)carbodiimide
Eq. or Equiv.	Equivalent
ESI	Electron spray ionization
EtOAc	Ethyl acetate
FDA	Food and drug administration
FOS	Function-oriented synthesis
g or gm	Gram
GA	Gibberellic acid
h or hr	Hour
H-bond	Hydrogen bond
HCl	Hydrochloric acid
HMBC	Heteronuclear multiple bond correlation
HOBt	Hydroxybenzotriazole
HRMS	High resolution mass spectrometer
HSQC	Heteronuclear single quantum coherence spectroscopy
IR	Infrared spectroscopy
J	Coupling constant (units of Hz)
kg	Kilogram

LC	Liquid chromatography
LCMS	Liquid chromatography–mass spectrometry
m	Metre
MD	Molecular dynamic
MeOH	Methanol
MHz	Megahertz
min	Minutes
mL or ml	Mililitre
mmol	Milimole
mp or m.p.	Melting point
mPT	Mitochondrial permeability transition
MS	Mass spectrometer
MTT	3-(4,5-Dimethylthiazol-2-yl)-2,5-diphenyltetrazolium bromide
NCI	National Cancer Institute
NF- κ B	Nuclear factor kappa-light-chain-enhancer of activated B cells
nM	Nanomole
NMR	Nuclear magnetic resonance
NOESY	Nuclear overhauser effect spectroscopy
NP	Natural product
No.	Number
PBA	Prussian blue analogs
PBS	Phosphate-buffered saline
PCF	Plant cell fermentation
PDB	Protein Data Bank
P-gp	P-glycoprotein
PI	Propidium iodide
Rf	Retention factor
RMSD	Root-mean-square deviation
RMSF	Root mean square fluctuation
SAR	Structure-activity relationship
SD	Standard deviation
TBHP	tert-Butyl hydroperoxide
TLC	Thin-layer chromatography
TMS	Trimethylsilane
v/v	Volume by volume
w/w	Weight by weight
μ l or μ L	Microlitre
μ M	Micromole

PREFACE

Natural products and their analogs comprise a major part of the approved drugs, mainly anticancer and anti-infectives, also in other therapeutic areas like cardiovascular diseases etc. Almost half of the approved drugs comprise either natural products, their semi-synthetic derivatives, or inspired from natural products. Natural products have a wide range of structural complexity and diversity. Despite the large number of successful drugs from natural sources, there is a declining trend in the number of NP-derived new drugs, raising renewed interest in the natural product-based drug discovery.

The present thesis work was aimed at the discovery of new anticancer lead molecules, derived from the natural products. In the present work, two plants were selected: *Ipomoea nil* (Convolvulaceae), and *Gloriosa superba* (Colchicaceae), based on degree of exploration and reported cytotoxic potential. These two plants were investigated for their phytochemistry and semi-synthetic derivatization of the major isolates.

This thesis consists of five chapters: (1) Introduction, (2) Isolation, synthesis, and *in-vitro* cytotoxic activity of compounds of *Ipomoea nil* seeds, (3) Isolation, semi-synthetic modification, and *in-vitro* cytotoxic evaluation of compounds of *Gloriosa superba* roots, (4) Conclusions and prospects, and References.

In the first chapter, the role of natural products in drug discovery is discussed, along with the examples and disadvantages of natural products. Semi-synthesis as an important tool to optimize the natural products in terms of biological activity as well as pharmacokinetic properties is discussed. The aims and objectives of the thesis and the plan of work is also discussed.

Chapter 2 describes the first objective: Isolation, total synthesis, and *in-vitro* cytotoxic activity of compounds of *Ipomoea nil* seeds. The phytochemical investigation

of *Ipomoea nil* seeds (commonly known as morning glory) is discussed. Eight compounds, including a novel diterpenoid containing bicyclo[3.2.1]octanone nucleus, were isolated from the acidified hydroalcoholic extract of *Ipomoea nil* seeds. All the isolated compounds were characterised on the basis of detailed NMR and mass spectral techniques. The novel compound, ipomone, showed structural similarity to the gibberellins that has been reported from the *I. nil* seeds. Ipomone was screened for cytotoxic activity in a panel of 12 cancer cell lines and found to have moderate activity ($IC_{50} = 34 - 86 \mu M$). The mechanistic investigation suggested that ipomone produced the cytotoxic activity mediated by apoptosis and autophagy.

We hypothesized, that ipomone is an acid-catalysed pinacol pinacolone type 1,2 alkyl rearrangement product of gibberellin, as it was obtained under acidic extraction conditions. Inspired by the fact that many gibberellins have already been reported from *I. nil* seeds, an efficient one-step method for the synthesis of ipomone was developed using commercially available gibberellic acid as a starting material and molecular iodine.

In the third chapter, the second objective i.e., isolation, semi-synthetic modification, and *in-vitro* cytotoxic evaluation of compounds of *Gloriosa superba* roots is discussed. The roots of *Gloriosa superba* were investigated for phytochemistry and 17 compounds were isolated including colchicine, gloriosine, and a new compound. All the known compounds were characterized by extensive NMR studies and comparing with reported data. The new compound was elucidated based on comprehensive spectroscopic analysis, including 1D, 2D NMR and HRMS.

Gloriosine has never been explored for its pharmacology, so we carried out its cytotoxic screening and found that gloriosine exhibited significant cytotoxicity ($IC_{50} = 32$ to 100 nM) that was comparable with colchicine. Since C-10 amino derivatives of

colchicine produced potent compounds, twenty semi-synthetic C-10 amino and amide derivatives of gloriosine were synthesized. Usual methods like benzoylation of amines, EDC coupling reactions did not provided the amide derivatives in an isolable yield. So, a new synthetic method for the synthesis of amides directly from aldehydes and amines using a nickel catalyst was developed. This new amidation method was found suitable for the synthesis of desired amide derivatives of gloriosine in isolable good yields.

Chapter 4 describes the conclusion and future prospects of the thesis. A new natural product, ipomone, was isolated from the seeds of *Ipomoea nil*. The total synthesis of ipomone led to the discovery of an intermediate (iodomethyl derivative of ipomone) that could be used as starting material to synthesise additional analogs for the future prospects. An anticancer lead, gloriosine, having *in-vitro* cytotoxic activity in nanomolar range was isolated from the roots of *Gloriosa superba*. Further, C-10 amino and amide derivatives of gloriosine were synthesized. The study suggested that gloriosine might serve as an interesting lead for further *in-vivo* and preclinical studies and future anticancer therapeutics.