

List of figures

1.1	The schematic represents the core, radiative, and convection zone in the solar interior, while the photosphere, chromosphere, and corona in its exterior. The thermonuclear reactions and transport of the energy by the mean of radiative and convective motions are illustrated in the interior. General perception of the entire electromagnetic radiations (e.g., γ -rays, X-rays, visible, UV, and radio emissions) are portrayed from different parts of the solar atmosphere. The different magnetic regions (e.g., active regions with the sunspot groups, coronal hole, quiet-Sun) are displayed. The prominence, coronal loops, and magnetic fluxtubes are also shown in the localized solar atmosphere. (Courtesy: www.solarviews.org)	3
1.2	Variation of temperature and density with height from the Sun's core to its surface. This image is adopted from Strong et al. (2012). (Courtesy: K. Strong)	5
1.3	Helioseismic measurements determine the internal rotation rate depicting that the longitudinal differential rotation is observed at the solar surface extending through the convection zone. The surface and tachocline regions possess strong radial shear. The blue, green, and red colours represent progressively faster rotations, collectively indicating a solar differential rotation. (Courtesy: NCAR/UCAR, HAO)	8

-
- 1.4 The full-disk visible surface of the Sun's photosphere is shown in the left-panel of the image. The zoomed region in right-panel displays the fine details of an active region (AR) sunspot, and the quiet-Sun (QS) granules in its surrounding. (Courtesy: SST; Luc Rouppe van der Voort and Shahin Jafarzadeh, University of Oslo, Norway) 11
- 1.5 Variation of temperature and density with height in the solar atmosphere as per VAL (Vernazza-Avrett-Loeset) solar atmospheric model. (Courtesy: Eugene Avrett, Smithsonian Astrophysical Observatory) 14
- 1.6 The full-disk image of the chromosphere is shown in $H\alpha$ 6563 Å . Different chromospheric features like sunspots, active regions, filaments, networks, plage regions appear. Right panel: Calcium II K 3933 Å line full disk image of the Sun's chromosphere. Several chromospheric features like sunspots, plage, tiny intra-networks, chromospheric networks, etc. evident in this image. [Courtesy: William Lee (<https://www.cloudynights.com/gallery/member/25174-bill-lee/>)] 15
- 1.7 The composite view of the EUV inner corona as observed by Solar Dynamics Observatory (SDO)/Atmospheric Imaging Assembly (AIA) 171Å filter and Solar and Heliospheric Observatory (SoHO)/The Large Angle and Spectrometric Coronagraph (LASCO-C2) showing the extended corona upto 6 $R_{\odot}$. Observational data is recorded on 21 April 2015. (Courtesy: SDO/AIA and SoHO/LASCO) 16
- 1.8 Left panel: Large-scale filaments as observed by the $H\alpha$ filter on the solar disk. Right panel: An eruptive prominence in He II 304 Å with an image of the Earth added for size comparison. (Courtesy: Solar and Heliospheric Observatory *SoHO*/EUV Imaging Telescope *EIT*; Big Bear Solar Observatory) 19

-
- 1.9 Images of the different types of solar prominences based on their location as observed by the SDO/AIA 304 Å filter. (Courtesy: Observational data from SDO/AIA) 21
- 1.10 A chromospheric filament is observed by the high resolution H α image on 30th October 2002. Dextral, sinistral, and prominence barbs patterns appear in the prominence. (Courtesy: USAF, ISOON; D. Neidig) 25
- 1.11 Rayleigh-Taylor unstable plumes in a solar prominence in the linear stage with the required conditions, e.g., self-similarity in same family of plumes, similarity of the wavelength/height ratio in different families. Schematic shows the method to estimate the observational growth rate of the instability. (Courtesy: Ryutova et al., 2010) 28
- 1.12 Upper panel: A basic schematic that shows the 2D spontaneous magnetic reconnection and release of energy. Different colors show the different connectivity of the field lines and arrows indicate their directions. Bottom panel: 2D schematic for the forced magnetic reconnection. Some external perturbation forced the coronal field lines to reconnect. The spring like structure indicates the external perturbations and star indicates the reconnection region where the energy release takes place. 32
- 1.13 CME image is associated with an eruption of the active region prominence on 07 June 2011. The MRT unstable prominence forms the core of CME that propagates in the outer corona. (Courtesy: STEREO/EUVI, COR-I) . 34
- 1.14 Three consecutive CMEs are erupted on 22 October 2011, which collectively interact with the Earth's atmosphere on 24-25 October 2011. They exhibit an intense geomagnetic storm (Dst index -146 nT) on 25 October 2011. (Courtesy: STEREO/COR-II, HI-I) 37

-
- 2.1 The Extreme Ultraviolet Imager (EUI) onboard Solar Orbiter will open a new window to understand many of the spectacular features that have not been observed before by other space satellites. The first image obtained from the Solar orbiter that shows the omnipresence nature of campfires on the Sun, signify the importance of EUV observations. (Courtesy: Solar Orbiter/EUI Team in ESA & NASA; CSL, IAS, MPS, PMOD/WRC, ROB, UCL/MSSL) 45
- 2.2 An image of SDO spacecraft. The SDO spacecraft consists of three instruments, the Atmospheric Imaging Assembly (AIA), the Heliospheric and Magnetic Imager, and the Extreme Ultraviolet Variability Experiment (EVE) that simultaneously probe the different layers of the solar atmosphere. (Courtesy: NASA) 49
- 2.3 Multiwavelength images of AIA instrument, which observe the different region of the Sun. (Courtesy: <https://solarsystem.nasa.gov/>) 50
- 2.4 Location of spacecraft STEREO-A and STEREO-B drifting away from each other to observe the different views of the Sun. In the combination of STEREO and SDO, these two spacecraft observe the entire Sun for several years. (Courtesy: NASA/GSFC) 53
- 2.5 Left panel: Image of different instruments onboard the STEREO spacecraft. Right panel: Image of the SECHHI instrument onboard the STEREO and its main components. (Courtesy: NRL/NASA) 54
- 2.6 Internal view of the EUVI telescope onboard STEREO. (Courtesy: Wuelser et al. (2004)). 55

-
- 2.7 The left panel shows the normal SDO/AIA 171 Å image. The right panel shows the Multi-scale Gaussian normalization processed SDO/AIA 171 Å image. The processed image shows the more fine-scale structures in the solar corona. 60
- 2.8 The left panel shows the STEREO-B image of an eruptive prominence. The right panel shows the STEREO-A image. The + signs in both images are lying on the same epipolar plane and provides the exact location of the feature. 62
- 2.9 Images of six EUV filters of SDO/AIA. These six optically thin EUV filters used to understand the thermal structure of the solar atmosphere. . . 65
- 2.10 Temperature response functions for the six EUV channels of the SDO/AIA that are optically thin and dominated by iron emission lines. (Courtesy: (Lemen et al., 2012)). 66
- 3.1 Upper panel: the full FOV of the loop-like prominence observed by SDO/AIA 304 Å on 2014 November 18th after applying multi-Gaussian normalized filter. The sub-region (white box) consists of an MRT-unstable plume. Middle panel: a dark cavity has been evident within the prominence. The spatio-temporal evolution of MRT-unstable plume (P1) has been observed. Lower panel: the second plume, “P2,” is tracked at different heights and times to estimate the observational growth rate of the MRT instability. The temporal image data in the middle and lower panels show the increments in time from right to left. This is the partial FOV of an eruptive prominence where all plasma processes related to MRT instability are evolved. 77

- 3.2 Upper panel: the sub-region of the eruptive prominence using a multi-Gaussian normalized filter on an AIA 304 Å image. A vertical line is drawn to indicate the initial location of the first plume (P1). The separation between the two consecutive plumes is the characteristic wavelength of an MRT instability. Lower panel: DEM of the same sub-region to observe the thermal structure of this MRT-unstable prominence. 81
- 3.3 Top panel: the spatiotemporal variation of the full FOV of the overlying prominence, which collapses by plasma downfall. The white boxes "a" and "b" indicate the region where KH unstable vortices and rolled plasma structures are formed. The evolutions of these vortices and rolled plasma structures are shown in the lower panels. 83
- 3.4 Top panel: Full FOV of multi-Gaussian normalized and fitted SDO/AIA 304 Å filter, which consists of Rayleigh-Taylor unstable prominence thread. Middle panel: The sequence of images of the sub-region in SDO/AIA 304 filter showing the temporal evolution of the plasma thread. Lower panel: The V-shaped dip as evident in the lower panel in the SDO/AIA 171 Å filter. 89
- 3.5 The sequence of the images of SDO/AIA 171 Å shows the boundary of the prominence-cavity, which became unstable. The middle panel shows that two bigger plasma bubbles are observed. Later they may be responsible for the shear motion at the interface. The shear motion initiates the Kelvin-Helmholtz unstable vortex formation and Rayleigh-Taylor unstable bubbles are also formed at the boundary. 90

- 3.6 The region of interest (ROI; white box in Figure 3.5) is displayed in the SDO/AIA (171+304)Å composite images. The composite images observed the characteristic of the bubble-prominence boundary. Shear motion, plasma downfall, MRT unstable bubbles associated with the Rayleigh-Taylor unstable thread are indicated by arrows at the interface. The complete evolution of shearing, bubbles formation, MRT unstable bubbles, and collapse of the bubbles at the prominence-cavity interface is also indicated. 92
- 3.7 The region of interest (ROI; white box in Figure 3.5) is displayed using SDO/AIA 304 Å images. MRT unstable bubbles, shear motion, and KH unstable vortex have been indicated by arrows at different time-epochs of the prominence-cavity boundary. 95
- 3.8 MRT unstable prominence bubbles are tracked at different height at the prominence-cavity interface. The observed bubbles propagate with different velocity and trying to merge within themselves. The collapse phase of MRT unstable bubbles has been observed in the lower panel. 98
- 4.1 The normalized multi-gaussian fitted SDO/AIA 171 Å images showing the evolution of solar tornadoes-like motions related to the polar prominence during 9–14 June 2018. Two dark pillars indicate the prominence barbs and the horizonatl structure that connects these barbs in the prominence spine. Several vertical threads and U-shaped horns-like structures have also evident (middle panel). Rotating plasma structure indicates that the growth of the prominence is related to the tornadoes-like motions. The evolution of a helical flux rope, apparent rotating plasma, swirling motion in the top part of the prominence is appeared. 104

- 4.2 Left panel: Full Field of View (FOV) of a polar prominence observed by SDO/AIA 304 Å on 14 June 2018. The sub-region (white-box) consists of a large-scale vortex motion and a swirl-like structure. Right panel: The spatio-temporal evolution of the large-scale vortex motion. The cool plasma and photospheric vortices (large scale vortex) channeled through the prominence body. 108
- 4.3 The region of interest (ROI; white box in Figure 4.2) is displayed in the SDO/AIA (171+304) Å composite images. The composite images display the large-scale vortex motion and swirl-like structure in the upper part of the prominence. We observed the spatio-temporal evolution of swirl indicated by white arrow within the prominence. 109
- 4.4 Left panel: The region of interest (ROI) of the polar prominence legs observed by SDO/AIA 171 Å on 14 June 2018. The oppositely directed prominence legs reconnect above the PIL. Right panel: Resistive tearing mode instability initiate the formation of multiple plasmoids within the current sheet. We observed three ejecting plasmoids and a bright cusp above the prominence legs (indicated by white arrows). The oppositely directed prominence legs reconnect over polarity inversion line (PIL) and initiate the plasmoid enhanced reconnection. We assume that the PIL is lying between the prominence legs. 110

- 4.5 The normalized multi-gaussian fitted SDO/AIA 171 Å images display the FOV of polar prominence. Upper panel: Southern segment of the prominence is rising while northern segment is falling along prominence legs. The oppositely directed prominence legs reconnect above the PIL. Multiple plasmoid ejection also observed during the reconnection. These plasmoid passes through the main body of the prominence. Simultaneously, plasma downfall has also evident along the magnetic arcades. 111
- 4.6 The Full FOV of the polar prominence in AIA 171 Å emission imager to analyses the inflow, outflow and nature of EUV intensity pattern at the reconnection site. We have estimated the EUV emission from the selected white box region. Two path (slit S1 and S2) has taken across the X-point to estimate the inflow and outflow velocity. Running difference images of spacetime maps from slits "a", "b", and "c" is used to understand the magnetic field structuring and swirl motion on the top of the prominence. 114
- 4.7 Top panel: The nature of EUV emission from six different SDO/AIA filters measure from white box of Figure 4.6. The temporal variation of EUV plasma intensity (131 Å, 171 Å, 193 Å, 211 Å, 304 Å and 335 Å) has estimated to understand the thermal behaviour at the reconnection site. We have analysed the inflow (rate of the thinning of the current sheet along slit S1) and outflow velocity (ejected plasmoids along slit S2) at the reconnection site. Middle panel: The thinning of current sheet after reconnection gives the inflow speed at the X-point. Bottom panel: We track the of plasmoids in H-T diagram to measure the outflow velocity. . . 116

- 4.8 The Differential Emission Measure (DEM) of the sub-region to observe thermal structure of the plasmoids and current sheet. The DEM (Cheung et al., 2015) map the temperature bins from $\log T=5.6$ to $\log T=6.7$ shows that plasmoid are best visible temperature range $\log T=5.8-6.1$ and an elongated current sheet observed between $\log T=5.8-6.7$ 119
- 4.9 The sub-region of eruptive prominence after applying normalized multi-Gaussian filter on AIA 171 Å imager. Several sheared magnetic arcades are evident and they collapse near the reconnection site. A sheared coronal arcade thread reconnect with the southern segment of the overlying prominence. This external reconnection may trigger to erupt the prominence. The eruptive prominence consists of the post-eruption arcade and flux rope, which are indicated by arrows. 120
- 4.10 The sub-region of eruptive prominence as observe by the composite images of AIA 171+304 Å . The composite images observed the characteristics of hot magnetic arcades (AIA 171 Å, green color) and cool prominence associated plasma (AIA 304 Å, red color). A sheared coronal arcade thread reconnects with the southern segment of the overlying prominence. 121
- 4.11 The eruptive prominence observed as composite image of AIA 171+211 Å. Southern segment of the overlying prominence reconnects with the collapsing magnetic arcades and lead to the external magnetic reconnection. A thin elongated current sheet is evident, which is best visible in AIA 211 Å (red color) waveband. The prominence associated hot plasma, magnetic arcades and post-eruption arcades are best visible in the AIA 171 Å (green color) waveband. This external magnetic reconnection trigger the prominence eruption. 123

- 5.1 The running difference image sequence in the SDO/AIA 171 Å shows an eruptive prominence and its impact on the surrounding coronal field lines. The eruptive prominence disturbs the surrounding magnetic fields (middle panel; indicated by cyan arrow), which is associated with a prominence. The prominence associated field lines are forced to bring the southward segment of prominence-corona transition region (PCTR) associated coronal field lines towards the reconnection-point. The reconnection region and region of interest (ROI) is indicated by cyan box, where prominence driven southward segment of the field lines reconnect with the northward branch of the magnetic field anchored at the limb and form a temporary X-point. 130
- 5.2 Direct imaging of forced reconnection in the Sun's corona is demonstrated in this composite picture. SDO/AIA imaging observations on 3rd May 2012 at 14:10:08 UT depict the formation of a temporary X-point and a forced reconnection region (a–b). Panel 'a' is the composite image of AIA 171 Å, 304 Å showing off-limb region. Panel 'b' is a zoomed view of the ROI as shown by the dotted-white box in panel 'a'. Panel 'c' is the schematic showing the formation of X-point in the far off-limb corona. 132
- 5.3 The Differential Emission Measure (DEM) (Hannah and Kontar, 2012) maps for different temperature bins between $\text{Log } T_e = 5.0$ and $\text{Log } T_e = 7.1$ (Figure 5.3a) display that a prominence brings the overlying southward coronal magnetic field lines towards oppositely directed northward fields, and force externally the formation of an inflow at a X-point to trigger the forced reconnection. The estimated average temperature of the current sheet (Figure 5.3a, black box region) using (Hannah and Kontar, 2012) DEM technique peaks at $\text{log } T_e = 6.4$ (Figure 5.3b). 133

- 5.4 Formation of X-point, and development of forced reconnection, and inflow/outflow regions are depicted in this figure. The running difference image sequence in the SDO/AIA 171 Å filter displays the motion of the prominence corona-transition region (PCTR) interface, which brings the southward branch of the coronal magnetic field towards the temporarily evolved X-point around 13:55 UT. The forced reconnection is subsequently triggered thereafter the initial onset of externally driven inflows. Plasma inflow occurs from the north-south direction (cf., Figure 5.5b, left middle image panel at 13:48 UT), and after the reconnection, the outflow is seen in the east-west direction (cf., Figure 5.5c; left lower panel during 13:57-14:45 UT). A dynamic current sheet is also formed during the forced reconnection process. 136
- 5.5 The analyses of inflows, outflows, and nature of EUV emissions at the reconnection site are described in this figure. The temporal variation of EUV plasma emissions (304 Å, 211 Å, 193 Å, 171 Å, 131 Å) has been measured to understand qualitatively the thermal response of the reconnection site (left top panel 'a'). The slit 'S₁' and 'S₂' are analyzed for inflow and outflow plasma motions ('d') across the X-point. These slits have been selected to deduce the distance-time maps ('b, c') for estimating their speeds close to the X-point. 142

5.6	Numerical model of the forced magnetic reconnection at X-point. Using FLASH code, the X-point coronal magnetic field configuration has been reproduced on the observed spatial scales. An external disturbance is imposed on the left side of the X-point, which creates an initial average inflow of $\approx 2.56 \text{ km s}^{-1}$ triggering the reconnection of the field lines. Thereafter plasma outflows with an average speed of $\approx 12.46 \text{ km s}^{-1}$ has been generated during the reconnection in the perpendicular direction of the inflows. The simulated inflows, outflows, and the reconnection rate match with the observations of the forced reconnection. The various contours with different colours denote the velocities in units of Mm s^{-1} . The top and bottom rows display respectively inflow and outflow velocity contours around X-point as overplotted on the reconnection region. The total duration of the reconnection process is $< 10 \text{ min}$ in the simulation, which match with the duration of physical processes (inflows, reconnection heating as well as outflows) as established initially in the reconnection region between 13:55 UT and 14:05 UT (cf., Figure 5.4).	148
5.7	Left: The magnitude of V_{in}/V_{out} with the change in resistivity. Right: The magnitude of V_{in}/V_{out} with the change in the strength of the external disturbance, i.e., the initial magnitude of velocity pulse. The third order quadratic fit is also drawn on these two plots.	149
6.1	Full disk image of COR1 onboard STEREO-A & B have captured the plasma segment of an eruptive prominence at 7 June 2011. Fingers are clearly visible at 08:45 UT on this upper northward MRT unstable segment in COR1 FOV.	161

- 6.2 Sub-region of the STEREO-A & B COR1 FOV on 7th June 2011 when a MRT unstable plasma segment is in eruptive state. The temporal evolution of MRT instability in form of finger structures and horn (H) have been observed. The evolution of magnetic Rayleigh-Taylor unstable finger structures, falling blobs, and associated horn-like structure is indicated by white arrows. 164
- 6.3 Right: Partial field of view of COR1 onboard STEREO-A that shows the MRT unstable fingers in the eruptive prominence at June 7th 2011. Left: A schematic representation of a MRT unstable finger structure. The outer most finger structure has been zoomed to describe the evolution of MRT instability. 165
- 6.4 A sub-region of the top: STEREO-A & bottom: STEREO-B COR1 FOV at a time 08:45 UT on 7th June 2011 when MRT unstable plasma segment is evident. This segment lies in the upper northward part of an eruptive prominence. Denser and bright finger structures (F1-F5) are seen in this plasma segment. Darker plasma regions (B1-B5) are also visible just below of these finger structures. 167
- 6.5 Right: The temporal sequence of images which shows the erupting upper northward plasma segment in the STEREO-A/COR2 on 7 June 2011. The evolution of the mushroom-like structures in the outer corona are observed in the plasma segment. A box has drawn to indicate the region of interest. Left: The box region is displayed to identify the mushroom-like MRT unstable structures. 175

- 6.6 Right: An STEREO-A/HI1 image to observe the lower inter-planetary region on 8 June 2011 at 00:49 UT. The eruptive plasma segment is fully extended upto $20 R_{\odot}$. Left: Small scale fragmented plasma structures are clearly visible in the inter-planetary space. 176
- 6.7 Left: The evolution of MRT unstable plasma spikes in the inter-planetary space coming towards 1.0 A.U. The aligned and composite images of STEREO/COR2A (2.5 to $15 R_{\odot}$), Heliospheric Imager (HI)-1A (15 to $80 R_{\odot}$), and Heliospheric Imager (HI)-2A (upto 4 A.U.) onboard Solar TERrestrial Relations Observatory (STEREO) on 9 June 2011 at 12:09 UT, display the formation of plasma spikes impinging around the Earth. Right: A schematic showing the propagation of MRT unstable plasma spikes in the inter-planetary space. 178
- 6.8 Evolution and propagation of MRT unstable plasma spikes into the inter-planetary space towards 1 A.U. Some representative plasma spikes (S1 & S2) show their motion in and around Earth. 179
- 7.1 Evolution of a quiescent filament as observed by GONG/H α . It evolved spatially and temporally and crossed the central meridian of the Sun on 26 August 2018. The filament was lifted and partially erupted when it passes over the northward lying coronal hole. The coronal plasma channel (CPC) is associated with the filament in its southwest direction, and its location is shown in the *white box*. It appears to be empty as no traces of cool plasma in H α has been found in CPC, and it is only visible in the coronal emissions. 189

- 7.2 The full field of view (FOV) of the coronal plasma channel (CPC) that connects the two coronal holes (CH1 and CH2) observed by SDO/AIA 211 Å on 20 August 2018. The spreading coronal plasma channel (*white box*) and post-eruption arcade indicate that an eruption from this region has already occurred. The spreading coronal plasma channel further interacts with an open field line of a northward lying coronal hole (CH1) and may trigger a jet-like eruption (*lower panel*). The *white-box-region* consists of a coronal plasma channel (CPC). The impact of an eruptive coronal channel could be observed in the form of spreading of this plasma channel and formation of the post-eruption arcade as seen within the *white box region*. 191
- 7.3 The region of interest (ROI) to observe the source location of the rotating jet-like eruption as observed by SDO/AIA 211 Å and STEREO-A/EUVI-304 Å on 20 August 2018. We adopt the tie-pointing method to locate the source region of the rotating jet-like eruption in the projected plane (EUVI data). We tracked the top, middle, and bottom part of the jet-like eruption in SDO/AIA 211 Å and STEREO-A/EUVI-304 Å (*plus sign in both panels*) by the triangulation technique. The *plus sign* indicates the corresponding location of the jet-like eruption in EUVI-304 Å projected image data. The ROI for the *right panel* corresponds to Figure 7.4 and ROI for the *left panel* is plotted in Figure 7.5. 192
- 7.4 The region of interest to observe the jet-like eruption as displayed in the difference images of SDO/AIA 171 Å. The difference images show that a ring-shaped rotating jet-like eruption occurred from the northward coronal hole (CH1). 193

- 7.5 The same region of interest as Figure 7.4 in the projected plane observed by STEREO-A/EUVI-304 Å. Initiation grew and an eruption of the jet-like structure was observed in the STEREO-A/EUVI-304 Å data. 194
- 7.6 The region of interest (ROI) to observe the source location of the flux-rope eruption using SDO/AIA 211 Å and STEREO-A/EUVI-171 Å imaging data on 20 August 2018. The tie-pointing method has been used to approximate the source region of the flux-rope eruption on the solar disk (SDO/AIA 211 Å). We tracked the top, the middle, and the bottom leg of the flux rope in the STEREO-A/EUVI 171Å and SDO/AIA 211Å (*green arrow* for the upper leg, the *blue arrow* for a middle part, and *black arrow* for the bottom leg of the flux rope) by triangulation technique. A diffuse and thin flux rope lying high in the lower corona (STEREO-A/EUVI) is situated above the CPC as observed on the solar disk (SDO/AIA data). The spatial extent of left panel is same as Figure 7.7 196
- 7.7 The region of interest is displayed in the STEREO-A/EUVI (171+195) Å composite images. The composite images observed the dynamics of thin and diffuse flux rope lying high in the lower corona. The top portion of the flux rope has erupted out and the lower part is connected with the solar disk (*lower panel*). 198
- 7.8 Half FOV of STEREO-A/COR2 to observe the eruption of the CMEs. A coronal-plasma channel associated CME appeared first with its low coronal signature (upper panel). The CMEs associated with the flux rope and rotating jet-like eruption have been observed using STEREO-A/COR2. Later these CMEs merge within each other and propagate towards the outer corona in the form of a compound CME. 200

- 7.9 Formation of the merged CME in interplanetary space as observed by STEREO-A/HI1. The compound CME has larger angular width in interplanetary space and propagated towards the Earth. It crossed the lower interplanetary space on 23 August 2018. 202
- 7.10 Aligned and composite images of STEREO-A/COR2 (2.5–15 $R_{\odot}$), *Heliospheric Imager* (HI1; 15–90 $R_{\odot}$), and *Heliospheric Imager* (HI2; up to 4 AU) onboard the *Solar TErrestrial Relations Observatory* (STEREO) on 26 August 2018 at 10:09 UT, display the interaction of CME with the Earth’s magnetosphere. 203
- 7.11 The propagation of collective CMEs towards 1 AU and interaction with the Earth’s magnetosphere. The sequence of STEREO-A/HI2 images was used to observe the merged CME in interplanetary space. The observed CME moves toward 1 AU and interacts with the Earth’s magnetosphere on 26 August 2018 and creates large-scale disturbances. 205
- 7.12 Dst index after the simultaneous arrival of CMEs. The Dst index has started to decrease from positive to negative values. The Dst index decreases to -176 nT at 12:09 UT on 26 August 2018 exhibiting an intense and long duration geomagnetic storm. The average magnetic field [**B**] and the component of the magnetic field [B_x , B_y , and B_z], plasma flow speed [**V**], and proton density [N_p]. The *black-dotted vertical lines* show the arrival time of the CME at 1 AU at 12:09 UT on 25 August 2018. The z -component of the magnetic field, Dst index, and other parameters simultaneously show the disturbances in the Earth magnetosphere. 206
- 7.13 Kp index (disturbances in the Earth’s magnetic field) from 25 August to 28 August 2018. The Kp index reaches up to 7, which indicates an intense geomagnetic storm at 12:00 UT on 26 August 2018. 208