
1.10 References

- [1] T.P. Huynh, H. Haick, Autonomous Flexible Sensors for Health Monitoring, *Adv. Mater.*, **30** (2018): 1802337. <https://doi.org/10.1002/adma.201802337>.
- [2] M. Javaid, A. Haleem, S. Rab, R. Pratap Singh, R. Suman, Sensors for daily life: A review, *Sens. Int.*, **2** (2021): 100121. <https://doi.org/10.1016/j.sintl.2021.100121>.
- [3] P.A. Oberg, T. Togawa, F.A. Spelman, Sensors in medicine and health care, *John Wiley & Sons*, 2006.
- [4] R.C. Dorf, The Electrical Engineering Handbook Third Edition Sensors, Nanoscience, Biomedical Engineering, and Instruments, *Taylor & Francis*, 2006.
- [5] X. Zhang, H. Ju, J. Wang, eds. Electrochemical Sensors, Biosensors and Their Biomedical Applications, *Academic Press*, 2011.
- [6] K. Cammann, U. Lemke, A. Rohen, J. Sander, H. Wilken, B. Winter, Chemical Sensors and Biosensors-Principles and Applications, *Angew. Chem. Int. Ed. Engl.*, (5) **30** (1991): 516-539. <https://doi.org/10.1002/anie.199105161>
- [7] H.Y. Neujahr, Biosensors for environmental control, *Biotechnol. Genet. Eng. Rev.*, **1** (1984) 167–186. <https://doi.org/10.1080/02648725.1984.10647785>.
- [8] D.R. Thévenot, K. Toth, R.A. Durst, G.S. Wilson, D.R. Thévenot, Electrochemical biosensors: recommended definitions and classification, *Biosens. Bioelectron.*, (1-2) **16** (2001): 121-131.
- [9] A.P.F. Turner, I. Karube, G.S. Wilson, Biosensors: fundamentals and applications, *Oxford University Press*, 1987.
- [10] P. Supraja, S. Tripathy, S.R. Krishna Vanjari, V. Singh, S.G. Singh, Label free, electrochemical detection of atrazine using electrospun Mn₂O₃ nanofibers: Towards ultrasensitive small molecule detection, *Sens. Actuators. B. Chem.*, **285** (2019): 317–325. <https://doi.org/10.1016/j.snb.2019.01.060>.
- [11] M.M. Pereira da Silva Neves, M.B. González-García, D. Hernández-Santos, P. Fanjul-Bolado, Future trends in the market for electrochemical biosensing, *Curr. Opin. Electrochem.*, **10** (2018): 107–111. <https://doi.org/10.1016/j.coelec.2018.05.002>.
- [12] U. Chakraborty, G. Bhanjana, J. Adam, Y.K. Mishra, G. Kaur, G.R. Chaudhary, A. Kaushik, A flower-like ZnO-Ag₂O nanocomposite for label and mediator free direct sensing of dinitrotoluene, *RSC Adv.*, **10** (2020): 27764–27774. <https://doi.org/10.1039/d0ra02826f>.
- [13] N. Jadon, R. Jain, S. Sharma, K. Singh, Recent trends in electrochemical sensors for multianalyte detection – A review, *Talanta*, **161** (2016): 894–916. <https://doi.org/10.1016/j.talanta.2016.08.084>.

- [14] J.R. Askim, M. Mahmoudi, K.S. Suslick, Optical sensor arrays for chemical sensing: The optoelectronic nose, *Chem. Soc. Rev.*, **42** (2013): 8649–8682. <https://doi.org/10.1039/c3cs60179j>.
- [15] A. Bigdeli, F. Ghasemi, H. Golmohammadi, S. Abbasi-Moayed, M.A.F. Nejad, N. Fahimi-Kashani, S. Jafarinejad, M. Shahrajabian, M.R. Hormozi-Nezhad, Nanoparticle-based optical sensor arrays, *Nanoscale*, **9** (2017): 16546–16563. <https://doi.org/10.1039/c7nr03311g>.
- [16] Y. Sun, J. Ma, Y. Wang, S. Qiao, Y. Feng, Z. Li, Z. Wang, Y. Han, Z. Zhu, A Wearable Patch Sensor for Simultaneous Detection of Dopamine and Glucose in Sweat, *Analytica*, **4** (2023): 170–181. <https://doi.org/10.3390/analytica4020014>.
- [17] G. Maduraiveeran, M. Sasidharan, V. Ganesan, Electrochemical sensor and biosensor platforms based on advanced nanomaterials for biological and biomedical applications, *Biosens. Bioelectron.*, **103** (2017): 113-129. <https://doi.org/10.1016/j.bios.2017.12.031>.
- [18] J. Wang, Analytical Electrochemistry Third Edition, 2006 *John Wiley & Sons, Inc.* <https://doi.org/10.1002/0471790303>.
- [19] M. Willander, A. Tahira, Z. Ibupoto, Potentiometric Biosensors Based on Metal Oxide Nanostructures, In: Reference Module in Chemistry, Molecular Sciences and Chemical Engineering, *Elsevier*, 2017. <https://doi.org/10.1016/b978-0-12-409547-2.13482-1>.
- [20] Y. Wang, Y. Zhou, J. Sokolov, B. Rigas, K. Levon, M. Rafailovich, A potentiometric protein sensor built with surface molecular imprinting method, *Biosens. Bioelectron.*, **24** (2008): 162–166. <https://doi.org/10.1016/j.bios.2008.04.010>.
- [21] H. Kaur, S.S. Siwal, R. V. Saini, N. Singh, V.K. Thakur, Significance of an Electrochemical Sensor and Nanocomposites: Toward the Electrocatalytic Detection of Neurotransmitters and Their Importance within the Physiological System, *ACS Nanosci. Au.*, **3** (2023): 1–27. <https://doi.org/10.1021/acsnanoscienceau.2c00039>.
- [22] M.P.S. Mousavi, M.K. Abd El-Rahman, A.M. Mahmoud, R.M. Abdelsalam, P. Bühlmann, In Situ Sensing of the Neurotransmitter Acetylcholine in a Dynamic Range of 1 nM to 1 mM, *ACS Sens.*, **3** (2018): 2581–2589. <https://doi.org/10.1021/acssensors.8b00950>.
- [23] S. Elhag, Z.H. Ibupoto, O. Nur, M. Willander, Incorporating β -Cyclodextrin with ZnO nanorods: A potentiometric strategy for selectivity and detection of dopamine, *Sensors*, **14** (2014): 1654–1664. <https://doi.org/10.3390/s140101654>.
- [24] E. Lv, J. Ding, W. Qin, Potentiometric aptasensing of small molecules based on surface charge change, *Sens. Actuators. B. Chem.*, **259** (2018): 463–466. <https://doi.org/10.1016/j.snb.2017.12.067>.
- [25] K.Y. Goud, H. Teymourian, S.S. Sandhu, N. Tostado, R.K. Mishra, L.C. Moore, S.P. Harvey, J. Wang, OPAA/fluoride biosensor chip towards field detection of G-type nerve

- agents, *Sens. Actuators. B. Chem.*, **320** (2020): 128344. <https://doi.org/10.1016/j.snb.2020.128344>.
- [26] J. Ding, W. Qin, Recent advances in potentiometric biosensors, *TrAC, Trends. Anal. Chem.*, **124** (2020): 115803. <https://doi.org/10.1016/j.trac.2019.115803>.
- [27] N. Coppedè, M. Giannetto, M. Villani, V. Lucchini, E. Battista, M. Careri, A. Zappettini, Ion selective textile organic electrochemical transistor for wearable sweat monitoring, *Org. Electron.*, **78** (2020): 105579. <https://doi.org/10.1016/j.orgel.2019.105579>.
- [28] M.A. Akl, E.S.R. El-Gharkawy, N.A. El-Mahdy, S.M. El-Sheikh, S.M. Sheta, A novel nano copper complex: Potentiometry, DFT and application as a cancer prostatic biomarker for the ultrasensitive detection of human PSA, *Dalton Trans.*, **49** (2020): 15769–15778. <https://doi.org/10.1039/d0dt03318a>.
- [29] H. Karimi-Maleh, Y. Orooji, F. Karimi, M. Alizadeh, M. Baghayeri, J. Rouhi, S. Tajik, H. Beitollahi, S. Agarwal, V.K. Gupta, S. Rajendran, A. Ayati, L. Fu, A.L. Sanati, B. Tanhaei, F. Sen, M. shabani-nooshabadi, P.N. Asrami, A. Al-Othman, A critical review on the use of potentiometric based biosensors for biomarkers detection, *Biosens. Bioelectron.*, **184** (2021): 113252. <https://doi.org/10.1016/j.bios.2021.113252>.
- [30] L.C.Jr. Qlark, Monitor and control of blood and tissue oxygen tension, *ASAIO Trans.*, (1) **2** (1956):41-48.
- [31] G. Xu, W. Wang, B. Li, Z. Luo, X. Luo, A dopamine sensor based on a carbon paste electrode modified with DNA-doped poly(3,4-ethylenedioxythiophene), *Microchim. Acta*, **182** (2015): 679–685. <https://doi.org/10.1007/s00604-014-1373-8>.
- [32] E. Katz, I. Willner, Probing biomolecular interactions at conductive and semiconductive surfaces by impedance spectroscopy: Routes to impedimetric immunosensors, DNA-sensors, and enzyme biosensors, *Electroanalysis*, **15** (2003): 913–947. <https://doi.org/10.1002/elan.200390114>.
- [33] X.-Z. Yuan, C. Song, H. Wang, J. Zhang, Electrochemical impedance spectroscopy in PEM fuel cells: fundamentals and applications, *Springer London*, (420) **13** (2010). <https://doi.org/10.1007/978-1-84882-846-9>.
- [34] J.-G. Guan, Y.-Q. Miao, Q.-J. Zhang, Impedimetric Biosensors, *J. Biosci. Bioeng.*, (4) **97** (2004): 219-226. [https://doi.org/10.1016/s1389-1723\(04\)70195-4](https://doi.org/10.1016/s1389-1723(04)70195-4).
- [35] E.B. Bahadir, M.K. Sezgintürk, A review on impedimetric biosensors, *Artif. Cells Nanomed. Biotechnol.*, **44** (2016): 248–262. <https://doi.org/10.3109/21691401.2014.942456>.
- [36] M. Holzinger, A. Le Goff, S. Cosnier, Nanomaterials for biosensing applications: A review, *Front. Chem.*, **2** (2014): 63. <https://doi.org/10.3389/fchem.2014.00063>.

- [37] O.S. Kwon, H.S. Song, T.H. Park, J. Jang, Conducting Nanomaterial Sensor Using Natural Receptors, *Chem. Rev.*, **119** (2019): 36–93. <https://doi.org/10.1021/acs.chemrev.8b00159>.
- [38] B. Öndeş, F. Akpınar, M. Uygun, M. Muti, D. Aktaş Uygun, High stability potentiometric urea biosensor based on enzyme attached nanoparticles, *Microchem. J.*, **160** (2021):105667. <https://doi.org/10.1016/j.microc.2020.105667>.
- [39] C.A. Cordeiro, A. Sias, T. Koster, B.H.C. Westerink, T.I.F.H. Cremers, In vivo “real-time” monitoring of glucose in the brain with an amperometric enzyme-based biosensor based on gold coated tungsten (W-Au) microelectrodes, *Sens. Actuators. B. Chem.*, **263** (2018): 605–613. <https://doi.org/10.1016/j.snb.2018.02.116>.
- [40] L. Torsi, M. Magliulo, K. Manoli, G. Palazzo, Organic field-effect transistor sensors: A tutorial review, *Chem. Soc. Rev.*, **42** (2013): 8612–8628. <https://doi.org/10.1039/c3cs60127g>.
- [41] B.K. Bansod, T. Kumar, R. Thakur, S. Rana, I. Singh, A review on various electrochemical techniques for heavy metal ions detection with different sensing platforms, *Biosens. Bioelectron.*, **94** (2017): 443–455. <https://doi.org/10.1016/j.bios.2017.03.031>.
- [42] Y.-Bong. Hahn, Ahmad. Umar, Metal oxide nanostructures and their applications, Vol. 5. *Los Angeles, CA, USA: American Scientific Publ.*, 2010.
- [43] R. Duan, X. Fang, D. Wang, A Methylene Blue Assisted Electrochemical Sensor for Determination of Drug Resistance of Escherichia coli, *Front. Chem.*, **9** (2021): 689735. <https://doi.org/10.3389/fchem.2021.689735>.
- [44] P.S. Guin, S. Das, P.C. Mandal, Electrochemical Reduction of Quinones in Different Media: A Review, *Int. J. Electrochem.*, (1) **2011** (2011): 1–22. <https://doi.org/10.4061/2011/816202>.
- [45] H. Liu, J. Deng, Amperometric glucose sensor using tetrathiafulvalene in Nafion gel as electron shuttle, *Anal. Chim. Acta*, (1-3) **300** (1995): 65-70. [https://doi.org/10.1016/0003-2670\(94\)00397-5](https://doi.org/10.1016/0003-2670(94)00397-5)
- [46] D. Astruc, Why is Ferrocene so Exceptional?, *Eur. J. Inorg. Chem.*, (1) **2017** (2017): 6–29. <https://doi.org/10.1002/ejic.201600983>.
- [47] T.J. Kealy, P.L. Pauson, A New Type of Organo-Iron Compound, *Nature*, **168** (1951): 1039–1040. <https://doi.org/10.1038/1681039b0>.
- [48] H. Beitollahi, M.A. Khalilzadeh, S. Tajik, M. Safaei, K. Zhang, H.W. Jang, M. Shokouhimehr, Recent Advances in Applications of Voltammetric Sensors Modified with Ferrocene and Its Derivatives, *ACS Omega*, **5** (2020): 2049–2059. <https://doi.org/10.1021/acsomega.9b03788>.

- [49] B. Fabre, Ferrocene-terminated monolayers covalently bound to hydrogen-terminated silicon surfaces. Toward the development of charge storage and communication devices, *Acc. Chem. Res.*, **43** (2010): 1509–1518. <https://doi.org/10.1021/ar100085q>.
- [50] J. Wu, L. Wang, H. Yu, Zain-ul-Abdin, R.U. Khan, M. Haroon, Ferrocene-based redox-responsive polymer gels: Synthesis, structures and applications, *J. Organomet. Chem.*, **828** (2017): 38–51. <https://doi.org/10.1016/j.jorganchem.2016.10.041>.
- [51] F.A. Larik, A. Saeed, T.A. Fattah, U. Muqadar, P.A. Channar, Recent advances in the synthesis, biological activities and various applications of ferrocene derivatives, *Appl. Organomet. Chem.*, **31** (2017): e3664. <https://doi.org/10.1002/aoc.3664>.
- [52] M.H.R. Borges, B.E. Nagay, R.C. Costa, J.G.S. Souza, M.T. Mathew, V.A.R. Barão, Recent advances of polypyrrole conducting polymer film for biomedical application: Toward a viable platform for cell-microbial interactions, *Adv. Colloid Interface Sci.*, **314** (2023): 102860. <https://doi.org/10.1016/j.cis.2023.102860>.
- [53] S.A. Hira, M. Yusuf, D. Annas, S. Nagappan, S. Song, S. Park, K.H. Park, Recent Advances on Conducting Polymer-Supported Nanocomposites for Nonenzymatic Electrochemical Sensing, *Ind. Eng. Chem. Res.*, **60** (2021): 13425–13437. <https://doi.org/10.1021/acs.iecr.1c02043>.
- [54] J.M. George, A. Antony, B. Mathew, Metal oxide nanoparticles in electrochemical sensing and biosensing: a review, *Microchim. Acta*, **185** (2018): 1-26. <https://doi.org/10.1007/s00604-018-2894-3>.
- [55] X. Luo, A. Morrin, A.J. Killard, M.R. Smyth, Application of nanoparticles in electrochemical sensors and biosensors, *Electroanalysis*, **18** (2006): 319–326. <https://doi.org/10.1002/elan.200503415>.
- [56] K. Singh, K.K. Maurya, M. Malviya, Review of Electrochemical Sensors and Biosensors Based on First-Row Transition Metals, Their Oxides, and Noble Metals Nanoparticles, *J. Anal. Test.*, (2) **8** (2023): 1-17. <https://doi.org/10.1007/s41664-023-00292-w>.
- [57] A.C. Power, B. Gorey, S. Chandra, J. Chapman, Carbon nanomaterials and their application to electrochemical sensors: A review, *Nanotechnol. Rev.*, **7** (2018): 19–41. <https://doi.org/10.1515/ntrev-2017-0160>.
- [58] C. Yang, M.E. Denno, P. Pyakurel, B.J. Venton, Recent trends in carbon nanomaterial-based electrochemical sensors for biomolecules: A review, *Anal. Chim. Acta.*, **887** (2015): 17–37. <https://doi.org/10.1016/j.aca.2015.05.049>.
- [59] A. Muhulet, F. Miculescu, S.I. Voicu, F. Schütt, V.K. Thakur, Y.K. Mishra, Fundamentals and scopes of doped carbon nanotubes towards energy and biosensing applications, *Mater. Today Energy*, **9** (2018): 154–186. <https://doi.org/10.1016/j.mtener.2018.05.002>.

- [60] H. Kaur, S.S. Siwal, G. Chauhan, A.K. Saini, A. Kumari, V.K. Thakur, Recent advances in electrochemical-based sensors amplified with carbon-based nanomaterials (CNMs) for sensing pharmaceutical and food pollutants, *Chemosphere*, **304** (2022): 135182. <https://doi.org/10.1016/j.chemosphere.2022.135182>.
- [61] A.D. da Silva, W.J. Paschoalino, J.P. V. Damasceno, L.T. Kubota, Structure, Properties, and Electrochemical Sensing Applications of Graphene-Based Materials, *Chem. Electro. Chem.*, **7** (2020): 4508–4525. <https://doi.org/10.1002/celec.202001168>.
- [62] Y. Shao, J. Wang, H. Wu, J. Liu, I.A. Aksay, Y. Lin, Graphene based electrochemical sensors and biosensors: A review, *Electroanalysis*, **22** (2010): 1027–1036. <https://doi.org/10.1002/elan.200900571>.
- [63] S. Wu, Q. He, C. Tan, Y. Wang, H. Zhang, Graphene-based electrochemical sensors, *Small*, **9** (2013): 1160–1172. <https://doi.org/10.1002/sml.201202896>.
- [64] S. Nigar, Z. Zhou, H. Wang, M. Imtiaz, Modulating the electronic and magnetic properties of graphene, *RSC Adv.*, **7** (2017): 51546–51580. <https://doi.org/10.1039/c7ra08917a>.
- [65] Z. Zhao, W. Zheng, W. Yu, B. Long, Electrical conductivity of poly(vinylidene fluoride)/carbon nanotube composites with a spherical substructure, *Carbon*, **47** (2009): 2118–2120. <https://doi.org/10.1016/j.carbon.2009.03.043>.
- [66] T. Thomas, R.J. Mascarenhas, O.J. D'Souza, S. Detriche, Z. Mekhalif, P. Martis, Pristine multi-walled carbon nanotubes/SDS modified carbon paste electrode as an amperometric sensor for epinephrine, *Talanta*, **125** (2014): 352–360. <https://doi.org/10.1016/j.talanta.2014.03.027>.
- [67] R.K.F. Lee, B.J. Cox, J.M. Hill, The geometric structure of single-walled nanotubes, *Nanoscale*, **2** (2010): 859–872. <https://doi.org/10.1039/b9nr00433e>.
- [68] O. Lupan, F. Schütt, V. Postica, D. Smazna, Y.K. Mishra, R. Adelung, Sensing performances of pure and hybridized carbon nanotubes-ZnO nanowire networks: A detailed study, *Sci. Rep.*, **7** (2017): 1-12. <https://doi.org/10.1038/s41598-017-14544-0>.
- [69] F.R. Baptista, S.A. Belhout, S. Giordani, S.J. Quinn, Recent developments in carbon nanomaterial sensors, *Chem. Soc. Rev.*, **44** (2015): 4433–4453. <https://doi.org/10.1039/c4cs00379a>.
- [70] A. Shahat, N.Y. Elamin, W. Abd El-Fattah, Spectrophotometric and Fluorometric Methods for the Determination of Fe(III) Ions in Water and Pharmaceutical Samples, *ACS Omega*, **7** (2022): 1288–1298. <https://doi.org/10.1021/acsomega.1c05899>.
- [71] A.H. Kamel, A.E.G.E. Amr, A.A. Almehizia, E.A. Elsayed, G.O. Moustafa, Low-cost potentiometric paper-based analytical device based on newly synthesized macrocyclic pyrido-pentapeptide derivatives as novel ionophores for point-of-care copper(ii) determination, *RSC Adv.*, **11** (2021): 27174–27182. <https://doi.org/10.1039/d1ra04712d>.

- [72] S. Jampasa, W. Khamcharoen, S. Traipop, W. Jesadabundit, T. Ozer, O. Chailapakul, Recent advances on nanomaterial-modified film-electrode-based sensors: Approach to clinical purpose, *Curr. Opin. Electrochem.*, **42** (2023): 101420. <https://doi.org/10.1016/j.coelec.2023.101420>.
- [73] S. Kempahanumakkagari, A. Deep, K.H. Kim, S. Kumar Kailasa, H.O. Yoon, Nanomaterial-based electrochemical sensors for arsenic - A review, *Biosens. Bioelectron.*, **95** (2017): 106–116. <https://doi.org/10.1016/j.bios.2017.04.013>.
- [74] H. Cheng, X. Wang, H. Wei, Ratiometric Electrochemical Sensor for Effective and Reliable Detection of Ascorbic Acid in Living Brains, *Anal. Chem.*, **87** (2015): 8889–8895. <https://doi.org/10.1021/acs.analchem.5b02014>.
- [75] A.M. Pisoschi, A. Pop, A.I. Serban, C. Fafaneata, Electrochemical methods for ascorbic acid determination, *Electrochim. Acta*, **121** (2014): 443–460. <https://doi.org/10.1016/j.electacta.2013.12.127>.
- [76] A. Joshi, W. Schuhmann, T.C. Nagaiah, Mesoporous nitrogen containing carbon materials for the simultaneous detection of ascorbic acid, dopamine and uric acid, *Sens. Actuators. B. Chem.*, **230** (2016): 544–555. <https://doi.org/10.1016/j.snb.2016.02.050>.
- [77] F. Palmisano, P.G. Zambonin, Ascorbic acid interferences in hydrogen peroxide detecting biosensors based on electrochemically immobilized enzymes, *Anal. Chem.*, **65** (1993): 2690–2692. <https://doi.org/10.1021/ac00067a024>
- [78] P.R. Roy, T. Okajima, T. Ohsaka, Simultaneous electrochemical detection of uric acid and ascorbic acid at a poly(N,N-dimethylaniline) film-coated GC electrode, *J. Electroanal. Chem.*, **561** (2004): 75–82. <https://doi.org/10.1016/j.jelechem.2003.07.003>.
- [79] N.S. Anuar, W.J. Basirun, M. Shalauddin, S. Akhter, A dopamine electrochemical sensor based on a platinum-silver graphene nanocomposite modified electrode, *RSC Adv.*, **10** (2020): 17336–17344. <https://doi.org/10.1039/c9ra11056a>.
- [80] M. Lakshmanakumar, N. Nesakumar, A.J. Kulandaisamy, J.B.B. Rayappan, Principles and recent developments in optical and electrochemical sensing of dopamine: A comprehensive review, *Measurement*, **183** (2021): 109873. <https://doi.org/10.1016/j.measurement.2021.109873>.
- [81] X. Liu, J. Liu, Biosensors and sensors for dopamine detection, *View*, **2** (2021): 20200102. <https://doi.org/10.1002/viw.20200102>.
- [82] P.K. Kannan, S.A. Moshkalev, C.S. Rout, Highly sensitive and selective electrochemical dopamine sensing properties of multilayer graphene nanobelts, *Nanotechnol.*, **27** (2016): 075504. <https://doi.org/10.1088/0957-4484/27/7/075504>.
- [83] G. Balkourani, A. Brouzgou, P. Tsiakaras, A review on recent advancements in electrochemical detection of dopamine using carbonaceous nanomaterials, *Carbon*, **213** (2023): 118281. <https://doi.org/10.1016/j.carbon.2023.118281>.

- [84] D. Yu, Y. Zeng, Y. Qi, T. Zhou, G. Shi, A novel electrochemical sensor for determination of dopamine based on AuNPs@SiO₂ core-shell imprinted composite, *Biosens. Bioelectron.*, **38** (2012): 270–277. <https://doi.org/10.1016/j.bios.2012.05.045>.
- [85] S. Han, T. Du, H. Jiang, X. Wang, Synergistic effect of pyrroloquinoline quinone and graphene nano-interface for facile fabrication of sensitive NADH biosensor, *Biosens. Bioelectron.*, **89** (2017): 422–429. <https://doi.org/10.1016/j.bios.2016.04.092>.
- [86] G. Weng, X. Zhao, J. Zhao, J. Li, J. Zhu, J. Zhao, Nanoplasmonic sensing of NADH by inhibiting the oxidative etching of gold nanorods, *Sens. Actuators. B. Chem.*, **299** (2019): 126982. <https://doi.org/10.1016/j.snb.2019.126982>.
- [87] P. Manusha, S. Yadav, J. Satija, S. Senthilkumar, Designing electrochemical NADH sensor using silver nanoparticles/phenothiazine nanohybrid and investigation on the shape dependent sensing behavior, *Sens. Actuators. B. Chem.*, **347** (2021): 130649. <https://doi.org/10.1016/j.snb.2021.130649>.
- [88] O.M. Istrate, L. Rotariu, V.E. Marinescu, C. Bala, NADH sensing platform based on electrochemically generated reduced graphene oxide-gold nanoparticles composite stabilized with poly(allylamine hydrochloride), *Sens. Actuators. B. Chem.*, **223** (2016): 697–704. <https://doi.org/10.1016/j.snb.2015.09.142>.
- [89] K.S. Prasad, J.C. Chen, C. Ay, J.M. Zen, Mediatorless catalytic oxidation of NADH at a disposable electrochemical sensor, *Sens. Actuators. B. Chem.*, **123** (2007): 715–719. <https://doi.org/10.1016/j.snb.2006.10.012>.
- [90] J. Zhu, X. Chen, W. Yang, A high performance electrochemical sensor for NADH based on graphite nanosheet modified electrode, *Sens. Actuators. B. Chem.*, **150** (2010): 564–568. <https://doi.org/10.1016/j.snb.2010.08.039>.

2.2 References

- [1] Martin. Ermrich, Detlef. Opper, PANalytical (Almelo), XRD for the analyst: getting acquainted with the principles, *PANalytical*, 2013.
- [2] J. Epp, 4 - X-ray diffraction (XRD) techniques for materials characterization, in: G. Hübschen, I. Altpeter, R. Tschuncky, H.-G. Herrmann (Eds.), *Materials Characterization Using Nondestructive Evaluation (NDE) Methods*, Woodhead Publishing, (2016) 81–124. [https://doi.org/https://doi.org/10.1016/b978-0-08-100040-3.00004-3](https://doi.org/10.1016/b978-0-08-100040-3.00004-3).
- [3] C. Hammond, *The basics of crystallography and diffraction*, Vol. **21**. Oxford University Press, USA, 2015.
- [4] C.F. Holder, R.E. Schaak, Tutorial on Powder X-ray Diffraction for Characterizing Nanoscale Materials, *ACS Nano*, **13** (2019) 7359–7365. <https://doi.org/10.1021/acsnano.9b05157>.
- [5] J.I. Goldstein, D.E. Newbury, J.R. Michael, N.W.M. Ritchie, J.H.J. Scott, D.C. Joy, *Scanning Electron Microscopy and X-Ray Microanalysis*, Springer New York, (2017). <https://doi.org/10.1007/978-1-4939-6676-9>.
- [6] K. Akhtar, S.A. Khan, S.B. Khan, A.M. Asiri, Scanning electron microscopy: Principle and applications in nanomaterials characterization, in: *Handbook of Materials Characterization*, Springer International Publishing, (2018) 113–145. https://doi.org/10.1007/978-3-319-92955-2_4.
- [7] B.J. Inkson, Scanning electron microscopy (SEM) and transmission electron microscopy (TEM) for materials characterization, *Materials Characterization Using Nondestructive Evaluation (NDE) Methods*, Woodhead publishing, (2016) 17–43. <https://doi.org/10.1016/b978-0-08-100040-3.00002-x>.
- [8] T. Walther, *Transmission Electron Microscopy of Nanostructures*, Microscopy Methods in Nanomaterials Characterization, Elsevier, (2017) 105–134. <https://doi.org/10.1016/B978-0-323-46141-2.00004-3>.
- [9] R. Bellotti, G.B. Picotto, L. Ribotta, AFM Measurements and Tip Characterization of Nanoparticles with Different Shapes, *Nanomanufacturing Metrol.*, **5** (2022) 127–138. <https://doi.org/10.1007/s41871-022-00125-x>.
- [10] D.N.G. Krishna, J. Philip, Review on surface-characterization applications of X-ray photoelectron spectroscopy (XPS): Recent developments and challenges, *Appl. Surf. Sci. Adv.*, **12** (2022). <https://doi.org/10.1016/j.apsadv.2022.100332>.

- [11] M.A. Isaacs, J. Davies-Jones, P.R. Davies, S. Guan, R. Lee, D.J. Morgan, R. Palgrave, Advanced XPS characterization: XPS-based multi-technique analyses for comprehensive understanding of functional materials, *Mater. Chem. Front.*, **5** (2021) 7931–7963. <https://doi.org/10.1039/d1qm00969a>.
- [12] E. Mazzotta, S. Rella, A. Turco, C. Malitesta, XPS in development of chemical sensors, *RSC Adv.*, **5** (2015) 83164–83186. <https://doi.org/10.1039/c5ra14139g>.
- [13] E. Bakker, M. Telting-Diaz, Electrochemical sensors, *Anal. Chem.*, **74** (2002) 2781–2800. <https://doi.org/10.1021/ac0202278>.
- [14] O. Gilbert, B.E. Kumara Swamy, U. Chandra, B.S. Sherigara, Simultaneous detection of dopamine and ascorbic acid using polyglycine modified carbon paste electrode: A cyclic voltammetric study, *J. Electroanal. Chem.*, **636** (2009) 80–85. <https://doi.org/10.1016/j.jelechem.2009.09.016>.
- [15] F. Harnisch, S. Freguia, A basic tutorial on cyclic voltammetry for the investigation of electroactive microbial biofilms, *Chem. Asian J.*, **7** (2012) 466–475. <https://doi.org/10.1002/asia.201100740>.
- [16] F. Marken, A. Neudeck, A.M. Bond, Cyclic voltammetry, in: *Electroanalytical Methods: Guide to Experiments and Applications*, Springer Berlin Heidelberg, (2010) 57–106. https://doi.org/10.1007/978-3-642-02915-8_4.
- [17] B. Kashyap, R. Kumar, A novel multi-set differential pulse voltammetry technique for improving precision in electrochemical sensing, *Biosens. Bioelectron.*, **216** (2022). <https://doi.org/10.1016/j.bios.2022.114628>.
- [18] S. Tajik, Z. Dourandish, P.M. Jahani, I. Sheikhshoae, H. Beitollahi, M. Shahedi Asl, H.W. Jang, M. Shokouhimehr, Recent developments in voltammetric and amperometric sensors for cysteine detection, *RSC Adv.*, **11** (2021) 5411–5425. <https://doi.org/10.1039/d0ra07614g>.
- [19] F. Lisdat, D. Schäfer, The use of electrochemical impedance spectroscopy for biosensing, *Anal. Bioanal. Chem.*, **391** (2008) 1555–1567. <https://doi.org/10.1007/s00216-008-1970-7>.
- [20] A.C. Lazanas, M.I. Prodromidis, Electrochemical Impedance Spectroscopy-A Tutorial, *ACS Meas. Sci. Au*, **3** (2023) 162–193. <https://doi.org/10.1021/acsmasuresciau.2c00070>.

3.6 References

- [1] M.J. Tierney, H.O.L. Kim, Electrochemical Gas Sensor with Extremely Fast Response Times, *Anal. Chem.*, **65** (1993): 3435–3440. <https://doi.org/10.1021/ac00071a017>.
- [2] N.G. Tsierkezos, U. Ritter, Electrochemical impedance spectroscopy and cyclic voltammetry of ferrocene in acetonitrile/acetone system, *J. Appl. Electrochem.*, **40** (2010): 409–417. <https://doi.org/10.1007/s10800-009-0011-3>.
- [3] N. Changsan, S. Chairam, P. Jarujamrus, M. Amatatongchai, Sensitive electrochemical sensor based on gold nanoparticles assembled ferrocene-functionalised graphene oxide modified glassy carbon electrode for simultaneous determination of dopamine and acetaminophen, *Adv. Nat. Sci.: Nanosci. Nanotechnol.*, **13** (2022): 015012. <https://doi.org/10.1088/2043-6262/ac5d44>.
- [4] P.C. Pandey, B.C. Upadhyay, Role of Palladium in the Redox Electrochemistry of Ferrocene Monocarboxylic Acid Encapsulated Within ORMOSIL Networks, *Molecules*, **10** (2005): 728–739. <https://doi.org/10.3390/10060728>.
- [5] H. Beitollahi, M.A. Khalilzadeh, S. Tajik, M. Safaei, K. Zhang, H.W. Jang, M. Shokouhimehr, Recent Advances in Applications of Voltammetric Sensors Modified with Ferrocene and Its Derivatives, *ACS Omega*, **5** (2020): 2049–2059. <https://doi.org/10.1021/acsomega.9b03788>.
- [6] L. Fernández, H. Carrero, Electrochemical evaluation of ferrocene carboxylic acids confined on surfactant–clay modified glassy carbon electrodes: oxidation of ascorbic acid and uric acid, *Electrochim. Acta*, **50** (2005): 1233–1240. <https://doi.org/10.1016/J.electacta.2004.08.016>.
- [7] G. huang Wu, Y. fang Wu, X. wei Liu, M. cong Rong, X. mei Chen, X. Chen, An electrochemical ascorbic acid sensor based on palladium nanoparticles supported on graphene oxide, *Anal. Chim. Acta*, **745** (2012): 33–37. <https://doi.org/10.1016/j.aca.2012.07.034>.
- [8] S. Qiu, S. Gao, L. Xie, H. Chen, Q. Liu, Z. Lin, B. Qiu, G. Chen, An ultra-sensitive electrochemical sensor for ascorbic acid based on click chemistry, *Analyst*, **136** (2011): 3962–3966. <https://doi.org/10.1039/c1an15316a>.
- [9] D. Sun, Y. Zhang, F. Wang, K. Wu, J. Chen, Y. Zhou, Electrochemical sensor for simultaneous detection of ascorbic acid, uric acid and xanthine based on the surface enhancement effect of mesoporous silica, *Sens. Actuators B: Chem.*, **2** (2009): 641–645. <https://doi.org/10.1016/j.snb.2009.07.043>.

- [10] P.R. Lima, P.R.B. de Miranda, A.B. de Oliveira, M.O.F. Goulart, L.T. Kubota, Modified Carbon Paste Electrode for Kinetic Investigation and Simultaneous Determination of Ascorbic and Uric Acids, *Electroanalysis*, **21** (2009): 2311. <https://doi.org/10.1002/elan.200904694>.
- [11] P.C. Pandey, A.K. Pandey, G. Pandey, Functionalized alkoxy silane mediated controlled synthesis of noble metal nanoparticles dispersible in aqueous and non-aqueous medium, *J. Nanosci. Nanotechnol.*, **14** (2014): 6606–6613. <https://doi.org/10.1166/jnn.2014.9378>.
- [12] P.C. Pandey, R. Singh, A.K. Pandey, Tetrahydrofuran hydroperoxide and 3-aminopropyltrimethoxysilane mediated controlled synthesis of Pd, Pd-Au, Au-Pd nanoparticles: Role of Palladium nanoparticles on the redox electrochemistry of ferrocene monocarboxylic acid, *Electrochim. Acta*, **138** (2014): 163–173. <https://doi.org/10.1016/j.electacta.2014.06.101>.
- [13] P.C. Pandey, R. Singh, Y. Pandey, Controlled synthesis of functional Ag, Ag–Au/Au–Ag nanoparticles and their Prussian blue nanocomposites for bioanalytical applications, *RSC Adv.*, **5** (2015): 49671–49679. <https://doi.org/10.1039/c5ra06251a>.
- [14] P.C. Pandey, M.D. Mitra, S. Shukla, R.J. Narayan, Organotrialkoxysilane-Functionalized Noble Metal Monometallic, Bimetallic, and Trimetallic Nanoparticle Mediated Non-Enzymatic Sensing of Glucose by Resonance Rayleigh Scattering, *Biosensors*, **11** (2021): 122. <https://doi.org/10.3390/bios11040122>.
- [15] L. Fabbri, The ferrocenium/ferrocene couple: a versatile redox switch, *ChemTexts*, **6** (2020): 1–20. <https://doi.org/10.1007/s40828-020-00119-6>.
- [16] J.B. Raoof, R. Ojani, F. Chekin, Electrochemical Analysis of D-Penicillamine Using a Carbon Paste Electrode Modified with Ferrocene Carboxylic Acid, *Electroanalysis*, **19** (2007): 1883–1889. <https://doi.org/10.1002/elan.200703947>.
- [17] P.C. Pandey, S. Upadhyay, N.K. Shukla, S. Sharma, Studies on the electrochemical performance of glucose biosensor based on ferrocene encapsulated ORMOSIL and glucose oxidase modified graphite paste electrode, *Biosens. Bioelectron.*, **18** (2003): 1257–1268. [https://doi.org/10.1016/s0956-5663\(03\)00075-7](https://doi.org/10.1016/s0956-5663(03)00075-7).
- [18] P.C. Pandey, V. Singh, P.C. Pandey, V. Singh, Electrochemical polymerization of aniline over tetracyanoquinodimethane encapsulated ormosil matrix: application in the electrocatalytic oxidation of ascorbic acid and acetylthiocholine, *Analyst*, **136** (2011): 1472. <https://doi.org/10.1039/c0an00491j>.

- [19] P.C. Pandey, S. Shukla, Solvent dependent fabrication of bifunctional nanoparticles and nanostructured thin films by self assembly of organosilanes, *J. Solgel. Sci. Technol.*, **86** (2018): 650–663. <https://doi.org/10.1007/s10971-018-4686-y>.
- [20] Chitra Singh, K. Singh, Prem.C. Pandey, Synthesis and Properties of Organotrialkoxysilane Functionalized Palladium–Cobalt Heterogeneous Catalysts for Oxygen Evolution Reaction, *Russ. J. Electrochem.*, **59** (2023): 604–615. <https://doi.org/10.1134/s1023193523080074>.
- [21] Y.C. Weng, Y.L. Hsiao, Comparison of Pt and Ni foil electrodes for amperometric sensing of ascorbic acid, *J. Electroanal. Chem.*, **651** (2011): 160–165. <https://doi.org/10.1016/j.jelechem.2010.11.031>.
- [22] L. Yang, D. Liu, J. Huang, T. You, Simultaneous determination of dopamine, ascorbic acid and uric acid at electrochemically reduced graphene oxide modified electrode, *Sens. Actuators B: Chem.*, **193** (2014): 166–172. <https://doi.org/10.1016/j.snb.2013.11.104>.
- [23] A. Singh, A. Sharma, A. Ahmed, S. Arya, Highly selective and efficient electrochemical sensing of ascorbic acid via CuO/rGO nanocomposites deposited on conductive fabric, *Appl. Phys. A: Mater. Sci. Process*, **128** (2022): 262. <https://doi.org/10.1007/s00339-022-05436-w>.
- [24] J. Yan, S. Liu, Z. Zhang, G. He, P. Zhou, H. Liang, L. Tian, X. Zhou, H. Jiang, Simultaneous electrochemical detection of ascorbic acid, dopamine and uric acid based on graphene anchored with Pd-Pt nanoparticles, *Colloids Surf. B: Biointerfaces*, **111** (2013): 392–397. <https://doi.org/10.1016/j.colsurfb.2013.06.030>.
- [25] L. Fu, A. Wang, G. Lai, W. Su, F. Malherbe, J. Yu, C. te Lin, A. Yu, Defects regulating of graphene ink for electrochemical determination of ascorbic acid, dopamine and uric acid, *Talanta*, **180** (2018): 248–253. <https://doi.org/10.1016/j.talanta.2017.12.058>.
- [26] S. Feng, L. Yu, M. Yan, J. Ye, J. Huang, X. Yang, Holey nitrogen-doped graphene aerogel for simultaneously electrochemical determination of ascorbic acid, dopamine and uric acid, *Talanta*, **224** (2021): 121851. <https://doi.org/10.1016/j.talanta.2020.121851>.
- [27] Y. Peng, D. Zhang, C. Zhang, Simultaneous voltammetric determination of ascorbic acid and uric acid using a seven-hole carbon nanotube paste multielectrode array, *Anal. Methods*, **6** (2014): 8965–8972. <https://doi.org/10.1039/c4ay01029a>.

$\mu\text{A}/\text{mM}$, the Co-NC/Pd /Dmfc electrodes demonstrate good analytical performance for the detection of NADH. This electrochemical sensor has been demonstrated to be extremely sensitive and accurate for detecting NADH in working electrolyte. One of the main elements of the upcoming sensor generation is reliable and producible catalytic electrodes for NADH oxidation.

4.5 References

- [1] M. Shivakumar, S. Manjunatha, M.S. Dharmaprakash, S.B. M, Biological activity of PdNPs derived from hemicellulose via microwave assisted green synthesis, *Curr. Res. Green Sustain. Chem.*, **4** (2021): 100150. <https://doi.org/10.1016/j.crgsc.2021.100150>.
- [2] G. Weng, X. Zhao, J. Zhao, J. Li, J. Zhu, J. Zhao, Nanoplasmonic sensing of NADH by inhibiting the oxidative etching of gold nanorods, *Sens. Actuators B: Chem.*, **299** (2019): 126982. <https://doi.org/10.1016/j.snb.2019.126982>.
- [3] S. Immanuel, R. Sivasubramanian, Electrochemical kinetic investigation of NADH oxidation on Prussian blue-mediated chemically reduced graphene oxide nanosheets, *J. Phys. Chem. Solids*, **161** (2022): 110471. <https://doi.org/10.1016/j.jpics.2021.110471>.
- [4] L. Wang, J. Zhang, B. Kim, J. Peng, S.N. Berry, Y. Ni, D. Su, J. Lee, L. Yuan, Y.-T. Chang, Boronic Acid: A Bio-Inspired strategy to increase the sensitivity and selectivity of fluorescent NADH probe, *J. Am. Chem. Soc.*, **138**, no. 33 (2016): 10394-10397. <https://doi.org/10.1021/jacs.6b05810>.
- [5] P. Manusha, S. Senthilkumar, Development of a ferrocene-tethered ionic liquid modified electrode for non-enzymatic electrochemical sensing of NADH, *J. Mater. Sci.: Mater. Electron.*, **33** (2022): 8576–8585. <https://doi.org/10.1007/s10854-021-06576-0>.
- [6] N.F. Atta, S.A. Abdel Gawad, E.H. El-Ads, A.R.M. El-Gohary, A. Galal, A new strategy for NADH sensing using ionic liquid crystals-carbon nanotubes/nano-magnetite composite platform, *Sens. Actuators B: Chem.*, **251** (2017): 65–73. <https://doi.org/10.1016/j.snb.2017.05.026>.

- [7] Y. Mie, Y. Yasutake, M. Ikegami, T. Tamura, Anodized gold surface enables mediator-free and low-overpotential electrochemical oxidation of NADH: A facile method for the development of an NAD⁺-dependent enzyme biosensor, *Sens. Actuators B: Chem.*, **288** (2019): 512–518. <https://doi.org/10.1016/j.snb.2019.03.039>.
- [8] W. Xie, A. Xu, E.S. Yeung, Determination of NAD⁺ and NADH in a single cell under hydrogen peroxide stress by capillary electrophoresis, *Anal. Chem.*, **81** (2009): 1280–1284. <https://doi.org/10.1021/ac802249m>.
- [9] M. Chu, Z. Bai, D. Zhu, W. Chen, G. Yang, J. Xin, H. Ma, H. Pang, L. Tan, X. Wang, A β -nicotinamide adenine dinucleotide electrochemical sensor based on polyoxometalate built by the combination of electrodeposition and self-assembly, *J. Electroanal. Chem.*, **907** (2022): 116083. <https://doi.org/10.1016/j.jelechem.2022.116083>.
- [10] M. Wang, X. Kan, Multilayer sensing platform: gold nanoparticles/prussian blue decorated graphite paper for NADH and H₂O₂ detection, *Analyst*, **143** (2018): 5278–5284. <https://doi.org/10.1039/c8an01502c>.
- [11] K.K. Maurya, K. Singh, M. Malviya, Effect of palladium and its nanogeometry on the redox electrochemistry of tetracyanoquinodimethane modified electrode; application in electrochemical sensing of ascorbic acid, *J. Appl. Electrochem.*, (9) **53** (2023): 1831–1842. <https://doi.org/10.1007/s10800-023-01878-z>.
- [12] A.S. Agnihotri, A. Varghese, N. M, Transition metal oxides in electrochemical and bio sensing: A state-of-art review, *Appl. Surf. Sci. Adv.*, **4** (2021): 100072. <https://doi.org/10.1016/j.apsadv.2021.100072>.
- [13] J.M. George, A. Antony, B. Mathew, Metal oxide nanoparticles in electrochemical sensing and biosensing: a review, *Microchim. Acta*, **185** (2018): 1–26. <https://doi.org/10.1007/s00604-018-2894-3>.
- [14] P. Liang, H. Yu, B. Guntupalli, Y. Xiao, Paper-based device for rapid visualization of NADH based on dissolution of gold nanoparticles, *ACS Appl. Mater. Interfaces*, **7** (2015): 15023–15030. <https://doi.org/10.1021/acsami.5b04104>.
- [15] H. Jaegfeldt, T. Kuwana, G. Johansson, Electrochemical Stability of Catechols with a Pyrene Side Chain Strongly Adsorbed on Graphite Electrodes for Catalytic Oxidation of Dihyronicotinamide Adenine Dinucleotide, *J. Am. Chem. Soc.*, (7) **105** (1983): 1805–1814. <https://doi.org/10.1021/ja00345a021>

- [16] G.P. Keeley, A. O'Neill, M. Holzinger, S. Cosnier, J.N. Coleman, G.S. Duesberg, DMF-exfoliated graphene for electrochemical NADH detection, *Phys. Chem. Chem. Phys.*, **13** (2011): 7747–7750. <https://doi.org/10.1039/c1cp20060g>.
- [17] C. Shan, H. Yang, D. Han, Q. Zhang, A. Ivaska, L. Niu, Electrochemical determination of NADH and ethanol based on ionic liquid-functionalized graphene, *Biosens. Bioelectron.*, **25** (2010): 1504–1508. <https://doi.org/10.1016/j.bios.2009.11.009>.
- [18] Y. Liu, R. Landick, S. Raman, A Regulatory NADH/NAD⁺ Redox Biosensor for Bacteria, *ACS Synth. Biol.*, **8** (2019): 264–273. <https://doi.org/10.1021/acssynbio.8b00485>.
- [19] R.D. Nagarajan, P. Murugan, A.K. Sundramoorthy, Selective Electrochemical Sensing of NADH and NAD⁺ Using Graphene/Tungstate Nanocomposite Modified Electrode, *ChemistrySelect*, **5** (2020): 14643–14651. <https://doi.org/10.1002/slct.202003554>.
- [20] A. Koyappayil, H.T. Kim, M.H. Lee, An efficient and rapid synthesis route to highly fluorescent copper microspheres for the selective and sensitive excitation wavelength-dependent dual-mode sensing of NADH, *Sens. Actuators B: Chem.*, **327** (2021): 128887. <https://doi.org/10.1016/j.snb.2020.128887>.
- [21] P. Manusha, S. Yadav, J. Satija, S. Senthilkumar, Designing electrochemical NADH sensor using silver nanoparticles/phenothiazine nano hybrid and investigation on the shape dependent sensing behavior, *Sens. Actuators B: Chem.*, **347** (2021):130649. <https://doi.org/10.1016/j.snb.2021.130649>.
- [22] L. Zhu, R. Yang, X. Jiang, D. Yang, Amperometric determination of NADH at a Nile blue/ordered mesoporous carbon composite electrode, *Electrochem. Commun.*, **11** (2009): 530–533. <https://doi.org/10.1016/j.elecom.2008.12.045>.
- [23] S. Shahrokhian, R. Salimian, S. Rastgar, Pd-Au nanoparticle decorated carbon nanotube as a sensing layer on the surface of glassy carbon electrode for electrochemical determination of ceftazidime, *Mater. Sci. Eng. C*, **34** (2014): 318–325. <https://doi.org/10.1016/j.msec.2013.09.014>.
- [24] G. Maduraiveeran, R. Ramaraj, Silver nanoparticles embedded in amine-functionalized silicate sol-gel network assembly for sensing cysteine, adenosine and NADH, *J. Nanoparticle Res.*, **13** (2011): 4267–4276. <https://doi.org/10.1007/s11051-011-0372-5>.
- [25] X.H. Pham, C.A. Li, K.N. Han, B.C. Huynh-Nguyen, T.H. Le, E. Ko, J.H. Kim, G.H. Seong, Electrochemical detection of nitrite using urchin-like palladium nanostructures on carbon nanotube thin film electrodes, *Sens. Actuators B: Chem.*, **193** (2014): 815–822. <https://doi.org/10.1016/j.snb.2013.12.034>.

- [26] S.H. Lim, J. Wei, J. Lin, Q. Li, J. KuaYou, A glucose biosensor based on electrodeposition of palladium nanoparticles and glucose oxidase onto Nafion-solubilized carbon nanotube electrode, *Biosens. Bioelectron.*, **20** (2005): 2341–2346. <https://doi.org/10.1016/j.bios.2004.08.005>.
- [27] B. Singh, N. Bhardwaj, V.K. Jain, V. Bhatia, Palladium nanoparticles decorated electrostatically functionalized MWCNTs as a non-enzymatic glucose sensor, *Sens. Actuators A: Phys.*, **220** (2014): 126–133. <https://doi.org/10.1016/j.sna.2014.09.030>.
- [28] B. Wang, K. Chen, G. Wang, X. Liu, H. Wang, J. Bai, A multidimensional and hierarchical carbon-confined cobalt phosphide nanocomposite as an advanced anode for lithium and sodium storage, *Nanoscale*, **11** (2019): 968–985. <https://doi.org/10.1039/c8nr07076h>.
- [29] G. Zhang, L. Liu, Q. Zhu, X. Kong, Chemoselective Hydrogenation of Nitroarenes by an Efficient Co@NC/AC Catalyst, *Catal. Letters*, **153** (2023): 1536–1542. <https://doi.org/10.1007/s10562-022-04085-1>.
- [30] K. Murugesan, T. Senthamarai, A.S. Alshammari, R.M. Altamimi, C. Kreyenschulte, M.M. Pohl, H. Lund, R. V. Jagadeesh, M. Beller, Cobalt-Nanoparticles Catalyzed Efficient and Selective Hydrogenation of Aromatic Hydrocarbons, *ACS Catal.*, **9** (2019): 8581–8591. <https://doi.org/10.1021/acscatal.9b02193>.
- [31] W. Yi, Z. Li, C. Dong, H.W. Li, J. Li, Electrochemical detection of chloramphenicol using palladium nanoparticles decorated reduced graphene oxide, *Microchem. J.* **148** (2019): 774–783. <https://doi.org/10.1016/j.microc.2019.05.049>.
- [32] R. Zhang, M. Tahir, S. Ding, M.A. Qadeer, H. Li, Q.X. Zeng, R. Gao, L. Wang, X. Zhang, L. Pan, J.J. Zou, Promotion of Nitrogen Reserve and Electronic Regulation in Bamboo-like Carbon Tubules by Cobalt Nanoparticles for Highly Efficient ORR, *ACS Appl. Energy Mater.*, **3** (2020): 2323–2330. <https://doi.org/10.1021/acsaem.9b01617>.
- [33] K. Zhu, C. Jin, Z. Klencsár, A.S. Ganeshraja, J. Wang, Cobalt-iron oxide, alloy and nitride: Synthesis, characterization and application in catalytic peroxydisulfate activation for orange II degradation, *Catalysts*, **7** (2017): 138. <https://doi.org/10.3390/catal7050138>.
- [34] S. Liang, C. Liang, High-density cobalt nanoparticles encapsulated with nitrogen-doped carbon nanoshells as a bifunctional catalyst for rechargeable zinc-air battery, *Materials*, **12** (2019): 243. <https://doi.org/10.3390/ma12020243>.

- [35] A. Chen, C. Ostrom, Palladium-Based Nanomaterials: Synthesis and Electrochemical Applications, *Chem. Rev.*, **115** (2015): 11999–12044. <https://doi.org/10.1021/acs.chemrev.5b00324>.
- [36] Y. Xiong, Y. Xia, Shape-controlled synthesis of metal nanostructures: The case of palladium, *Adv. Mater.*, **19** (2007): 3385–3391. <https://doi.org/10.1002/adma.200701301>.
- [37] K. Białaś, D. Moschou, F. Marken, P. Estrela, Electrochemical sensors based on metal nanoparticles with biocatalytic activity, *Microchim. Acta*, **189** (2022): 172. <https://doi.org/10.1007/s00604-022-05252-2>.
- [38] B. Fall, D.D. Sall, M. Hémadi, A.K.D. Diaw, M. Fall, H. Randriamahazaka, S. Thomas, Highly efficient non-enzymatic electrochemical glucose sensor based on carbon nanotubes functionalized by molybdenum disulfide and decorated with nickel nanoparticles (GCE/CNT/MoS₂/NiNPs), *Sens. Actuators Rep.* **5** (2023): 100136. <https://doi.org/10.1016/j.snr.2022.100136>.
- [39] H. Elsaywy, B.M. Thamer, A. Sedky, M.H. El-Newehy, Facile two-step synthesis of nickel nanoparticles supported on 3D porous carbon frameworks as an effective electrocatalyst for urea and methanol oxidation, *Mater. Chem. Phys.*, **297** (2023): 127361. <https://doi.org/10.1016/j.matchemphys.2023.127361>.
- [40] S. Chen, K. Shang, X. Gao, X. Wang, The development of NAD⁺-dependent dehydrogenase screen-printed biosensor based on enzyme and nanoporous gold co-catalytic strategy, *Biosens. Bioelectron.*, **211** (2022): 114376. <https://doi.org/10.1016/j.bios.2022.114376>.
- [41] T. Anusha, K.S. Bhavani, J. V. Shanmukha Kumar, P.K. Brahman, R.Y.A. Hassan, Fabrication of electrochemical immunosensor based on GCN-β-CD/Au nanocomposite for the monitoring of vitamin D deficiency, *Bioelectrochemistry*, **143** (2022): 107935. <https://doi.org/10.1016/j.bioelechem.2021.107935>.
- [42] H.S. Magar, R.Y.A. Hassan, A. Mulchandani, Electrochemical impedance spectroscopy (EIS): Principles, construction, and biosensing applications, *Sensors*, **21** (2021): 6578. <https://doi.org/10.3390/s21196578>.
- [43] R. Aghajari, A. Azadbakht, Amplified detection of streptomycin using aptamer-conjugated palladium nanoparticles decorated on chitosan-carbon nanotube, *Anal. Biochem.*, **547** (2018): 57–65. <https://doi.org/10.1016/j.ab.2018.02.005>.

- [44] C.E. Banks, R.G. Compton, Exploring the electrocatalytic sites of carbon nanotubes for NADH detection: An edge plane pyrolytic graphite electrode study, *Analyst*, **130** (2005): 1232–1239. <https://doi.org/10.1039/b508702c>.
- [45] P. Jain, B. Chakma, S. Patra, P. Goswami, Hairpin stabilized fluorescent silver nanoclusters for quantitative detection of NAD⁺ and monitoring NAD⁺/NADH based enzymatic reactions, *Anal. Chim. Acta*, **956** (2017): 48–56. <https://doi.org/10.1016/j.aca.2016.12.030>.
- [46] L. Meng, A.P.F. Turner, W.C. Mak, Positively-charged hierarchical PEDOT interface with enhanced electrode kinetics for NADH-based biosensors, *Biosens. Bioelectron.*, **120** (2018): 115–121. <https://doi.org/10.1016/j.bios.2018.08.017>.
- [47] M. Sahin, E. Ayrançi, Electrooxidation of NADH on Modified Screen-Printed Electrodes: Effects of Conducting Polymer and Nanomaterials, *Electrochim. Acta*, **166** (2015): 261–270. <https://doi.org/10.1016/j.electacta.2015.03.030>.
- [48] A.R. Marlinda, S. Sagadevan, N. Yusoff, A. Pandikumar, N.M. Huang, O. Akbarzadeh, M.R. Johan, Gold nanorods-coated reduced graphene oxide as a modified electrode for the electrochemical sensory detection of NADH, *J. Alloys Compd.*, **847** (2020): 156552. <https://doi.org/10.1016/j.jallcom.2020.156552>.
- [49] S.N. Prashanth, K.C. Ramesh, J. Seetharamappa, Electrochemical Oxidation of an Immunosuppressant, Mycophenolate Mofetil, and Its Assay in Pharmaceutical Formulations, *Int. J. Electrochem.*, **2011** (2011): 1–7. <https://doi.org/10.4061/2011/193041>.
- [50] M. Elanchezian, K. Theyagarajan, V.K. Ponnusamy, K. Thenmozhi, S. Senthilkumar, Porous graphene oxide based disposable non-enzymatic electrochemical sensor for the determination of nicotinamide adenine dinucleotide, *Micro Nano Eng.*, **15** (2022): 100133. <https://doi.org/10.1016/j.mne.2022.100133>.

5.5 References

- [1] D. Kim, S. Lee, Y. Piao, Electrochemical determination of dopamine and acetaminophen using activated graphene-Nafion modified glassy carbon electrode, *J. Electroanal. Chem.*, **794** (2017): 221–228. <https://doi.org/10.1016/j.jelechem.2017.04.018>.
- [2] M. Wei, Y. Qiao, H. Zhao, J. Liang, T. Li, Y. Luo, S. Lu, X. Shi, W. Lu, X. Sun, Electrochemical non-enzymatic glucose sensors: recent progress and perspectives, *Chem. Commun.*, **56** (2020): 14553–14569. <https://doi.org/10.1039/d0cc05650b>.
- [3] J. Feng, Q. Li, J. Cai, T. Yang, J. Chen, X. Hou, Electrochemical detection mechanism of dopamine and uric acid on titanium nitride-reduced graphene oxide composite with and without ascorbic acid, *Sens. Actuators B: Chem.*, **298** (2019): 126872. <https://doi.org/10.1016/j.snb.2019.126872>.
- [4] R. Sha, N. Vishnu, S. Badhulika, MoS₂ based ultra-low-cost, flexible, non-enzymatic and non-invasive electrochemical sensor for highly selective detection of Uric acid in human urine samples, *Sens. Actuators B: Chem.*, **279** (2019): 53–60. <https://doi.org/10.1016/j.snb.2018.09.106>.
- [5] L.Q. Xie, Y.H. Zhang, F. Gao, Q.A. Wu, P.Y. Xu, S.S. Wang, N.N. Gao, Q.X. Wang, A highly sensitive dopamine sensor based on a polyaniline/reduced graphene oxide/Nafion nanocomposite, *Chin. Chem. Lett.* **28** (2017): 41–48. <https://doi.org/10.1016/j.ccllet.2016.05.015>.
- [6] T. Kokulnathan, T.J. Wang, E.A. Kumar, N. Duraisamy, An-Ting Lee, An electrochemical platform based on yttrium oxide/boron nitride nanocomposite for the detection of dopamine, *Sens. Actuators B: Chem.*, **349** (2021): 130787. <https://doi.org/10.1016/j.snb.2021.130787>.
- [7] C. Wang, J. Chen, L. Zhang, Y. Yang, M. Huang, C. Chen, C. Li, Y. Xie, P. Zhao, J. Fei, An ultra-sensitive dopamine photoelectrochemical sensing platform based on two-dimensional Zn carbon nanosheets, hollow Cu₂O and CdTe QDs composite films, *Carbon*, **198** (2022) 101–109. <https://doi.org/10.1016/j.carbon.2022.07.007>.
- [8] B. Patella, A. Sortino, F. Mazzara, G. Aiello, G. Drago, C. Torino, A. Vilasi, A. O’Riordan, R. Inguanta, Electrochemical detection of dopamine with negligible interference from ascorbic and uric acid by means of reduced graphene oxide and metals-NPs based electrodes, *Anal. Chim. Acta*, **1187** (2021): 339124. <https://doi.org/10.1016/j.aca.2021.339124>.
- [9] L.O. Orzari, M.H.M.T. Assumpção, J. Nandenha, A.O. Neto, L.H.M. Junior, M. Bergamini, B.C. Janegitz, Pd, Ag and Bi carbon-supported electrocatalysts as electrochemical multifunctional materials for ethanol oxidation and dopamine determination, *Electrochim. Acta*, **428** (2022): 140932. <https://doi.org/10.1016/j.electacta.2022.140932>.

- [10] H. Ouyang, W. Li, Y. Long, Carbon-doped h-BN for the enhanced electrochemical determination of dopamine, *Electrochim. Acta*, **369** (2021): 137682. <https://doi.org/10.1016/j.electacta.2020.137682>.
- [11] A. Arroquia, I. Acosta, M.P.G. Armada, Self-assembled gold decorated polydopamine nanospheres as electrochemical sensor for simultaneous determination of ascorbic acid, dopamine, uric acid and tryptophan, *Mater. Sci. Eng. C*, **109** (2020): 110610. <https://doi.org/10.1016/j.msec.2019.110602>.
- [12] A. Savk, B. Özdil, B. Demirkan, M.S. Nas, M.H. Calimli, M.H. Alma, Inamuddin, A.M. Asiri, F. Şen, Multiwalled carbon nanotube-based nanosensor for ultrasensitive detection of uric acid, dopamine, and ascorbic acid, *Mater. Sci. Eng. C*, **99** (2019): 248–254. <https://doi.org/10.1016/j.msec.2019.01.113>.
- [13] Y.H. Chang, P.M. Woi, Y. Alias, The selective electrochemical detection of dopamine in the presence of ascorbic acid and uric acid using electro-polymerised- β -cyclodextrin incorporated f-MWCNTs/polyaniline modified glassy carbon electrode, *Microchem. J.*, **148** (2019): 322–330. <https://doi.org/10.1016/j.microc.2019.04.081>.
- [14] K. Kaewket, C. Karuwan, S. Sonsupap, S. Maensiri, K. Ngamchuea, Anti-Fouling Effects of Carbon Nanofiber in Electrochemical Sensing of Phenolic Compounds, *J. Electrochem. Soc.*, **168** (2021): 067501. <https://doi.org/10.1149/1945-7111/ac0358>.
- [15] B.L. Hanssen, S. Siraj, D.K.Y. Wong, Recent strategies to minimise fouling in electrochemical detection systems, *Rev. Anal. Chem.*, **35** (2016): 1–28. <https://doi.org/10.1515/revac-2015-0008>.
- [16] E. Peltola, S. Sainio, K.B. Holt, T. Palomäki, J. Koskinen, T. Laurila, Electrochemical Fouling of Dopamine and Recovery of Carbon Electrodes, *Anal. Chem.*, **90** (2018): 1408–1416. <https://doi.org/10.1021/acs.analchem.7b04793>.
- [17] B. Liu, H. Guo, L. Sun, Z. Pan, L. Peng, M. Wang, N. Wu, Y. Chen, X. Wei, W. Yang, Electrochemical sensor based on covalent organic frameworks/MWCNT for simultaneous detection of catechol and hydroquinone, *Colloids Surf. A: Physicochem. Eng. Asp.*, **639** (2022): 128335. <https://doi.org/10.1016/j.colsurfa.2022.128335>.
- [18] J. Cheng, X. Wang, T. Nie, L. Yin, S. Wang, Y. Zhao, H. Wu, H. Mei, A novel electrochemical sensing platform for detection of dopamine based on gold nanobipyramid/multi-walled carbon nanotube hybrids, *Anal. Bioanal. Chem.*, **412** (2020): 2433–2441. <https://doi.org/10.1007/s00216-020-02455-5>.
- [19] G. Li, B. Yuan, L. Zhao, W. Gao, C. Xu, G. Liu, Fouling-resistant electrode for electrochemical sensing based on covalent-organic frameworks TpPA-1 dispersed carbon nanotubes, *Talanta*, **267** (2024): 125162. <https://doi.org/10.1016/j.talanta.2023.125162>.
- [20] F. Heshmatpour, R. Abazari, Formation of dispersed palladium-nickel bimetallic nanoparticles in microemulsions: Synthesis, characterization, and their use as efficient heterogeneous recyclable catalysts for the amination reactions of aryl chlorides under

- mild conditions, *RSC Adv.*, **4** (2014): 55815–55826. <https://doi.org/10.1039/c4ra06958g>.
- [21] R.K. Shervedani, M. Karevan, A. Amini, Prickly nickel nanowires grown on Cu substrate as a supersensitive enzyme-free electrochemical glucose sensor, *Sens. Actuators B: Chem.* **204** (2014): 783–790. <https://doi.org/10.1016/j.snb.2014.08.033>.
- [22] J.A.S.B. Cardoso, B. Šljukić, E. Kayhan, C.A.C. Sequeira, D.M.F. Santos, Palladium-nickel on tin oxide-carbon composite supports for electrocatalytic hydrogen evolution, *Catal. Today*, **357** (2020): 302–310. <https://doi.org/10.1016/j.cattod.2019.05.056>.
- [23] Y. Zhang, Y. Liao, G. Shi, W. Wang, B. Su, Preparation, characterization, and catalytic performance of Pd-Ni/AC bimetallic nano-catalysts, *Green Process. Synth.*, **9** (2020): 760–769. <https://doi.org/10.1515/gps-2020-0071>.
- [24] S. Liu, S.K. Tam, K.M. Ng, Dual-reductant synthesis of nickel nanoparticles for use in screen-printing conductive paste, *J. Nanoparticle Res.*, **23** (2021): 78. <https://doi.org/10.1007/s11051-021-05191-8>.
- [25] M.R. Ahghari, V. Soltaninejad, A. Maleki, Synthesis of nickel nanoparticles by a green and convenient method as a magnetic mirror with antibacterial activities, *Sci. Rep.*, **10** (2020): 12627. <https://doi.org/10.1038/s41598-020-69679-4>.
- [26] M.J. Huhn-Ibarra, M.I. Loría-Bastarrachea, S. Duarte-Aranda, A. de J. Montes-Luna, J. Ortiz-Espinoza, M.O. González-Díaz, M. Aguilar-Vega, PPSU dual layer hollow fiber mixed matrix membranes with functionalized MWCNT for enhanced antifouling, salt and dye rejection in water treatment, *J. Appl. Polym. Sci.* **139** (2022): e53203. <https://doi.org/10.1002/app.53203>.
- [27] H. Cheng, N. Yang, G. Liu, Y. Ge, J. Huang, Q. Yun, Y. Du, C.J. Sun, B. Chen, J. Liu, H. Zhang, Ligand-Exchange-Induced Amorphization of Pd Nanomaterials for Highly Efficient Electrocatalytic Hydrogen Evolution Reaction, *Adv. Mater.*, **32** (2020): 1902964. <https://doi.org/10.1002/adma.201902964>.
- [28] X.L. Cai, C.H. Liu, J. Liu, Y. Lu, Y.N. Zhong, K.Q. Nie, J.L. Xu, X. Gao, X.H. Sun, S.D. Wang, Synergistic effects in CNTs-Pd Au/Pt trimetallic nanoparticles with high electrocatalytic activity and stability, *Nanomicro. Lett.*, **9** (2017): 1–10. <https://doi.org/10.1007/s40820-017-0149-1>.
- [29] M.G. Hosseini, F. Hosseinzadeh, P. Zardari, O. Mermer, Pd-Ni nanoparticle supported on reduced graphene oxide and multi-walled carbon nanotubes as electrocatalyst for oxygen reduction reaction, *Fuller. Nanotub. Carbon Nanostructures*, **26** (2018): 675–687. <https://doi.org/10.1080/1536383X.2018.1465049>.
- [30] F. Gulbagca, A. Aygün, M. Gülcan, S. Ozdemir, S. Gonca, F. Şen, Green synthesis of palladium nanoparticles: Preparation, characterization, and investigation of antioxidant, antimicrobial, anticancer, and DNA cleavage activities, *Appl. Organomet. Chem.*, **35** (2021): e6272. <https://doi.org/10.1002/aoc.6272>.

- [31] G. Dutta, F.C.B. Fernandes, P. Estrela, D. Moschou, P.R. Bueno, Impact of surface roughness on the self-assembling of molecular films onto gold electrodes for label-free biosensing applications, *Electrochim. Acta*, **378** (2021): 138137. <https://doi.org/10.1016/j.electacta.2021.138137>.
- [32] A.J. Bard, L.R. Faulkner, H.S. White, *Electrochemical methods: fundamentals and applications*, John Wiley & Sons, 2022.
- [33] X. Zheng, X. Zhou, X. Ji, R. Lin, W. Lin, Simultaneous determination of ascorbic acid, dopamine and uric acid using poly(4-aminobutyric acid) modified glassy carbon electrode, *Sens. Actuators B: Chem.*, **178** (2013): 359–365. <https://doi.org/10.1016/j.snb.2012.12.115>.
- [34] G. Venkata Prasad, S.J. Jang, Y. Chandra Sekhar, T. Madhusudana Reddy, L. Subramanyam Sarma, H.B. Kim, T. Hyun Kim, Fine-tuning of Pd–CeO₂/rGO nanocomposite: A facile synergetic strategy for effective electrochemical detection of dopamine in pharmaceutical and biological samples, *J. Electroanal. Chem.*, **941** (2023): 117544. <https://doi.org/10.1016/j.jelechem.2023.117544>.
- [35] M. Dai, Q. Zhu, D. Han, L. Niu, Z. Wang, Sensitive and selective electrochemical sensor for the detection of dopamine by using AuPd@Fe₂O₃ nanoparticles as catalyst, *Adv. Sensor. Energy Mater.*, **2** (2023): 100048. <https://doi.org/10.1016/j.asems.2023.100048>.
- [36] K.J. Huang, J.Z. Zhang, Y.J. Liu, L.L. Wang, Novel electrochemical sensing platform based on molybdenum disulfide nanosheets-polyaniline composites and Au nanoparticles, *Sens. Actuators B: Chem.*, **194** (2014): 303–310. <https://doi.org/10.1016/j.snb.2013.12.106>.
- [37] K.K. Maurya, K. Singh, M. Malviya, Fabrication of dopamine sensor based on a carbon paste electrode modified with tetracyanoquinodimethane and cobalt tungstate nanoparticles, *J. Mater. Sci.: Mater. Electron.*, **35** (2024): 13. <https://doi.org/10.1007/s10854-023-11721-y>.
- [38] T.K. Aparna, R. Sivasubramanian, M.A. Dar, One-pot synthesis of Au-Cu₂O/rGO nanocomposite based electrochemical sensor for selective and simultaneous detection of dopamine and uric acid, *J. Alloys Compd.*, **741** (2018): 1130–1141. <https://doi.org/10.1016/j.jallcom.2018.01.205>.
- [39] S. Immanuel, T.K. Aparna, R. Sivasubramanian, A facile preparation of Au—SiO₂ nanocomposite for simultaneous electrochemical detection of dopamine and uric acid, *Surf. Interfaces*, **14** (2019): 82–91. <https://doi.org/10.1016/j.surfin.2018.11.010>.
- [40] C. Zhang, J. Ren, J. Zhou, M. Cui, N. Li, B. Han, Q. Chen, Facile fabrication of a 3,4,9,10-perylene tetracarboxylic acid functionalized graphene-multiwalled carbon nanotube-gold nanoparticle nanocomposite for highly sensitive and selective electrochemical detection of dopamine, *Analyst*, **143** (2018): 3075–3084. <https://doi.org/10.1039/c8an00559a>.

- [41] A. Xie, H. Wang, J. Lin, J. Pan, M. Li, J. Wang, S. Jiang, S. Luo, 3D rGO/MWCNTs-loaded bimetallic-organic gel derived $ZrFeO_x$ as an electrochemical sensor for simultaneous detection of dopamine and paracetamol, *J. Alloys Compd.*, **938** (2023): 168647. <https://doi.org/10.1016/j.jallcom.2022.168647>.
- [42] S.E. Elugoke, O.E. Fayemi, A.S. Adekunle, B.B. Mamba, T.T.I. Nkambule, E.E. Ebenso, Electrochemical sensor for the detection of dopamine using carbon quantum dots/copper oxide nanocomposite modified electrode, *FlatChem*, **33** (2022): 100372. <https://doi.org/10.1016/j.flatc.2022.100372>.
- [43] S. Ponnada, D.B. Gorle, M.S. Kiai, S. Rajagopal, R.K. Sharma, A. Nowduri, A facile, cost-effective, rapid, single-step synthesis of Ag-Cu decorated ZnO nanoflower-like composites (NFLCs) for electrochemical sensing of dopamine, *Mater. Adv.*, **2** (2021): 5986–5996. <https://doi.org/10.1039/d1ma00319d>.
- [44] Y.J. Yang, W. Li, CTAB functionalized graphene oxide/multiwalled carbon nanotube composite modified electrode for the simultaneous determination of ascorbic acid, dopamine, uric acid and nitrite, *Biosens. Bioelectron.*, **56** (2014): 300–306. <https://doi.org/10.1016/j.bios.2014.01.037>.
- [45] C. Yang, X. Sun, C. Zhang, M. Liu, Green synthesis of Co-Ni hollow spheres for its electrochemical detection of dopamine, *J. Nanoparticle Res.*, **22** (2020): 55. <https://doi.org/10.1007/s11051-020-4775-z>.
- [46] K. Reddaiah, T.M. Reddy, K. Mallikarjuna, G. Narasimha, Electrochemical detection of dopamine at poly (solochrome cyanine)/Pd nanoparticles doped modified carbon paste electrode and simultaneous resolution in the presence of ascorbic acid and uric acid: A voltammetric method, *Anal. Methods*, **5** (2013): 5627–5636. <https://doi.org/10.1039/c3ay41039k>.